# **Explanation of Key Trends - Sulfur Dioxide**

# **Obligations**

Under the terms of the UN ECE Geneva Convention on Long-range Transboundary Air Pollution Control (CLTRAP, 1979), the Federal Republic of Germany was obliged by the UN ECE Helsinki Protocol to reduce its annual sulphur emissions by at least 30% by 1993, as compared to 1980 levels. In 1993, the SO<sub>2</sub> emissions were 2.9 Mt, compared to approximately 7.5 Mt in 1980. This represents a reduction of 61%. The second UN ECE protocol on the reduction of sulphur emissions obliged Germany to reduce SO<sub>2</sub> emissions to 1,300kt by 2000, and to 990kt by 2005. The targets set for 2000 & 2005 have already been achieved by 1998.

More recently, Germany has made a commitment under the multicomponent protocol to further minimise  $SO_2$  emissions. Since 2010, it is no longer permissible to exceed a National Emission Ceiling of 550kt  $SO_2$  for Germany as whole. The revised Gothenburg Protocol and the revised NEC Directive both define emission reduction targets relative to a 2005 base year, mandating 21% (2020) and 58% (2030) reductions respectively.

Further details can be found in Chapter 9 - Projections and Chapter 11 - Adjustments and Emission Ceiling Exceedance.

## Main drivers


By far the largest proportion of SO<sub>2</sub> is produced by the oxidation of the sulphur contained in the fuels used in combustion processes, Fuel Combustion (NFR 1.A) with close to 96% of total SO<sub>2</sub> emissions in 1990 and a 96% reduction between 1990-2019. In 1990, the biggest source of emissions therein is Public Electricity and Heat Production (NFR 1.A.1.a) with about 60% of the emissions from Fuel Combustion (NFR 1.A). Other sources substantially influencing the SO<sub>2</sub> emission trend are Manufacturing Industries and Construction (NFR 1.A.2) and Other Sectors (NFR 1.A.4, including commercial/institutional and residential sources), each adding about one fifth of 1990 emissions from Fuel Combustion (NFR 1.A). All of these sub-categories show a reduction >95 per cent between 1990 and 2019, due to stricter regulations of West Germany that applied to the New German Länder after the German Reunification and changed the fuel mix from sulphur-rich solid fuels to liquid and gaseous fuels.

### **SO<sub>2</sub> Emissions 1990-2019**

Total Emissions (kt)														Trend: latest compared to	
1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	1990	last years
2.854	2.198	1.907	1.642	1.470	1.444	1.435	1.434	1.389	1.362	1.338	1.289	1.207	1.133	-60,3%	<b>1</b>

#### Sulfur Dioxide / Schwefeldioxid

Emissions per Sector / Sektorale Emissionen



Quelle-German Emission Inventory (08.01.2021)

 $NO_{\times}$  trend by sector