

1.A.5.b i - Military Ground Vehicles and Vehicles

Short description

In sub-category *1.A.5.b i - Other, Mobile (including Military)* emissions from military ground-vehicles and mobile machinery are reported.

Method	AD	EF	Key Category
T1, T2	NS	CS, D	see superordinate chapter

Method

Activity data

Basically, all fuel consumption in military vehicles is included in the primary activity data provided by the National Energy Balances (NEB) (AGEB, 2019).

As the NEB does not provide specific data for military use, the following additional sources are used:

For the years as of 1995, the official mineral-oil data of the Federal Republic of Germany (Amtliche Mineralölstatistik der Bundesrepublik Deutschland), prepared by the Federal Office of Economics and Export Control (BAFA), are used (BAFA, 2019)¹⁾. Provided in units of [1,000 t], these amounts have to be converted into [TJ] on the basis of the relevant net calorific values given by²⁾.

As the official mineral-oil data does not distinguish into fossil and biofuels but does provide amounts for inland deliveries of total diesel and gasoline fuels, no data on the consumption of biodiesel and bioethanol is available directly at the moment. Therefore, activity data for biofuels used in military vehicles are calculated by applying Germany's official annual biofuel shares to the named total deliveries (see also: info on EF).

Table 1: Annual fuel deliveries to the military for ground-vehicles and machinery, in terajoules

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Diesel Oil	15,037	8,001	1,364	3,366	990	622	972	681	683	580	578	415	279	
Biodiesel	0	0	0	74	64	41	63	39	41	31	30	22	16	
Gasoline	21,508	9,800	7,477	6,857	4,862	4,696	4,175	4,092	3,695	3,342	3,009	2,502	2,341	
Biogasoline	0	0	0	47	188	192	185	175	161	145	131	107	105	
Σ 1.A.5.b i	36,545	17,801	8,841	10,343	6,103	5,551	5,395	4,988	4,580	4,099	3,748	3,046	2,741	

[gallery size="medium" : 1A5bi_AD.png : 1A5bi_AD_bio.png gallery](#)

Emission factors

Table 2: Annual country-specific emission factors¹, in kg/TJ

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

DIESEL FUELS																		
NH ₃	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00
NM VOC	316	274	274	274	274	274	274	274	274	274	274	274	274	274	274	274	274	274
NO _x	1,195	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360	1,360
SO _x	125	60.5	14.0	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37	0.37
PM ²	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
BC ³	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0	53.0
CO	515	350	350	350	350	350	350	350	350	350	350	350	350	350	350	350	350	350
GASOLINE FUELS																		
NH ₃	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00
NM VOC	594	373	373	373	373	373	373	373	373	373	373	373	373	373	373	373	373	373
NO _x	682	725	725	725	725	725	725	725	725	725	725	725	725	725	725	725	725	725
SO _x	11.8	8.30	3.20	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40
PM ²	3.63	3.55	3.13	2.66	2.66	2.51	2.39	2.27	2.14	2.09	2.03	1.97	1.91	1.91	1.91	1.91	1.91	1.91
BC ³	0.44	0.43	0.38	0.32	0.32	0.30	0.29	0.27	0.26	0.25	0.24	0.24	0.23	0.23	0.23	0.23	0.23	0.23
CO	4,199	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010	4,010
TSP ⁴	2.46	0.82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Pb ⁴	1.54	0.52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

¹ Due to lack of better information: similar EF are applied for fossil fuels and biofuels.

² EF(PM_{2.5}) also applied for PM₁₀ and TSP (assumption: > 99% of TSP from diesel oil combustion consists of PM_{2.5})

³ EF(BC) estimated from tier1 default f-BC values provided in ³⁾, chapter 1.A.3.b, table 3-11 for gasoline passenger cars (f-BC: 0.12) and diesel heavy duty vehicles (f-BC: 0.53)

⁴ from leaded gasoline (until 1997), based upon country-specific emission factors from ⁴⁾

NOTE: With respect to the emission factors applied for particulate matter, given the circumstances during test-bench measurements, condensables are most likely included at least partly.[footnote](#) During test-bench measurements, temperatures are likely to be significantly higher than under real-world conditions, thus reducing condensation. On the contrary, smaller dilution (higher number of primary particles acting as condensation germs) together with higher pressures increase the likeliness of condensation. So over-all condensables are very likely to occur but different to real-world conditions. [footnote](#)

For information on the **emission factors for heavy-metal and POP exhaust emissions**, please refer to [Appendix 2.3 - Heavy Metal \(HM\) exhaust emissions from mobile sources](#)] and [Appendix 2.4 - Persistent Organic Pollutant \(POP\) exhaust emissions from mobile sources](#)].

Discussion of emission trends

This sub-category is **not considered separately in the key category analysis**.

Due to the application of very several tier1 emission factors, most emission trends reported for this sub-category only reflect the trend in fuel deliveries. Therefore, the fuel-consumption dependent trends in emission estimates are only influenced by the annual fuel mix.

[gallery size="medium" : 1A5bi_EM_NH3.png : 1A5bi_EM_NOx.png gallery](#)

Here, for **sulphur dioxide**, this consumption-based falling trend is intensified by the impact of fuel-sulphur legislation.

[gallery size="medium" : 1A5bi_EM_SO2.png gallery](#)

Over-all **particulate matter** emissions are by far dominated by emissions from diesel oil combustion with the falling trend basically following the decline in fuel consumption. Here, until 1997, the emission values reported for **total suspended particles (TSP)** are slightly higher than those reported for PM_{2.5}, and PM₁₀, due to the additional TSP emissions from leaded gasoline that was banned in 1997.

[gallery size="medium" : 1A5bi_EM_PM.png gallery](#)

Recalculations

activity data

Table: Revised activity data, in terajoules

	2009	2010	2011	2012	2013	2014	2015	2016	2017				
1.A.5.b i TOTAL													
Submission 2021	6,119	6,103	5,551	5,395	4,988	4,580	4,099	3,748	3,046				
Submission 2020	6,119	6,103	5,551	5,395	4,988	4,580	4,102	3,752	3,155				
absolute change	-0.09	0.02	0.11	0.0	0.0	0.0	-3.6	-4.2	-109				
relative change	-0.001%	0.0004%	0.002%	0.00%	0.00%	0.00%	-0.09%	-0.11%	-3.46%				
DIESEL OIL													
Submission 2021	1,003	990	622	972	681	683	580	578	415				
Submission 2020	1,003	990	622	972	681	683	583	582	421				
absolute change	0.02	0.02	0.00	0.00	0.00	0.00	-3.50	-4.10	-5.76				
relative change	0.002%	0.002%	0.00%	0.00%	0.00%	0.00%	-0.60%	-0.70%	-1.37%				
BIODIESEL													
Submission 2021	69	64	41	63	39	41	31	30	22				
Submission 2020	69	64	41	63	39	41	32	31	22				
absolute change	0.00	0.00	0.00	0.00	0.00	0.00	-0.19	-0.22	-0.31				
relative change	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	-0.60%	-0.70%	-1.37%				
GASOLINE													
Submission 2021	4,907	4,862	4,696	4,175	4,092	3,695	3,342	3,009	2,502				
Submission 2020	4,907	4,862	4,695	4,175	4,092	3,695	3,342	3,009	2,605				
absolute change	-0.11	0.00	0.10	0.00	0.00	0.00	0.08	0.07	-103				
relative change	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	-3.96%				
BIOGASOLINE													
Submission 2021	140	188	192	185	175	161	145	131	107				
Submission 2020	140	188	192	185	175	161	145	131	106				
absolute change	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13				
relative change	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.12%				



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter [8.1 - Recalculations](#).

Planned improvements

Given the limited quality of the emission factors applied, the inventory compiler will check a possible revision at least for the main pollutants.

FAQs

bibliography : 1 : BAFA, 2019: Federal Office of Economics and Export Control (Bundesamt für Wirtschaft und Ausfuhrkontrolle, BAFA): Amtliche Mineralöl-daten für die Bundesrepublik Deutschland; URL: https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel_amtliche_daten_2017_dezember.html, Eschborn, 2019. : 2 : AGEb, 2019b: Working Group on Energy Balances (Arbeitsgemeinschaft Energiebilanzen (Hrsg.), AGEb): Daten - Sondertabellen - Heizwerte der Energieträger und Faktoren für die Umrechnung von spezifischen Mengeneinheiten in Wärmeinheiten (2005-2017); URL: <https://ag-energiebilanzen.de/#heizwerte2005bis2016>, Köln & Berlin, 2019. : 3 : Knörr et al. (2019b): Knörr, W., Heidt, C., Gores, S., & Bergk, F. (2018b): ifeu Institute for Energy and Environmental Research (Institut für Energie- und Umweltforschung Heidelberg gGmbH, ifeu): Aktualisierung des Modells TREMOD-Mobile Machinery (TREMOD MM) 2019, Heidelberg, 2019 : 4 : EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook 2019; Copenhagen, 2019. : 5 : Rentz et al., 2008: Nationaler Durchführungsplan unter dem Stockholmer Abkommen zu persistenten organischen Schadstoffen (POPs), im Auftrag des Umweltbundesamtes, FKZ 205 67 444, UBA Texte | 01/2008, January 2008, URL: <http://www.umweltbundesamt.de/en/publikationen/nationaler-durchfuehrungsplan-unter-stockholmer-bibliography>

¹⁾ (bibcite 1)

²⁾ (bibcite 2)

³⁾ (bibcite 4)

⁴⁾ (bibcite 3)