

# 1.A.5.a - Other, Stationary (including Military)

## Short description

In sub-category *1.A.5.a - Other, Stationary (including Military)* emissions from stationary fuel combustion in military facilities are reported.

Category Code	Method					AD					EF				
1.A.5.a	T2, T3					NS					CS				
Key Category	SO <sub>2</sub>	NO <sub>x</sub>	NH <sub>3</sub>	NMVOG	CO	BC	Pb	Hg	Cd	Diox	PAH	HCB	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
1.A.5.a	-/-	-/-	-/-	L/T	-/-	-/-	-	-/-	-	-/-	-/-	-/-	-/-	-/-	-/-

## Methodology

### Activity Data

The National Energy Balance of the Federal Republic of Germany (NEB) provides the basis for the activity data used.

However, since 1995 the NEB does not provide separate listings of military agencies' final energy consumption. Instead, these data is included in NEB line 67: 'commerce, trade, services and other consumers'. Therefore, additional energy statistics and fuel-specific data from the Federal Ministry of Defence (BMVg, 2020)<sup>1)</sup> is being used, providing the "Energy input for heat production in the German Federal Armed Forces" for all years as of 2000.

For liquid fuels, data is derived from the official mineral-oil data of the Federal Republic of Germany (Amtliche Mineralöl-daten der Bundesrepublik Deutschland), compiled annually by the Federal Office of Economics and Export Control (BAFA, 2020) The consumption figures given in units of [1000 t] are converted into [TJ] based on the relevant heating statistics published by the Working Group on Energy Balances.

All non-NEB figures are deducted from the figures in NEB line 67 (commerce, trade, services) and are reported in *1.A.5*, rather than in *1.A.4*.

### Emission Factors


The database for the emission factors used for source category *1.A.5.a* consists of the results of a research project carried out by the University of Stuttgart, under commission to the Federal Environment Agency (Struschka, 2008)<sup>2)</sup>. Within that project, device-related and source-category-specific emission factors for combustion systems in military agencies were calculated, with a high level of detail, for all important emissions components for the reference year 2005. The method used to determine the factors conforms to that described for source category *1.A.4*. The following table shows the sectoral emission factors used.

Table 1: Emission factors for Military stationary combustion plants

	NO <sub>x</sub>	SO <sub>x</sub>	NMVOC
	[kg/T]		
Hard coal	46	403	1
Brown coal briquettes	86	289	332
Light heating oil	46	77	2.8
Gaseous fuels	25	0.5	0.34

## Trend discussion for key sources

The following diagrams give an overview and assistance for explaining dominant emission trends of selected pollutants



NMVOC emissions show a remarkable falling trend due to the closure of military agencies especially at the beginning of the 1990s as well as the reducing use of lignite for heating purposes.

## Recalculations

Recalculations of PAH Emission due to a Revision of PAH Emission factors for all small combustion

sources.



For more information on recalculated emission estimates for Base Year and 2018, please see the pollutant-specific recalculation tables following chapter [8.1 - Recalculations](#).

## Planned improvements

Currently no improvements are planned.

---

<sup>1)</sup> Bundesministerium der Verteidigung: Energieeinsatz zur Wärmeversorgung der Bundeswehr (unpublished).

<sup>2)</sup> Struschka, Dr. M., Kilgus, D., Springmann, M.; Baumbach, Prof.Dr. Günter: Effiziente Bereitstellung aktueller Emissionsdaten für die Luftreinhaltung; UBA Forschungsbericht 205 42 322; Dessau, 2008. URL:

\*<https://www.umweltbundesamt.de/publikationen/effiziente-bereitstellung-aktueller-emissionsdaten>