

1.A.4.c ii (b) - Off-road Vehicles and other Machinery: Forestry

Short description

Under sub-category *1.A.4.c ii (b)* fuel combustion activities and resulting emissions from off-road vehicles and mobile machinery used in forestry are reported.

NFR-Code	Source category	Method	AD	EF	Key Category Analysis
1.A.4.c ii (a)	Off-road Vehicles and Other Machinery: Agriculture	T1, T2	NS, M	CS, D, M	see superordinate chapter

Methodology

Activity data

Primary activity data (PAD) are taken from National Energy Balances (NEBs) line 67: 'Commercial, trade, services and other consumers' (AGEB, 2020) ¹⁾.

Following the deduction of energy inputs for military vehicles as provided in (BAFA, 2020) ²⁾, the remaining amounts of gasoline and diesel oil are apportioned onto off-road construction vehicles (NFR 1.A.2.g vii) and off-road vehicles in commercial/institutional use (1.A.4. ii) as well as agriculture and forestry (NFR 1.A.4.c ii) based upon annual shares derived from TREMOD MM (Knörr et al. (2020b) ³⁾ (cf. [NFR 1.A.4 - mobile](#)]).

Table 1: Annual contribution of forestry vehicles and mobile machinery to the primary fuel delivery data provided in NEB line 67

1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Diesel fuels													
2,41%	1,36%	2,16%	2,88%	2,92%	2,99%	2,77%	2,76%	2,81%	2,89%	2,72%	2,79%	3,35%	3,54%
Gasoline fuels													
68,5%	40,3%	44,9%	41,4%	35,5%	35,6%	33,1%	32,9%	33,1%	33,3%	31,6%	31,9%	35,8%	36,8%

source: own estimates based on ⁴⁾

Table 2: Annual mobile fuel consumption in forestry, in terajoules

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Diesel Oil	2.695	1.332	2.051	2.368	2.583	2.712	2.472	2.559	2.718	2.931	2.865	3.024	3.393	3.625
Biodiesel	3.093	3.004	3.325	3.022	1.543	1.404	392	383	412	1.660	1.575	1.588	1.741	1.739
Gasoline	0	0	0	151	198	189	174	152	167	160	152	161	197	206
Biogasoline	0	0	0	21	60	58	17	16	18	72	68	67	78	75
Σ 1.A.4.c ii (ii)	5.788	4.336	5.375	5.562	4.383	4.364	3.055	3.110	3.316	4.824	4.660	4.840	5.409	5.646

Emission factors

The emission factors used here are of rather different quality: For all **main pollutants, carbon monoxide** and **particulate matter**, annually changing values computed within TREMOD MM (Knörr et al. (2020b))⁵⁾ are used, representing the development of mitigation technologies and the effect of fuel-quality legislation.

Table 3: Annual country-specific emission factors from TREMOD MM¹⁾

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Gasoline fuels														
NH₃	0,075	0,083	0,083	0,086	0,087	0,088	0,091	0,092	0,092	0,092	0,092	0,092	0,092	0,092
NM VOC²⁾	3,04	6,37	4,67	4,56	4,83	4,74	5,23	5,22	5,11	5,00	5,32	5,19	4,30	4,07
NM VOC³⁾	5795	5099	5099	5320	5424	4858	3596	2897	2897	2897	2897	2897	2897	2901
NO_x	42,6	49,4	49,4	76,4	86,0	78,5	63,1	55,1	55,1	55,1	55,1	55,1	55,1	55,1
SO_x	10,1	8,27	3,22	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37
BC⁵⁾	5,09	3,73	3,73	3,86	3,91	3,96	4,08	4,13	4,13	4,13	4,13	4,13	4,13	4,13
PM⁴⁾	102	74,6	74,6	77,2	78,1	79,2	81,5	82,7	82,7	82,7	82,7	82,7	82,7	82,7
TSP⁶⁾	2,35	0,82	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
CO	16813	14796	14796	15371	15609	15827	16279	16514	16514	16514	16514	16514	16514	16514
Pb	1,471	0,516	0	0	0	0	0	0	0	0	0	0	0	0
Diesel fuels														
NH₃	0,164	0,169	0,174	0,176	0,177	0,177	0,177	0,177	0,177	0,177	0,177	0,177	0,177	0,178
NM VOC	191	157	130	80,2	46,5	42,4	38,7	35,0	31,4	28,1	25,3	22,9	20,9	19,1
NO_x	982	1.053	1.071	834	543	495	454	422	397	375	351	326	305	287
SO_x	79,6	60,5	14,0	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37	0,37
BC⁵⁾	84,6	60,8	41,8	27,7	21,6	21,0	20,4	19,3	17,8	16,1	14,5	13,0	11,5	10,1
PM⁴⁾	155	112	75,8	45,3	30,4	28,8	27,3	25,3	23,1	20,8	18,6	16,5	14,6	12,9
CO	689	619	554	395	282	269	258	252	249	248	249	251	254	257

¹⁾ due to lack of better information: similar EF are applied for fossil and biofuels

²⁾ from fuel combustion

³⁾ from gasoline evaporation

⁴⁾ EF(PM_{2,5}) also applied for PM₁₀ and TSP (assumption: > 99% of TSP consists of PM_{2,5})

⁵⁾ estimated via a f-BCs as provided in ⁶⁾, Chapter 1.A.2.g vii, 1.A.4.a ii, b ii, c ii, 1.A.5.b i - Non-road, note to Table 3-1: Tier 1 emission factors for off-road machinery

⁶⁾ from leaded gasoline (until 1997)



With respect to the emission factors applied for particulate matter, given the circumstances during test-bench measurements, condensables are most likely included at least partly. ¹⁾

For information on the **emission factors for heavy-metal and POP exhaust emissions**, please refer to [Appendix 2.3 - Heavy Metal \(HM\) exhaust emissions from mobile sources](#) and [Appendix 2.4 - Persistent Organic Pollutant \(POP\) exhaust emissions from mobile sources](#)].

Recalculations

Revisions in **activity data** result from slightly adapted EBZ67 shares as well as the implementation of primary activity data from the now finalised NEB 2018.

Table 5: Revised annual contribution of 1.A.2.g vii to fuel-specific over-all fuel deliveries provided in NEB line 67

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018
diesel fuels													
Submission 2021	0,024	0,014	0,022	0,029	0,029	0,030	0,028	0,028	0,028	0,029	0,027	0,028	0,034
Submission 2020	0,024	0,013	0,020	0,027	0,027	0,027	0,025	0,026	0,026	0,027	0,025	0,026	0,026
absolute change	0,0003	0,0002	0,0011	0,0023	0,0024	0,0025	0,0024	0,0020	0,0022	0,0023	0,0021	0,0022	0,0076
relative change	1,08%	1,62%	5,38%	8,71%	9,16%	9,27%	9,36%	7,86%	8,29%	8,56%	8,54%	8,63%	29,4%
gasoline fuels													
Submission 2021	0,685	0,403	0,449	0,414	0,355	0,356	0,331	0,329	0,331	0,333	0,316	0,319	0,358
Submission 2020	0,685	0,403	0,449	0,416	0,360	0,362	0,337	0,335	0,338	0,341	0,324	0,327	0,327
absolute change	0,000	0,000	0,000	-0,002	-0,005	-0,005	-0,006	-0,006	-0,007	-0,008	-0,008	-0,009	0,031
relative change	0,00%	0,00%	0,00%	-0,46%	-1,29%	-1,46%	-1,70%	-1,88%	-2,06%	-2,23%	-2,48%	-2,66%	9,34%

Table 6: Revised activity data, in terajoules

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018
diesel fuels													
Submission 2021	2.695	1.332	2.051	2.519	2.781	2.902	2.646	2.711	2.886	3.092	3.016	3.185	3.590
Submission 2020	2.666	1.311	1.946	2.221	2.519	2.645	2.408	2.510	2.662	2.846	2.778	2.931	2.758
absolute change	28,9	21,2	105	298	262	257	239	201	223	245	239	254	831
relative change	1,08%	1,62%	5,38%	13,4%	10,4%	9,70%	9,91%	8,00%	8,38%	8,62%	8,59%	8,67%	30,1%
gasoline fuels													
Submission 2021	3.093	3.004	3.325	3.043	1.603	1.462	409	400	430	1.732	1.644	1.655	1.819
Submission 2020	3.093	3.004	3.325	3.057	1.624	1.484	416	407	439	1.772	1.685	1.700	1.663
absolute change	0,00	0,00	0,00	-14,0	-21,0	-21,7	-7,1	-7,7	-9,0	-39,6	-41,8	-45,1	155
relative change	0,00%	0,00%	0,00%	-0,46%	-1,29%	-1,46%	-1,70%	-1,88%	-2,06%	-2,23%	-2,48%	-2,66%	9,34%
over-all fuel consumption													
Submission 2021	5.788	4.336	5.375	5.562	4.383	4.364	3.055	3.110	3.316	4.824	4.660	4.840	5.409

Submission 2020	5.759	4.315	5.271	5.278	4.143	4.129	2.824	2.917	3.102	4.618	4.463	4.631	4.422
absolute change	28,9	21,2	105	284	241	235	231	193	214	206	197	209	987
relative change	0,50%	0,49%	1,99%	5,38%	5,81%	5,69%	8,20%	6,62%	6,90%	4,46%	4,41%	4,51%	22,3%

As in contrast, all **emission factors** remain unrevised compared to last year's submission, emission estimates for the years as of 2015 change in accordance with the underlying activity data.



For specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter [8.1 - Recalculations](#).

¹⁾ AGEb, 2020: Working Group on Energy Balances (Arbeitsgemeinschaft Energiebilanzen (Hrsg.), AGEb): Energiebilanz für die Bundesrepublik Deutschland; URL: <http://www.ag-energiebilanzen.de/7-0-Bilanzen-1990-2018.html>, (Aufruf: 29.11.2020), Köln & Berlin, 2020.

²⁾ BAFA, 2020: Federal Office of Economics and Export Control (Bundesamt für Wirtschaft und Ausfuhrkontrolle, BAFA): Amtliche Mineralöl-daten für die Bundesrepublik Deutschland; URL: https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel_amtliche_daten_2018_dezember.html, Eschborn, 2020.

^{3), 4), 5)} Knörr et al. (2020b): Knörr, W., Heidt, C., Gores, S., & Bergk, F.: ifeu Institute for Energy and Environmental Research (Institut für Energie- und Umweltforschung Heidelberg gGmbH, ifeu): Aktualisierung des Modells TREMOD-Mobile Machinery (TREMOD MM) 2020, Heidelberg, 2020.

⁶⁾ EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook – 2019, Copenhagen, 2019.

¹⁾ During test-bench measurements, temperatures are likely to be significantly higher than under real-world conditions, thus reducing condensation. On the contrary, smaller dilution (higher number of primary particles acting as condensation germs) together with higher pressures increase the likelihood of condensation. So over-all condensables are very likely to occur but different to real-world conditions.