

## **1.A.3.a ii (ii) - Domestic Civil Aviation: Cruise**

## **Short description**

In NFR category 1.A.3.a ii (ii) - Domestic Civil Aviation: Cruise emissions from domestic flights between German airports during cruise stage (above 3,000 feet of altitude) are reported.

In the following, information on sub-category specific activity data, (implied) emission factors and emission estimates are provided.

# Methodology

## Activity Data

Specific fuel consumption during LTO-stage is calculated within TREMOD AV as described in the [superordinate chapter](#).

Table 1: annual jet kerosene & avgas consumption during cruise-stage, in terajoules

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>Kerosene</b>	21.690	19.937	25.301	24.071	24.736	25.337	25.111	24.048	22.503	20.552	21.026	19.762	19.038	19.195	20.067	20.793	21.067	21.573
<b>Avgas</b>	1.580	614	614	291	260	228	259	244	237	283	246	199	180	233	145	142	116	72

source: Knörr et al. (2019c)<sup>1)</sup> & Gores (2019)<sup>2)</sup>

# Emission factors

All country specific emission factors used for emission reporting were basically ascertained within UBA project FKZ 360 16 029<sup>3)</sup> and have since then been compiled, revised and maintained in TREMOD AV<sup>4)</sup>

For more information, please see the [superordinate chapter](#) on civil aviation.

**Table 2: Annual country-specific emission factors, in kg/TJ**

<b>BC<sup>2</sup></b>														
<b>CO</b>														

<sup>1</sup> EF(TSP) also applied for PM<sub>10</sub> and PM<sub>2.5</sub> (assumption: > 99% of TSP consists of PM<sub>2.5</sub>)

<sup>2</sup> estimated via a f-BC of 0.48 as provided in <sup>5)</sup>, Chapter: 1.A.3.a, 1.A.5.b Aviation, page 49: "Conclusion".



For the country-specific emission factors applied for particulate matter, no clear indication is available, whether or not condensables are included.

## Trend discussion for Key Sources

NFR 1.A.3.a ii (ii) - Domestic Civil Aviation - Cruise is **not included in the national emission totals** and hence **not included in the key category analysis**.

## Recalculations

**Activity data** have been revised for all years within TREMOD AV to keep in line with information available from the 2019 EMEP/EEA Guidebook <sup>6)</sup> and Eurocontrol's AEM model <sup>7)</sup>.

Table 3: Revised kerosene consumption in 1.A.3.a ii (ii), in terajoules

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
<b>JET KEROSENE</b>																	
<b>Submission 2021</b>	21.690	19.937	25.301	24.071	24.736	25.337	25.111	24.048	22.503	20.552	21.026	19.762	19.038	19.195	20.067	20.793	21.067
<b>Submission 2020</b>	20.024	20.875	22.967	21.565	22.122	22.904	23.145	21.771	21.579	21.776	20.673	18.717	19.614	19.730	19.746	19.074	19.178
<b>absolute change</b>	1.667	-938	2.334	2.505	2.615	2.433	1.966	2.277	923	-1.224	354	1.045	-576	-535	321	1.719	1.888
<b>relative change</b>	8,32%	-4,49%	10,16%	11,62%	11,82%	10,62%	8,49%	10,46%	4,28%	-5,62%	1,71%	5,58%	-2,94%	-2,71%	1,63%	9,01%	9,85%
<b>AVGAS</b>																	
<b>Submission 2021</b>	1.580	614	614	291	260	228	259	244	237	283	246	199	180	233	145	142	116
<b>Submission 2020</b>	IE																
<b>absolute change</b>	1.580	614	614	291	260	228	259	244	237	283	246	199	180	233	145	142	116

In parallel, the majority of **country-specific emission factors** has been revised within TREMOD AV based on information available from Eurocontrol's AEM model <sup>8)</sup>.

Table 4: Revised emission-factor values, in [kg/TJ]

< NMVOC																	
~ Submission 2020	> 15.1	> 17.0	> 17.2	> 19.9	> 20.9	> 21.5	> 21.7	> 21.6	> 21.3	> 21.4	> 21.7	> 23.3	> 23.6	> 23.8	> 19.1	> 19.5	
~ Submission 2019	> 9.3	> 11.6	> 10.5	> 12.1	> 13.2	> 13.3	> 13.0	> 13.1	> 13.1	> 13.3	> 13.2	> 13.7	> 14.1	> 13.5	> 13.4	> 13.6	

~ absolute change	> 5.84	> 5.46	> 6.68	> 7.76	> 7.69	> 8.21	> 8.69	> 8.49	> 8.24	> 8.10	> 8.48	> 9.68	> 9.48	> 10.28	> 5.66	> 5.83
~ relative change	63.0%	47.2%	63.8%	64.0%	58.1%	61.7%	66.7%	64.8%	63.1%	61.1%	64.0%	70.9%	67.1%	75.9%	42.2%	42.7%
<b>&lt; Nitrogen oxides - NO<sub>x</sub>,x,,</b>																
~ Submission 2020	> 330	> 367	> 332	> 319	> 317	> 320	> 333	> 346	> 354	> 356	> 360	> 359	> 366	> 365	> 379	> 380
~ Submission 2019	> 342	> 381	> 351	> 338	> 339	> 346	> 358	> 366	> 372	> 375	> 380	> 385	> 384	> 388	> 396	> 402
~ absolute change	-12.51	-14.42	-18.74	-19.36	-21.48	-26.16	-25.15	-20.71	-18.06	-19.21	-20.09	-26.37	-17.21	-22.82	-17.11	-21.59
~ relative change	-3.65%	-3.79%	-5.34%	-5.72%	-6.34%	-7.55%	-7.02%	-5.65%	-4.86%	-5.13%	-5.29%	-6.85%	-4.48%	-5.88%	-4.33%	-5.38%
<b>&lt; Sulphur oxides - SO<sub>x</sub>,x,,</b>																
~ Submission 2020	> 19.7	> 19.5	> 19.5	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	> 19.6	
~ Submission 2019	> 25.1	> 15.2	> 8.5	> 6.3	> 5.9	> 5.5	> 5.1	> 4.7	> 4.7	> 4.7	> 4.7	> 4.7	> 4.7	> 4.7	> 4.7	
~ absolute change	> -5.45	> 4.30	> 11.08	> 13.28	> 13.71	> 14.13	> 14.55	> 14.98	> 14.98	> 14.98	> 14.98	> 14.98	> 14.98	> 14.98	> 14.98	
~ relative change	-21.7%	28.2%	131%	209%	232%	257%	287%	322%	322%	322%	322%	322%	322%	322%	322%	
<b>&lt; Black carbon - BC</b>																
~ Submission 2020	> 2.02	> 2.49	> 2.44	> 2.53	> 2.56	> 2.47	> 2.20	> 2.22	> 2.28	> 2.25	> 2.26	> 2.27	> 2.38	> 2.40	> 2.24	> 2.21
~ Submission 2019	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	> 2.23	
~ absolute change	> -0.21	> 0.26	> 0.21	> 0.30	> 0.32	> 0.24	> -0.03	> -0.01	> 0.05	> 0.02	> 0.03	> 0.03	> 0.15	> 0.17	> 0.01	> -0.02
~ relative change	-9.52%	11.61%	9.50%	13.39%	14.49%	10.85%	-1.42%	-0.49%	2.32%	0.82%	1.34%	1.55%	6.53%	7.57%	0.29%	-0.95%
<b>&lt; Particulate matter - PM</b>																
~ Submission 2020	> 4.21	> 5.19	> 5.09	> 5.27	> 5.33	> 5.16	> 4.58	> 4.63	> 4.76	> 4.69	> 4.71	> 4.72	> 4.95	> 5.00	> 4.66	> 4.61
~ Submission 2019	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	> 4.65	
~ absolute change	> -0.44	> 0.54	> 0.44	> 0.62	> 0.67	> 0.50	> -0.07	> -0.02	> 0.11	> 0.04	> 0.06	> 0.07	> 0.30	> 0.35	> 0.01	> -0.04
~ relative change	-9.52%	11.6%	9.50%	13.4%	14.5%	10.8%	-1.42%	-0.49%	2.32%	0.82%	1.34%	1.55%	6.53%	7.57%	0.29%	-0.95%
<b>&lt; Carbon monoxide - CO</b>																
~ Submission 2020	> 144.5	> 145.2	> 179.0	> 195.3	> 194.5	> 193.3	> 195.1	> 194.2	> 190.2	> 190.0	> 194.1	> 205.0	> 206.2	> 208.5	> 149.0	> 145.3
~ Submission 2019	> 85.5	> 88.2	> 111.7	> 111.0	> 111.0	> 105.2	> 102.4	> 104.8	> 105.8	> 105.2	> 102.1	> 98.7	> 100.4	> 104.3	> 98.2	> 91.7
~ absolute change	58.93	56.97	67.33	84.30	83.50	88.08	92.71	89.41	84.38	84.77	91.96	106.30	105.85	104.16	50.81	53.62
~ relative change	68.9%	64.6%	60.3%	75.9%	75.2%	83.7%	90.5%	85.3%	79.7%	80.6%	90.1%	108%	105.4%	99.9%	51.7%	58.5%



For more information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter [8.1 - Recalculations](#).

## Uncertainties

For uncertainties information, please see [main chapter](#) on civil aviation.

## Planned improvements

For information on planned improvements, please see [main chapter](#) on civil aviation.

## FAQs

**bibliography** : 1 : Knörr, W., Schacht, A., & Gores, S. (2010): Entwicklung eines eigenständigen Modells zur Berechnung des Flugverkehrs (TREMOD-AV) : Endbericht. Endbericht zum F+E-Vorhaben 360 16 029, URL:

<https://www.umweltbundesamt.de/publikationen/entwicklung-eines-modells-zur-berechnung>; Berlin & Heidelberg, 2012. : 2 : Knörr et al. (2019c): Knörr, W., Schacht, A., & Gores, S.: TREMOD Aviation (TREMOD AV) 2019 - Revision des Modells zur Berechnung des Flugverkehrs (TREMOD-AV).

Heidelberg, Berlin: Ifeu Institut für Energie- und Umweltforschung Heidelberg GmbH & Öko-Institut e.V., Berlin & Heidelberg, 2019. : 3 : Gores (2018): Inventartool zum deutschen Flugverkehrsinventory 1990-2018, im Rahmen der Aktualisierung des Moduls TREMOD-AV im Transportemissionsmodell TREMOD, Berlin, 2019. : 4 : EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook 2019,

<https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-b-sectoral-guidance-chapters/1-energy/1-a-combustion/1-a-3-a-aviation/view>; Copenhagen, 2019. : 5 : Eurocontrol (2019): Advanced emission model (AEM); <https://www.eurocontrol.int/model/advanced-emission-model>; 2019

<sup>1)</sup> (bibcite 2)

<sup>2)</sup> (bibcite 3)

<sup>3)</sup> (bibcite 1)

<sup>4)</sup> (bibcite 2)

<sup>5)</sup> (bibcite 4)

<sup>6)</sup> (bibcite 4)

<sup>7)</sup> (bibcite 5)

<sup>8)</sup> (bibcite 5)