

1.A.3.a ii (i) - Domestic Civil Aviation: LTO

Short description

In NFR category *1.A.3.a ii (i) - Domestic Civil Aviation: LTO* emissions from domestic flights between German airports occurring during LTO stage (Landing/Take-off: 0-3,000 feet) are reported.

Category Code	Method					AD					EF				
1.A.3.a ii(i)	T1, T2, T3					NS, M					CS, D, M				
Key Category	SO ₂	NO _x	NH ₃	NM VOC	CO	BC	Pb	Hg	Cd	Diox	PAH	HCb	TSP	PM ₁₀	PM _{2.5}
1.A.3.a ii(i)	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-	-/-	-	-/-	-/-	-/-

T = key source by Trend **L** = key source by Level

Methods	
D	Default
RA	Reference Approach
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
T3	Tier 3 / Detailed Methodology *
C	CORINAIR
CS	Country Specific
M	Model

* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

AD - Data Source for Activity Data	
NS	National Statistics
RS	Regional Statistics
IS	International Statistics
PS	Plant Specific data
AS	Associations, business organisations
Q	specific questionnaires, surveys
EF - Emission Factors	
D	Default (EMEP Guidebook)
C	Confidential
CS	Country Specific
PS	Plant Specific data

In the following, information on sub-category specific AD, (implied) emission factors and emission estimates are provided.

Methodology

Activity Data

Specific jet kerosene consumption during LTO-stage is calculated within TREMOD AV as described in the [superordinate chapter](#).

Table 1: Percentual annual fuel consumption during LTO-stage of domestic flights

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Jet Kerosene	30,2	29,4	27,9	27,6	27,5	27,3	27,3	27,3	27,6	27,7	28,0	27,9	27,7	27,7	28,1	28,3	28,4	28,1
Aviation Gasoline	18,9	36,0	33,6	50,2	52,1	55,4	51,6	51,2	49,9	46,7	49,1	54,0	56,8	51,4	61,8	62,0	68,0	76,0


source: Knörr et al. (2019c) ¹⁾ & Gores (2019) ²⁾

As explained above, the use of aviation gasoline is - due to a lack of further information - assumed to entirely take place within the LTO-range.

Table 2: annual LTO fuel consumption for domestic flights, in terajoule

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Jet Kerosene	9,380	8,303	9,811	9,187	9,402	9,493	9,422	9,021	8,589	7,869	8,171	7,633	7,297	7,358	7,844	8,210	8,362	8,417
Aviation Gasoline	368	346	311	293	283	283	276	255	236	248	237	234	237	246	234	232	248	229
Σ 1.A.3.a ii (i)	9,748	8,649	10,122	9,481	9,686	9,776	9,698	9,277	8,825	8,117	8,408	7,868	7,534	7,604	8,078	8,442	8,610	8,647

source: Knörr et al. (2019c) ³⁾ & Gores (2019) ⁴⁾



Emission factors

All country-specific emission factors used for emission reporting were basically ascertained within UBA project FKZ 360 16 029 (Knörr, W., Schacht, A., & Gores, S. (2010))⁵⁾ and have since then been compiled, revised and maintained in TREMOD AV⁶⁾.

Furthermore, the **newly implemented EF(BC)** have been estimated via f-BCs as provided in the 2019 EMEP/EEA Guidebook⁷⁾, Chapter 1.A.3.a, 1.A.5.b Aviation, page 49: "Conclusion".

For more details, please see [superordinate chapter](#) on civil aviation.

Table 3: Country-specific emission factors, in kg/TJ

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
JET KEROSENE																		
NH₃	3,98	3,95	3,95	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97	3,97
NMVOC	28,4	28,9	30,5	32,4	33,9	34,4	34,7	33,2	32,3	31,9	32,0	34,9	37,0	36,9	36,5	38,3	39,1	43,2
NO_x	295	324	287	277	276	281	290	300	304	309	312	311	310	312	321	322	316	308
SO_x	19,7	19,5	19,5	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6	19,6
BC¹	1,43	1,57	1,54	1,61	1,62	1,59	1,47	1,48	1,51	1,50	1,52	1,53	1,50	1,52	1,44	1,44	1,56	1,55
PM²	2,99	3,28	3,21	3,36	3,38	3,32	3,06	3,07	3,14	3,13	3,17	3,18	3,12	3,17	3,01	2,99	3,25	3,22
CO	212	211	275	291	292	286	280	266	260	254	252	260	265	265	252	255	262	276
AVIATION GASOLINE																		
NH₃																		
NMVOC																		
NO_x																		
SO_x																		
BC¹																		
PM²																		
TSP³																		
CO																		

¹ estimated via a f-BCs (avgas: 0.15, jet kerosene: 0.48) as provided in⁸⁾

² EF(PM_{2.5}) also applied for PM₁₀ and TSP (assumption: > 99% of TSP from diesel oil combustion consists of PM_{2.5})

³ also including TSP from lead: EF(TSP) = 1.6 x EF(Pb) - see road transport



For the country-specific emission factors applied for particulate matter, no clear indication is available, whether or not condensables are included.

Table 4: Tier1 emission factors for heavy-metal and POP exhaust emissions

	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	B[a]P	B[b]F	B[k]F	I[...]p	PAH 1-4	PCDD/F
	[g/TJ]									[mg/TJ]				[µg/TJ]	
Kerosene	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Aviation gasoline	9,481 ¹	0.005	0.200	0.007	0.145	0.103	0.053	0.005	0.758	126	182	90	205	602	NE

¹ calculated from the average lead content of AvGas 100 LL (low-lead) of 0.56 g Pb/liter


Trend discussion for Key Sources

NFR 1.A.3.a ii (i) - Domestic Civil Aviation - LTO is **no key source**.

Where **sulphur oxides** emissions are dominated by jet kerosene due to the amount of fuel used, the majority of **carbon monoxide** stems from the consumption of avgas given the much higher emission factor applied to this fuel.

[gallery size="medium" : 1A3aii\(i\)_SOx.png : 1A3aii\(i\)_CO.png gallery](#)

Lead emissions on the other hand, with no emission factor available for jet kerosene, are only calculated for avgas.



Recalculations

Activity data

In order to keep in line with the regularly updated data sets provided to the EEA by Eurocontrol, the average fuel use per LTO cycle has been updated again within TREMOD Aviation but with much smaller impact as in last year's submission.

Furthermore, as explained in the superordinate chapter, avgas consumption for international flights and outside the LTO range has been estimated for the first time for this submission, with the

Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	
NITROGEN OXIDES																	
Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	
SULPHUR OXIDES																	
Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	
BLACK CARBON - BC																	
Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	
PARTICULATE MATTER - PM																	
Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	
CARBON MONOXIDE - CO																	
Submission 2021																	
Submission 2020																	
absolute change																	
relative change																	

The TSP emissions calculated depend directly on the reported lead emissions: The emission factor for TSP is 1.6 times the emission factor used for lead: $EF(TSP) = 1.6 \times EF(Pb)$. The applied procedure is similar to the one used for calculating TSP emissions from leaded gasoline used in road transport.

bibliography : 1 : Knörr, W., Schacht, A., & Gores, S. (2010): Entwicklung eines eigenständigen Modells zur Berechnung des Flugverkehrs (TREMOT-AV) : Endbericht. Endbericht zum F+E-Vorhaben 360 16 029, URL:

<https://www.umweltbundesamt.de/publikationen/entwicklung-eines-modells-zur-berechnung>; Berlin & Heidelberg, 2012. : 2 : Knörr et al. (2019c): Knörr, W., Schacht, A., & Gores, S.: TREMOD Aviation (TREMOT AV) 2018 - Revision des Modells zur Berechnung des Flugverkehrs (TREMOT-AV).

Heidelberg, Berlin: Ifeu Institut für Energie- und Umweltforschung Heidelberg GmbH & Öko-Institut e.V., Berlin & Heidelberg, 2019. : 3 : Gores (2019): Inventartool zum deutschen Flugverkehrsinventar 1990-2018, im Rahmen der Aktualisierung des Moduls TREMOD-AV im Transportemissionsmodell TREMOD, Berlin, 2019. : 4 : EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook 2019,

<https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-b-sectoral-guidance-chapters/1-energy/1-a-combustion/1-a-3-a-aviation/view>; Copenhagen, 2019. : 5 : Eurocontrol (2019):

Advanced emission model (AEM); <https://www.eurocontrol.int/model/advanced-emission-model>; 2019
bibliography

¹⁾ (bibcite 2)

²⁾ (bibcite 3)

³⁾ (bibcite 2)

⁴⁾ (bibcite 3)

⁵⁾ (bibcite 1)

⁶⁾ (bibcite 2)

⁷⁾ (bibcite 4)

⁸⁾ (bibcite 4)

⁹⁾ (bibcite 4)

¹⁰⁾ (bibcite 5)