

1.A.3.c - Transport: Railways

Short description

In category 1.A.3.c - Railways, emissions from fuel combustion in German railways and from the related abrasion and wear of contact line, braking systems and tyres on rails are reported.

Method	AD	EF	Key Category
T1, T2	NS, M	CS, D, M	L: TSP, PM _{2.5} , L&T: PM ₁₀ , PM _{2.5} , L: TSP

T = key source by Trend **L** = key source by Level

Methods	
D	Default
RA	Reference Approach
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
T3	Tier 3 / Detailed Methodology *
C	CORINAIR
CS	Country Specific
M	Model
* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.	
AD - Data Source for Activity Data	
NS	National Statistics
RS	Regional Statistics
IS	International Statistics
PS	Plant Specific data
AS	Associations, business organisations
Q	specific questionnaires, surveys
EF - Emission Factors	
D	Default (EMEP Guidebook)
C	Confidential
CS	Country Specific
PS	Plant Specific data

Germany's railway sector is undergoing a long-term modernisation process, aimed at making electricity the main energy source for rail transports. Use of electricity, instead of diesel fuel, to power locomotives has been continually increased, and electricity now provides 80% of all railway traction power. Railways' power stations for generation of traction current are allocated to the stationary component of the energy sector (1.A.1.a) and are not included in the further description that follows here. In energy input for trains of German railways, diesel fuel is the only energy source that plays a significant role apart from electric power.

Method

Activity Data

Basically, total inland deliveries of *diesel oil* are available from the National Energy Balances (NEBs) (AGEB, 2019) ¹⁾. This data is based upon sales data of the Association of the German Petroleum Industry (MWV) ²⁾. As a recent revision of MWV data on diesel oil sales for the years 2005 to 2009 has not yet been adopted to the respective NEBs, this original MWV data has been used for this five years.

Data on the consumption of biodiesel in railways is provided in the NEBs as well, from 2004 onward. But as the NEBs do not provide a solid time series regarding most recent years, the data used for the inventory is estimated based on the prescribed shares of biodiesel to be added to diesel oil.

Small quantities of *solid fuels* are used for historical steam engines vehicles operated mostly for tourism and exhibition purposes. Official fuel delivery data are available for lignite, through 2002, and for hard coal, through 2000, from the NEBs. In order to complete these time series, a study was carried out in 2012 by Hedel, R., and Kunze, J. (2012) ³⁾. During this study, questionnaires were provided to any known operator of historical steam engines in Germany. Here, due to limited data archiving, nearly complete data could only be gained for years as of 2005. For earlier years, in order to achieve a solid time series, conservative gap filling was applied. A follow-up study to gain original consumption data for 2015 was carried out in 2016 by Illichmann, S. (2016) ⁴⁾.

Table 1: Overview of activity-data sources for domestic fuel sales to railway operators

Activity	data source / quality of activity data
combustion of:	
Diesel oil	1990-2004: NEB lines 74 and 61: 'Schienenverkehr' / 2005-2009: MWV annual report, table: 'Sektoraler Verbrauch von Dieselmotorkraftstoff' / from 2010: NEB line 61
Biodiesel	calculated from official blending rates
Hard coal	1990-1994: NEB lines 74; 1995-2004: interpolated data; from 2005: original data from studies; 2016: forward extrapolation
Hard coal coke	1990-1997: NEB lines 74 and 61; 1998-2004: interpolated data; from 2005: original data from studies; 2016: forward extrapolation
Raw lignite	from 1990: NEB lines 74 and 61
Lignite briquettes	from 1990: NEB lines 74 and 61
abrasion and wear of contact line, braking systems and tyres on rails:	
transport performance data	in Mio ptkm (performance-ton-kilometers) derived from the TREMOD model

Table 2: Annual fuel consumption in German railways, in terajoules

=	= 1990	= 1995	= 2000	= 2005	= 2010	= 2011	= 2012	= 2013	= 2014	= 2015	= 2016	= 2017	= 2018							
~ Diesel Oil	> 38,458	> 31,054	> 25,410	> 18,142	> 14,626	> 14,730	> 13,514	> 13,771	> 12,283	> 13,321	> 13,775	> 11,344	> 10,961							
~ Biodiesel	> 0	> 0	> 0	> 397	> 949	> 966	> 882	> 798	> 745	> 720	> 724	> 602	> 633							

~ Liquids TOTAL	> 38,458	> 31,054	> 25,410	> 18,539	> 15,575	> 15,696	> 14,396	> 14,569	> 13,028	> 14,041	> 14,499	> 11,946	> 11,594						
~ Lignite Briquettes	> 0.00	> 0.00	> 431.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00						
~ Raw Lignite	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00	> 0.00						
~ Hard Coal	> 576	> 250	> 250	> 255	> 314	> 345	> 357	> 352	> 341	> 339	> 340	> 340	> 340						
~ Hard Coal Coke	> 0	> 86	> 1	> 1	> 1	> 1	> 1	> 1	> 1	> 1	> 1	> 1	> 1						
Solids TOTAL	> 576	> 336	> 682	> 256	> 315	> 346	> 357	> 353	> 342	> 340	> 341	> 341	> 341						
Σ 1.A.3.c	~ 39,034	~ 31,390	~ 26,092	~ 18,795	~ 15,890	~ 16,041	~ 14,754	~ 14,921	~ 13,370	~ 14,381	~ 14,839	~ 12,287	~ 11,934						

The use of other fuels – such as vegetable oils or gas – in private narrow-gauge railway vehicles has not been included to date and may still be considered negligible.

Table 3: Annual transport performance, in Mio tkm (ton-kilometers)

	1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Electric Traction	361,515	337,853	361,633	356,605	344,546	342,701	350,085	335,298	331,235	323,387	295,798	296,280	288,336	
Diesel Traction	98,812	58,805	37,237	26,540	26,702	27,403	26,791	23,768	23,734	21,397	21,484	21,365	19,580	
Σ 1.A.3.c	460,326	396,658	398,870	383,145	371,248	370,104	376,876	359,065	354,970	344,785	317,282	317,645	307,916	

[gallery size="medium" : 1A3c_AD\(TJ\).png : 1A3c_AD\(km\).png](#) [gallery](#)

Regarding particulate-matter and heavy-metal emissions from **abrasion and wear of contact line, braking systems, tyres on rails**, annual transport performances of railway vehicles with electrical and Diesel traction derived from Knörr et al. (2019a) ⁵⁾ are applied as activity data.

Emission factors

The (implied) emission factors used here for estimating **emissions from diesel fuel combustion** of very different quality: For main pollutants, CO and PM, annual tier2 IEF computed within the TREMOD model are used, representing the development of German railway fleet, fuel quality and mitigation technologies ⁶⁾. On the other hand, constant default values from (EMEP/EEA, 2019) ⁷⁾ are used for all reported PAHs and heavy metals and from Rentz et al. (2008) ⁸⁾ regarding PCDD/F. As no emission factors are available for HCB and PCBs, no such emissions have been calculated yet.

Regarding **emissions from solid fuels** used in historic steam engines, all emission factors displayed below have been adopted from small-scale stationary combustion.

Furthermore, regarding **emissions from abrasion and wear**, emission factors are calculated from PM₁₀, emission estimates directly provided by the German railroad company Deutsche Bahn AG. As these original emissions are only available as of 2013, implied EF(PM₁₀) were calculated from the emission estimates extrapolated backwards from 2013 to 1990 and the transport performance data available from TREMOD. Regarding PM_{2.5}, and TSP, due to lack of better information, a fractional distribution of 0.5 : 1 : 1 (PM_{2.5} : PM₁₀ : TSP) is assumed for now. Emission factors for emissions

[gallery size="medium" : 1A3c_EM_SO2.png](#) [gallery](#)

Regarding **heavy metals**, emissions from combustion of diesel oil and from abrasion and wear are estimated from tier1 default emission factors. Therefore, the emission trends reflect the development of diesel use and - for copper, chromium and nickel emissions resulting from the abrasion & wear of contact line and braking systems - the annual transport performance (see description of activity data above).

[gallery size="medium" : 1A3c_EM_Cu.png : 1A3c_EM_Cd.png](#) [gallery](#)

Recalculations

Given the revised NEB 2018, both the **activity data** for diesel oil and the annual amounts of blended biodiesel were revised accordingly.

Table 5: Revised 2017 fuel consumption, in terajoule

>	> Diesel Oil	> Biodiesel
~ Submission 2020	> 11,344	> 602
~ Submission 2019	> 13,690	> 726
~ absolute change	> -2,346	> -124
~ relative change	> -17.1%	> -17.1%

Due to the routine revision of the TREMOD model ¹⁰⁾, tier2 **emission factors** changed for recent years. Here, the revision results mainly from the consideration of revised NCvs for diesel oil as provided by the AGEb.

Table 6: Revised country-specific emission factors for diesel fuels, in kg/TJ

=	= 2005	= 2006	= 2007	= 2008	= 2009	= 2010	= 2011	= 2012	= 2013	= 2014	= 2015	= 2016	= 2017
< Nitrogen oxides - NO _x													
~ Submission 2020	> 1,111	> 1,058	> 1,028	> 1,010	> 991	> 970	> 990	> 919	> 899	> 886	> 826	> 801	> 775
~ Submission 2019	> 1,111	> 1,058	> 1,029	> 1,011	> 1,001	> 986	> 1,010	> 921	> 882	> 897	> 851	> 836	> 814
~ absolute change	> 0.00	> -0.75	> -0.44	> -1.81	> -9.91	> -15.63	> -19.97	> -2.05	> 16.27	> -10.45	> -25.27	> -34.82	> -38.70
~ relative change	> 0.00%	> -0.07%	> -0.04%	> -0.18%	> -0.99%	> -1.59%	> -1.98%	> -0.22%	> 1.84%	> -1.16%	> -2.97%	> -4.16%	> -4.75%
< Non-methane volatile organic compounds - NMVOC													
~ Submission 2020	> 64.8	> 61.8	> 57.3	> 55.6	> 51.2	> 52.0	> 54.3	> 44.8	> 42.2	> 41.2	> 38.5	> 38.2	> 37.2
~ Submission 2019	> 64.8	> 62.1	> 57.8	> 56.7	> 53.8	> 55.7	> 59.2	> 46.9	> 43.5	> 43.1	> 41.4	> 40.9	> 39.3

~ absolute change	> -0.04	> -0.33	> -0.48	> -1.05	> -2.60	> -3.79	> -4.84	> -2.09	> -1.33	> -1.95	> -2.87	> -2.66	> -2.08
~ relative change	> -0.06%	> -0.52%	> -0.83%	> -1.85%	> -4.85%	> -6.80%	> -8.18%	> -4.46%	> -3.06%	> -4.52%	> -6.93%	> -6.50%	> -5.30%
< Particulate matter - PM (PM _{2.5} = PM ₁₀ = TSP)													
~ Submission 2020	> 23.4	> 22.4	> 20.9	> 19.5	> 17.6	> 17.7	> 18.5	> 16.0	> 14.7	> 14.3	> 13.3	> 13.1	> 12.4
~ Submission 2019	> 23.4	> 22.5	> 21.1	> 19.9	> 18.2	> 18.6	> 19.8	> 16.6	> 14.8	> 15.4	> 14.7	> 14.6	> 13.7
~ absolute change	> -0.02	> -0.14	> -0.21	> -0.40	> -0.68	> -0.95	> -1.33	> -0.58	> -0.14	> -1.12	> -1.37	> -1.58	> -1.33
~ relative change	> -0.08%	> -0.62%	> -1.01%	> -2.03%	> -3.75%	> -5.07%	> -6.72%	> -3.48%	> -0.95%	> -7.25%	> -9.31%	> -10.79%	> -9.73%
< Black carbon - BC													
~ Submission 2020	> 15.2	> 14.5	> 13.6	> 12.7	> 11.4	> 11.5	> 12.0	> 10.4	> 9.5	> 9.3	> 8.6	> 8.5	> 8.0
~ Submission 2019	> 15.2	> 14.6	> 13.7	> 12.9	> 11.9	> 12.1	> 12.9	> 10.8	> 9.6	> 10.0	> 9.5	> 9.5	> 8.9
~ absolute change	> -0.01	> -0.09	> -0.14	> -0.26	> -0.45	> -0.61	> -0.87	> -0.38	> -0.09	> -0.73	> -0.89	> -1.03	> -0.87
~ relative change	> -0.08%	> -0.62%	> -1.01%	> -2.03%	> -3.75%	> -5.07%	> -6.72%	> -3.48%	> -0.95%	> -7.25%	> -9.31%	> -10.79%	> -9.73%
< Carbon monoxide - CO													
~ Submission 2020	> 162	> 152	> 141	> 134	> 123	> 121	> 121	> 105	> 101	> 98.9	> 94.7	> 93.3	> 92.6
~ Submission 2019	> 162	> 153	> 142	> 136	> 129	> 129	> 129	> 109	> 104	> 104	> 101	> 98.1	> 94.8
~ absolute change	> -0.09	> -0.73	> -1.08	> -2.26	> -6.12	> -8.14	> -8.30	> -3.77	> -2.33	> -4.92	> -5.81	> -4.83	> -2.26
~ relative change	> -0.05%	> -0.48%	> -0.76%	> -1.66%	> -4.75%	> -6.31%	> -6.42%	> -3.46%	> -2.24%	> -4.74%	> -5.78%	> -4.93%	> -2.38%



For more information on **recalculated emission estimates for Base Year and 2018**, please see the pollutant-specific recalculation tables following chapter [8.1 - Recalculations](#).

Uncertainties

Uncertainty estimates for **activity data** of mobile sources derive from research project FKZ 360 16 023 (title: "Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland") carried out by Knörr et al. (2009) ¹¹⁾.

Planned improvements

Besides the scheduled **routine revision** of TREMOD, no further improvements are planned for the next annual submission.

FAQs

Why are similar EF applied for estimating exhaust heavy metal emissions from both fossil and biofuels?

The EF provided in ¹²⁾ represent summatory values for (i) the fuel's and (ii) the lubricant's heavy-metal content as well as (iii) engine wear. Here, there might be no heavy metals contained in the biofuels. But since the specific shares of (i), (ii) and (iii) cannot be separated, and since the contributions of lubricant and engine wear might be dominant, the same emission factors are applied to biodiesel.

bibliography : 1 : AGEb (2019): Working Group on Energy Balances (Arbeitsgemeinschaft Energiebilanzen (Hrsg.), AGEb): Energiebilanz für die Bundesrepublik Deutschland; URL: <https://ag-energiebilanzen.de/7-0-Bilanzen-1990-2017.html>, Köln & Berlin, 2019. : 2 : MWV (2019): Association of the German Petroleum Industry (Mineralölwirtschaftsverband, MWV): Annual Report 2018, page 65, Table 'Sektoraler Verbrauch von Dieselkraftstoff 2012-2016'; URL: https://www.mwv.de/wp-content/uploads/2016/06/180830_MWV_Jahresbericht-2018_RZ_Web_es_small.pdf, Berlin, 2019. : 3 : Hedel, R., & Kunze, J. (2012): Recherche des jährlichen Kohleeinsatzes in historischen Schienenfahrzeugen seit 1990. Probst & Consorten Marketing-Beratung. Dresden, 2012. : 4 : Illichmann, S. (2016): Recherche des Festbrennstoffeinsatzes historischer Schienenfahrzeuge in Deutschland 2015, Probst & Consorten Marketing-Beratung. Study carried out for UBA; FKZ 363 01 392; not yet published; Dresden, 2016. : 5 : Knörr et al. (2019a): Knörr, W., Heidt, C., Gores, S., & Bergk, F.: ifeu Institute for Energy and Environmental Research (Institut für Energie- und Umweltforschung Heidelberg gGmbH, ifeu): Fortschreibung des Daten- und Rechenmodells: Energieverbrauch und Schadstoffemissionen des motorisierten Verkehrs in Deutschland 1960-2035, sowie TREMOD 5.81, im Auftrag des Umweltbundesamtes, Heidelberg & Berlin, 2019. : 6 : EMEP/EEA (2019): EMEP/EEA air pollutant emission inventory guidebook 2019, <https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-b-sectoral-guidance-chapters/1-energy/1-a-combustion/1-a-3-c-railways/view>; Copenhagen, 2019. : 7 : Rentz et al. (2008): Nationaler Durchführungsplan unter dem Stockholmer Abkommen zu persistenten organischen Schadstoffen (POPs), im Auftrag des Umweltbundesamtes, FKZ 205 67 444, UBA Texte | 01/2008, January 2008 - URL: <http://www.umweltbundesamt.de/en/publikationen/nationaler-durchfuehrungsplan-unter-stockholmer> : 7 : Knörr et al. (2009): Knörr, W., Heldstab, J., & Kasser, F.: Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland; final report; URL: <https://www.umweltbundesamt.de/sites/default/files/medien/461/publikationen/3937.pdf>, FKZ 360 16 023, Heidelberg & Zürich, 2009. **bibliography**

¹⁾ (bibcite 1)

²⁾ (bibcite 2)

³⁾ (bibcite 3)

- ⁴⁾ (bibcite 4)
- ⁵⁾ (bibcite 5)
- ⁶⁾ (bibcite 4)
- ⁷⁾ (bibcite 6)
- ⁸⁾ (bibcite 7)
- ⁹⁾ (bibcite 5)
- ¹⁰⁾ (bibcite 5)
- ¹¹⁾ (bibcite 7)
- ¹²⁾ (bibcite 5)