# **2.G.** - Use of Charcoal for barbecues

In sub-category NFR 2.G.4( c) - Charcoal TSP,  $PM_{10}$  and  $PM_{2.5}$  emissions from charcoal used for barbecue are reported.

Method	AD E	F	Key Categ	gory	
T1	NS [	D For <b><u>2.G.</u> L:</b> TSP, Cd, PM <sub>10</sub> / <b>L &amp; T:</b> PM <sub>2.5</sub> , Pb			
<b>T</b> = key	sourc	e by Trend <b>L</b> =	key source	by Level	
Method	ls				
D			Defau	Default	
RA		Refere	Reference Approach		
T1		Tier 1	Tier 1 / Simple Methodology *		
T2		Tier 2 <sup>3</sup>	Tier 2*		
Т3		Tier 3	Tier 3 / Detailed Methodology *		
C		CORIN	CORINAIR		
CS Cour			Count	ry Specific	
M Model			Model		
1		in the EMEP/C	ORINAIR En	nission Inventory Guidebook - 2007, in the group specific	
chapter					
AD - Data Source for Activity Data			vity Data		
NS National Statistics					
RS Regional Statistics					
IS International Statistics				-	
PS Plant Specific data					
<ul><li>AS Associations, business organisations</li><li>Q specific questionnaires, surveys</li></ul>			-		
	-		surveys		
		n Factors			
		MEP Guidebook	()		
C Conf		-	_		
CS Cour		pecific			
		ific data			

# Methodology

#### Activity data

The annual charcoal consumption for barbecue is calculated as annual import + production - export, and the relevant volumes of charcoal are extracted from national statistics by the Federal Statistical Office. Other applications for charcoal are not included.

The model is based on the two assumptions that there is no storage of charcoal and that all charcoal

is burned.

The amount of charcoal used for barbecue has been ever-expanding from 1990 to 2012 and is predominantly imported. As there is only one big producer, produced amounts and resulting emissions are confidential.

#### **Emission factors**

The emission factors are derived from the CEPMEIP Database (SNAP: 060508).<sup>1)</sup>

#### Uncertainties

The uncertainties of emissions are 54% for the lower and upper bounds.

### **Recalculations**

The import and export data for 2018 were changed as revised activity data for the foreign sale was available from the Federal Statistical Office.



For specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

## **Planned improvements**

No improvements are planned.

1)

CEPMEIP, 2018: Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP), CEPMEIP Database, SNAP code : 060508; URL: http://www.air.sk/tno/cepmeip/em\_factors.php?PHPSESSID=cc235582eb4e09bf725d6f859deb382d