# **2.G.** - Use of Charcoal for barbecues

In sub-category NFR 2.G. - Use of Charcoal for barbecues TSP,  $PM_{10}$  and  $PM_{2.5}$  emissions from charcoal used for barbecue are reported.

u	AD EI	- Ke	y Category
T1	NS D	For <b><u>2.G.</u> L:</b> TSP,	Cd, PM <sub>10</sub> / <b>L &amp; T:</b> PM <sub>2.5</sub> , Pb
<b>T</b> = key	source	by Trend <b>L</b> = key	v source by Level
Method	ls		
		D	Default
		RA	Reference Approach
		Т1	Tier 1 / Simple Methodology *
		T2	Tier 2*
		Т3	Tier 3 / Detailed Methodology *
		C	CORINAIR
		CS	Country Specific
		М	Model
		in the EMEP/CORI	NAIR Emission Inventory Guidebook - 2007, in the group specific
chapters			
		urce for Activity	Data
NS Natio			
	onal St		
<b>RS</b> Regi			
IS Inter	nation	al Statistics	
IS Inter PS Plan	nation t Speci	al Statistics fic data	
IS Inter PS Plan AS Asso	nation t Speci ociatior	al Statistics fic data is, business organ	
IS Inter PS Plan AS Asso	nation t Speci ociatior	al Statistics fic data	
IS Inter PS Plan AS Asso Q spec	nation t Speci ociation ific qu	al Statistics fic data is, business organ	
IS Inter PS Plan AS Asso Q spec EF - Em	nation t Speci ciatior ific qu <b>ission</b>	al Statistics fic data is, business organ estionnaires, surv	
IS Inter PS Plan AS Asso Q spec EF - Em	nation t Speci ciatior ific qu <b>ission</b> ult (EM	al Statistics fic data is, business organ estionnaires, surv Factors IEP Guidebook)	
IS Inter PS Plan AS Asso Q spec EF - Em D Defa	nation t Speci ciatior ific qu <b>ission</b> ult (EM identia	al Statistics fic data ns, business organ estionnaires, surv <b>Factors</b> IEP Guidebook)	

# Methodology

#### Activity data

The annual charcoal consumption for barbecue is calculated as annual import + production - export, and the relevant volumes of charcoal are extracted from national statistics by the Federal Statistical Office. Other applications for charcoal are not included.

The model is based on the two assumptions that there is no storage of charcoal and that all charcoal

is burned.

The amount of charcoal used for barbecue has been ever-expanding from 1990 to 2012 and is predominantly imported. As there is only one big producer, produced amounts and resulting emissions are confidential.

#### **Emission factors**

The emission factors are derived from the CEPMEIP Database (SNAP: 060508).<sup>1)</sup>

#### Uncertainties

The uncertainties of emissions are 54% for the lower and upper bounds.

### **Recalculations**

The import and export data for 2018 were changed as revised activity data for the foreign sale was available from the Federal Statistical Office.



For specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

## **Planned improvements**

No improvements are planned.

1)

CEPMEIP, 2018: Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP), CEPMEIP Database, SNAP code : 060508; URL: http://www.air.sk/tno/cepmeip/em\_factors.php?PHPSESSID=cc235582eb4e09bf725d6f859deb382d