2.G. - Use of Charcoal for barbecues

In sub-category NFR 2.G. - Use of Charcoal for barbecues TSP, PM_{10} and $PM_{2.5}$ emissions from charcoal used for barbecue are reported.

Method	AD I	EF	Key Ca	tegory		
T1	NS	D	For <u>2.G.</u> L: Cd, Pl	M ₁₀ / L &	T: PM _{2.5}	
T = key	souro	ce b	oy Trend L = key	source b	by Level	
Method	ls					
D			D	Default		
RA			RA	Reference Approach		
T1			F1	Tier 1 / Simple Methodology *		
T2			۲ 2	Tier 2*		
Т3			F3	Tier 3 / Detailed Methodology *		
C			с	CORINAIR		
CS Count			CS	Country	y Specific	
M Mode				Model		
1		d ir	h the EMEP/CORIN	IAIR Emis	ission Inventory Guidebook - 2007, in the group specific	
chapters			a a fa u A atia ita a	Data		
AD - Data Source for Activity Da			-	Data		
NS National Statistics						
RS Regional Statistics						
PS Plant Specific data AS Associations, business organisations				sations		
Q specific questionnaires, surveys						
EF - Em				ys		
			P Guidebook)			
C Conf		-	-: <i>f</i> i -			
CS Cour		· .				
PS Plant	c Spe	CITIC	c data			

Method

Activity data

The annual charcoal consumption for barbecue is calculated as annual import + production - export, and the relevant volumes of charcoal are extracted from national statistics by the Federal Statistical Office. Other applications for charcoal are not included.

The model is based on the two assumptions that there is no storage of charcoal and that all charcoal

is burned.

The amount of charcoal used for barbecue has been ever-expanding from 1990 to 2012 and is predominantly imported. As there is only one big producer, produced amounts and resulting emissions are confidential.

Emission factors

The emission factors are derived from the CEPMEIP Database (SNAP: 060508).¹⁾

Uncertainties

The uncertainties of emissions are 54% for the lower and upper bounds.

Recalculations

The import and export data for 2018 were changed as revised activity data for the foreign sale was available from the Federal Statistical Office.



For specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

Planned improvements

No improvements are planned.

1)

CEPMEIP, 2018: Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP), CEPMEIP Database, SNAP code : 060508; URL: http://www.air.sk/tno/cepmeip/em_factors.php?PHPSESSID=cc235582eb4e09bf725d6f859deb382d