

11.B - Forest Fires

Short Description

Method	AD	EF	Key Category
CS, T2, T1	CS	D	<i>not included in key category analysis</i>

Category 11.B - Forest fires include emissions from forest fires occurring naturally or caused by humans.

In Germany's forests prescribed burning is not applied. Therefore, all forest fires are categorized as wildfires. - Note that emissions reported here are not accounted for the national totals.

Methodology

For calculating the emissions of wildfires a country specific Tier2 approach was used. The mass of carbon emitted $M(C)$ was calculated using the adapted equation follows the methodology of Seiler and Crutzen (1980) ¹⁾.



$$M(C) = 0.45 * A * B * \beta$$

where:

0.45 = average fraction of carbon in fuel wood;

A = forest area burnt in [m²];

B = mean above-ground biomass of fuel material per unit area in [kg/m²];

β = burning efficiency (fraction burnt) of the above-ground biomass.

The data on forest areas burnt for the period 1990 to 2018 have been taken from the German forest fire statistic (BLE, 2020) ²⁾ managed by the Federal Agency for Agriculture and Food. The mean above-

ground biomass was derived for each year by linear extrapolation and interpolation between the

- German National Forest Inventories of 1987, 2002, 2012 (Bundeswaldinventuren 1987, 2002, 2012),
- the inventory study 2008 ³⁾ and,
- the carbon inventory 2017 (Kohlenstoffinventur 2017) ⁴⁾.

Pursuant to König (2007) ⁵⁾, 80% of the forest fires in Germany are surface fires and 20% crown fires. In accordance to the IPCC Good Practice Guidance for LULUCF (2003) a burning efficiency of 0.15 was used for surface fires and an efficiency of 0.45 was used for crown fires.

The emissions for the pollutants were calculated by multiplying the mass of carbon with the respective emission factors from table 3-3 (EMEP/EEA, 2019) ⁶⁾.

For the calculation of particulate emissions (TSP, PM₁₀ and PM_{2.5}) the burnt biomass was multiplied with the respective emission factors from table 3-5 (EMEP/EEA, 2019). Those particulate emission factors have been estimated by averaging the emission factors from the US Environmental Protection Agency (USEPA, 1996) ⁷⁾ methodology, since no better information is available. Those emission factors are assumed to be the same for all types of forest.

The Guidebook does not indicate whether EFs have considered the condensable component (with or without).

As a first estimate black carbon emissions are reported the first time.

Activity data

The data on forest areas burnt for the period 1990 to 2019 are based on the German forest fire statistic (BLE, 2020) managed by the Federal Agency for Agriculture and Food.

Table 1: Area of forest burnt from 1990 until the latest reporting year, in [ha]

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Area of forest burnt	1606	920	4908	1493	1114	592	1381	599	397	415	581
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Area of forest burnt	122	122	1315	274	183	482	256	539	757	522	214
	2012	2013	2014	2015	2016	2017	2018	2019			
Area of forest burnt	269	199	120	526	283	395	2349	2711			

Emission factors

For the year 2019 the estimated emission factors from table 2 were applied.

Table 2: Emission factors applied for 2019

Pollutant	EF ₂₀₁₉ [kg/ha forest area burnt]
NO _x	123.63
CO	4,409.60

Pollutant	EF₂₀₁₉ [kg/ha forest area burnt]
NMVOC	389.45
SO _x	29.67
NH ₃	33.38
TSP	700.59
PM ₁₀	453.32
PM _{2.5}	370.90
BC	33.38

In addition, a large-scale fire, which occurred in September 2018, is reported under 11.B. A detailed description can be found in the NIR 2020 in Chapter 6.8.2.5 ⁸⁾, because a large amount of CO₂ emissions were released.

The burned area of the drained moor, which is used as a military facility, covered 1,221 ha. This fire was extensively investigated and documented by the Federal Office for Infrastructure, Environmental Protection and Services of the German Armed Forces. The emissions are calculated according to IPCC GL (2006), chapter 2, form 2.27 ⁹⁾. The product MB×Cf is set to 336 t dm ha⁻¹ according to Table 2.6 and formula 2.7 2013 IPCC Wetlands Supplement ¹⁰⁾, i.e. it is assumed that the moor was completely drained during the fire.

For the calculation of CO emissions the EF according to Table 2.7, 2013 IPCC Wetlands Supplement 207 g (kg dm)⁻¹, is taken into account. This results in 85 kt CO. For other emissions from land fires on drained organic soils no Tier-1 emission factors exist and are therefore not reported (NO)".

¹⁾

Seiler, W., Crutzen P.J.: Estimates of gross and net fluxes of carbon between the biosphere and the atmosphere from biomass burning. Climate Change. 1980

²⁾

BLE (Bundesanstalt für Landwirtschaft und Ernährung), 2020: Waldbrandstatistik der Bundesrepublik Deutschland für das Jahr 2019, Bonn: 21 S.

³⁾

[<https://www.thuenen.de/en/wo/projects/forest-monitoring/greenhouse-gas-inventory-for-forests/inventory-study-2008/> Inventurstudie 2008]

⁴⁾

[<https://www.thuenen.de/en/wo/projects/forest-monitoring/greenhouse-gas-inventory-for-forests/carbon-inventory-2017>]

⁵⁾

König, H.-C., 2007. Waldbrandschutz - Kompendium für Forst und Feuerwehr. 1. Fachverlag Matthias Grimm, Berlin, 197 S.

⁶⁾

EMEP/EEA, 2019:

[*<https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-b-sectoral-guidance-chapters/11-natural-sources/11-b-forest-fires/view>]

⁷⁾

USEPA, 1996: Compilation of Air Pollutant Emission Factors Vol.1. Stationary, Point and Area Sources. Report AP-42 (fifth edition)

⁸⁾

NIR (2020): National Inventory Report 2020 for the German Greenhouse Gas Inventory 1990-2018. Available in April 2020

⁹⁾

IPCC (Intergovernmental Panel on Climate Change) (2006): 2006 IPCC Guidelines for National

Greenhouse Gas Inventories, Volume 4: Agriculture, Forestry and Other Land Use. Eds.: Eggleston S., Buendia L., Miwa K., Ngara T., Tanabe K. (Eds). IEA/OECD, IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

<<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>>

¹⁰⁾

IPCC (Intergovernmental Panel on Climate Change) (2014b): 2013 Supplement to the IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G.(eds). Published: IPCC, Switzerland

<<http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html>>