

## 5.C.2 - Open Burning of Waste

Category Code	Method					AD					EF				
5.C.2	CS					Q					D, CS				
Key Category	SO <sub>2</sub>	NO <sub>x</sub>	NH <sub>3</sub>	NMVOC	CO	BC	Pb	Hg	Cd	Diox	PAH	HCB	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
5.C.2	-/-	-/-	-	-/-	-/-	-/-	-/-	-	-/-	-/-	-/-	-	-/-	L/-	L/-

**T** = key source by Trend **L** = key source by Level

### Methods

<b>D</b>	Default
<b>RA</b>	Reference Approach
<b>T1</b>	Tier 1 / Simple Methodology *
<b>T2</b>	Tier 2*
<b>T3</b>	Tier 3 / Detailed Methodology *
<b>C</b>	CORINAIR
<b>CS</b>	Country Specific
<b>M</b>	Model

\* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

### AD - Data Source for Activity Data

<b>NS</b>	National Statistics
<b>RS</b>	Regional Statistics
<b>IS</b>	International Statistics
<b>PS</b>	Plant Specific data
<b>AS</b>	Associations, business organisations
<b>Q</b>	specific questionnaires, surveys

### EF - Emission Factors

<b>D</b>	Default (EMEP Guidebook)
<b>C</b>	Confidential
<b>CS</b>	Country Specific
<b>PS</b>	Plant Specific data

Within NFR sub-category 5.C.2 - Open Burning of Waste, the German emissions inventory provides emissions from registered bonfires and other wooden materials burnt outdoors. Emissions from bonfires are key source for PM<sub>2.5</sub> and PM<sub>10</sub>, but in principle of minor priority due to discontinuous appearance.

Emissions from open burning of wood and green waste for traditional purposes, so-called bonfires such as Easter fires, are reported model-based. In addition to biogenic carbon dioxide, emissions of NO<sub>x</sub>, SO<sub>2</sub>, CO, NMVOC, particulate matter (PM<sub>2.5</sub>, PM<sub>10</sub> and TSP) and Polycyclic Aromatic Hydrocarbons (PAHs) are covered so far.

## Method

For developing of a estimation frame a survey regarding the number of such bonfires was carried out by an expert work (Wagner & Steinmetzer, 2018) [1]. As the result, questionnaires from municipalities and statistical projections for Germany for the year 2016 were checked. The project has shown a declining trend since 1990. On the basis of expert judgement, a further reduction of emissions in the future is expected.

## Activity data

Activity data for this category are based on data from a step by step calculation: After the evaluation of the questionnaires an extrapolation of the volume and the number of bonfires was made for Germany. The median values of clusters of city-sizes were used for the calculation, resulting in the following values <sup>1)</sup>:

fire	resulting number	resulting quantity in kt of wooden wastes
easter fires et.	54	343.3
other open burning of wood	49	59.3

## Emission factors

Emission factors used were taken from different sources:


pollutant	figure	reference
CO	58.0	GB 2016 small combustion Table 3-6: Tier 1 emission factors for NFR source category 1.A.4.b, using biomass
NO <sub>x</sub>	0.9	Research results for charcoal
SO <sub>2</sub>	0.2	Research results for charcoal
NM <sub>VOC</sub>	47.0	GB 2016 Forest fires, table 3-1, according 'wood burned'
TSP	17.0	GB 2016 Forest fires, table 3-1, according 'wood burned'
PM <sub>10</sub>	11.0	GB 2016 Forest fires, table 3-1, according 'wood burned'
PM <sub>2.5</sub>	9.0	GB 2016 Forest fires, table 3-1, according 'wood burned'
BC	0.81	GB 2016 Forest fires, table 3-1, according 'wood burned'
PCDD/F	10.0 µg/ t	GB 2016 Forest fires, table 3-2
PAH	0.00339	sum of single compounds
BaP	0.0013	Research results for charcoal
BbF	0.0015	Research results for charcoal
BkF	0.0005	Research results for charcoal
IxP	0.00009	Research results for charcoal
Pb	0.32 g/ t	GB 2016 Forest fires, table 3-2
Cd	0.13 g/ t	GB 2016 Forest fires, table 3-2

## Trends in emissions

All trends in emissions correspond to trends of AD. No rising trends are to identify. In 2019, there were many bans on open fires due to increased forest fire danger.

## trends of emissions of bonfires

Emissions by pollutant / Emissionen nach Schadstoff



\* Base Year for PM = 1995 / Basisjahr für Feinstäube (PM) ist 1995

German Emission Inventory (12.02.2021)

### Emission trends of bonfires

## Recalculations

With **activity data** and **emission factors** remaining unrevised, no recalculations have been carried out compared to last year's submission.



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following [chapter 8.1 - Recalculations](#).

<sup>1)</sup> Wagner & Steinmetzer, 2018: Jörg Wagner, Sonja Steinmetzer, INTECUS GmbH Abfallwirtschaft und

umweltintegratives Management: Erhebung der Größen und Zusammensetzung von Brauchtums- und Lagerfeuern durch kommunale Befragungen; URL:  
[https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2018-02-19\\_texte\\_11-2018\\_lager-brauchtumsfeuer.pdf](https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2018-02-19_texte_11-2018_lager-brauchtumsfeuer.pdf); UBA-Texte 11/2018