

Explanation of Key Trends - Ammonia

Obligations

Germany has made a commitment under the Gothenburg Protocol to reduce ammonia emissions. Since 2010, it is no longer permissible to exceed a National Emission Ceiling of 550kt NH₃ for Germany as whole. The revised Gothenburg Protocol and the revised NEC Directive both define emission reduction targets relative to a 2005 base year, mandating 5% (2020) and 29% (2030) reductions respectively.

Further details can be found in [Chapter 9 - Projections](#) and [Chapter 11 - Adjustments and Emission Ceiling Exceedance](#).

Main drivers


The Main Drivers for NH₃ emissions are agricultural emissions from **Manure Management (NFR 4.B)** with 42% of total 1990 emissions and a 20% reduction between 1990-2019 and **Agricultural Soils (NFR 4.D)** with even 53% of total 1990 emissions and a 16% decrease between 1990-2018. The overall emission trend mainly follows the agricultural emissions closely with a total reduction of 18% between 1990 and 2019. The decrease of NH₃ emission in the year 1991 is due to a reduced livestock population that followed after the German reunification, while no explicit trend is discernible for the years up to 2016. Between 2016 and 2019 there is a slight reduction of about 8%, but it has to be seen how stable this trend will be.

NH₃ Emissions 1990-2019

| Total Emissions (kt) | | | | | | | | | | | | | | Trend: latest compared to | |
|----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------------------------|------------|
| 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 1990 | last years |
| 715 | 613 | 627 | 607 | 619 | 625 | 631 | 637 | 645 | 641 | 638 | 624 | 601 | 587 | ▼ -18.0% | ▼ |

Ammonia / Ammoniak

Emissions per Sector / Sektorale Emissionen



Quelle: German Emission Inventory (01.02.2022)

NH₃ trend by sector