

3.B - Manure Management

Short description

| NFR-Code | Name of Category | Method | AD | EF | State of reporting |
|--|--------------------------|--|--------|--|--|
| 3.B | Manure Management | see sub-category details | | | |
| consisting of / including source categories | | | | | |
| 3.B.1.a & 3.B.1.b | Cattle | T3 (NH ₃), T2 (NO _x , TSP, PM ₁₀ , PM _{2.5} , NMVOC) | NS, RS | CS (NH ₃ , NO _x), D (TSP, PM ₁₀ , PM _{2.5} , NMVOC) | L: NH ₃ (for 3.B.1.a) |
| 3.B.2, 3.B.4.d, 3.B.4.e | Sheep, Goats, Horses | T2 (NH ₃ , NO _x , TSP, PM ₁₀ , PM _{2.5}), T1 (NMVOC) | NS, RS | CS (NH ₃ , NO _x), D (TSP, PM ₁₀ , PM _{2.5} , NMVOC) | |
| 3.B.3 | Swine | T3 (NH ₃), T2 (NO _x , TSP, PM ₁₀ , PM _{2.5}), T1 (NMVOC) | NS, RS | CS (NH ₃ , NO _x), D (TSP, PM ₁₀ , PM _{2.5} , NMVOC) | |
| 3.B.4.a | Buffalo | | | | NO, from 1990 until 1995, since 1996 IE, considered in 3.B.1.b |
| 3.B.4.f | Mules and asses | | | | IE, considered in 3.B.4.e |
| 3.B.4.g i-iv | Poultry | T2 (NH ₃ , NO _x , TSP, PM ₁₀ , PM _{2.5}), T1 (NMVOC) | NS, RS | CS (NH ₃ , NO _x), D (TSP, PM ₁₀ , PM _{2.5} , NMVOC) | T: NH ₃ (for 3.B.4.g iii) |
| 3.B.4.h | Other animals | | | | NE |

| Key Category | NO _x | NMVOC | SO ₂ | NH ₃ | PM _{2.5} | PM ₁₀ | TSP | BC | CO | Pb | Cd | Hg | Diox | PAH | HCB |
|--------------|-----------------|-------|-----------------|-----------------|-------------------|------------------|-----|----|----|----|----|----|------|-----|-----|
| 3.B.1.a | -/- | L/- | - | L/- | L/- | L/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.1.b | -/- | L/- | - | L/T | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.2 | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.3 | -/- | -/- | - | L/T | -/- | -/- | L/- | - | - | - | - | - | - | - | - |
| 3.B.4.d | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.4.e | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.4.g.i | -/- | -/- | - | -/- | -/- | -/- | L/- | - | - | - | - | - | - | - | - |
| 3.B.4.g.ii | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.4.g.iii | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |
| 3.B.4.g.iv | -/- | -/- | - | -/- | -/- | -/- | -/- | - | - | - | - | - | - | - | - |

T = key source by Trend L = key source by Level

| Methods | |
|-----------|---------------------------------|
| D | Default |
| RA | Reference Approach |
| T1 | Tier 1 / Simple Methodology * |
| T2 | Tier 2* |
| T3 | Tier 3 / Detailed Methodology * |
| C | CORINAIR |
| CS | Country Specific |
| M | Model |

* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

| AD - Data Source for Activity Data | |
|------------------------------------|--------------------------------------|
| NS | National Statistics |
| RS | Regional Statistics |
| IS | International Statistics |
| PS | Plant Specific data |
| AS | Associations, business organisations |
| Q | specific questionnaires, surveys |
| EF - Emission Factors | |
| D | Default (EMEP Guidebook) |
| C | Confidential |

| |
|-------------------------------|
| EF - Emission Factors |
| CS Country Specific |
| PS Plant Specific data |

Country specifics



In 2020, NH_3 emissions from category 3.B (manure management) were 48.6 % from total agricultural emissions, which is equal to ~ 249.2 kt NH_3 . Within those emissions 52.0 % originate from cattle manure (~ 129.5 kt), 34.0 % from pig manure (ca. 84.7 kt), and 10.8 % from poultry manure (~ 27.0 kt). Calculations take into account the impact of anaerobic digestion of manure on the emissions.

NO_x emissions from category 3.B (manure management) contribute only 1.3 % (~ 1.4 kt) to the total agricultural NO_x emissions. They are calculated proportionally to N_2O emissions, see Vos et al. (2022) ¹⁾.

NM VOC emissions from category 3.B (manure management) contributed 96.9 % (289.8 kt) from total agricultural NM VOC emissions (298.9 kt).

In 2020, manure management contributed, respectively, 71.3 % (43.0 kt), 42.8 % (12.9 kt) and 84.8 % (3.7 kt) to the total agricultural TSP, PM_{10} and $\text{PM}_{2.5}$ emissions (TSP: 60.3 kt, PM_{10} : 30.2 kt, $\text{PM}_{2.5}$: 4.4 kt, respectively).

Activity data for all pollutants

The Federal Statistical Agency and the Statistical Agencies of the federal states carry out surveys in order to collect, along with other data, the head counts of animals. The results of these surveys are used for emission calculations, for details see Vos et al, 2022, Chapter 3.4.2.

The animal population figures used in the inventory are presented in Table 1. From 2017 to 2019 the animal population figures for horses, goats and poultry differ from the figures presented in the last IIR. In the last year's submission these figures were extrapolated from the year 2016. Now there are new figures available for the year 2020 and the years 2017-2019 have been interpolated. Buffaloes are included in the cattle population figures, mules and asses are included in the horse population figures (IE), see Vos et al. (2022), Chapters 4.1 and 7.1. In the first years after the German reunification in 1990 animal livestock decreased markedly. The head counts for cattle continued to decrease significantly until 2006/2007, followed by a more or less stable period until 2014. Since 2015 a slight decrease occurred. In 2020, dairy cattle numbers are 61.7 % of 1990 numbers, while the total population of other cattle is at 56.2 % of 1990. Swine numbers decreased until 1995 and then increased slightly. Since 2014 a slight decrease occurred (2020: 81.6 % of 1990). The 2020 numbers of horses, sheep and goats are, respectively, at 90,8 %, 54.5 % and 172.1 % of 1990.

Figures for broilers and turkeys are showing a massive increase since 1990. Since the year 2013, there have been only minor changes of total poultry numbers. In total, 2020 poultry population figures are at 152.0 % of 1990. A detailed description of the animal numbers used can be found in the National Inventory Report 2022 ²⁾, Chapter 5.1.3.2.3.

Table 1: Population of animals

| Population of animals (in 1000) | | | | | | | | | | | | | | | |
|-------------------------------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| dairy cattle | 6'354.6 | 5'229.4 | 4'569.8 | 4'236.4 | 4'183.1 | 4'190.1 | 4'190.5 | 4'267.6 | 4'295.7 | 4'284.6 | 4'217.7 | 4'199.0 | 4'100.9 | 4'011.7 | 3'921.4 |
| other cattle | 13'133.4 | 10'660.5 | 9'968.9 | 8'800.4 | 8'628.7 | 8'340.4 | 8'319.1 | 8'418.4 | 8'446.5 | 8'350.8 | 8'248.9 | 8'082.2 | 7'848.2 | 7'627.9 | 7'380.5 |
| buffalo | NO | NO | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE |
| mules and asses | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE | IE |
| horses | 499.5 | 634.1 | 499.5 | 508.4 | 461.8 | 461.6 | 461.5 | 461.3 | 454.9 | 448.4 | 442.0 | 444.9 | 447.8 | 450.7 | 453.7 |
| sheep | 3'266.1 | 2'990.7 | 2'743.3 | 2'643.1 | 2'245.0 | 1'979.7 | 1'965.9 | 1'877.2 | 1'892.4 | 1'866.9 | 1'851.0 | 1'863.2 | 1'846.0 | 1'813.6 | 1'780.3 |
| goats | 90.0 | 100.0 | 140.0 | 170.0 | 149.9 | 143.4 | 136.8 | 130.2 | 133.1 | 135.9 | 138.8 | 142.8 | 146.9 | 150.9 | 154.9 |
| swine | 26'502.5 | 20'387.3 | 21'767.7 | 22'742.8 | 22'244.4 | 22'787.9 | 23'648.3 | 23'391.2 | 23'666.9 | 22'978.5 | 22'761.2 | 22'920.8 | 22'019.2 | 21'596.4 | 21'622.0 |
| laying hens | 53'450.5 | 45'317.3 | 44'225.6 | 38'203.6 | 35'279.0 | 39'514.9 | 43'750.8 | 47'986.7 | 49'303.0 | 50'619.3 | 51'935.5 | 52'571.1 | 53'206.6 | 53'842.1 | 54'477.6 |
| broilers | 35'393.0 | 42'025.8 | 50'359.9 | 56'762.5 | 67'531.1 | 77'402.6 | 87'274.1 | 97'145.6 | 96'027.5 | 94'909.4 | 93'791.3 | 93'458.7 | 93'126.1 | 92'793.5 | 92'461.0 |
| turkeys | 5'029.2 | 6'742.0 | 8'893.1 | 10'611.1 | 11'344.0 | 11'981.2 | 12'618.5 | 13'255.7 | 12'957.1 | 12'658.5 | 12'359.9 | 12'164.7 | 11'969.5 | 11'774.3 | 11'579.1 |
| pullets | 17'210.8 | 14'592.0 | 14'240.5 | 12'301.4 | 11'303.3 | 12'749.3 | 14'195.2 | 15'641.2 | 14'734.7 | 13'828.3 | 12'921.8 | 12'736.3 | 12'550.7 | 12'365.1 | 12'179.6 |
| ducks | 2'013.7 | 1'933.7 | 2'055.7 | 2'352.2 | 3'164.3 | 3'029.5 | 2'894.6 | 2'759.7 | 2'585.3 | 2'410.8 | 2'236.4 | 2'209.1 | 2'181.9 | 2'154.6 | 2'127.4 |
| geese | 781.5 | 617.0 | 404.8 | 329.5 | 278.1 | 366.8 | 455.5 | 544.2 | 472.5 | 400.8 | 329.0 | 327.7 | 326.3 | 324.9 | 323.5 |
| other animals: no data available a) | | | | | | | | | | | | | | | |

a) Animal numbers of other animals are not available. Emissions of other animals were approximated with estimated population figures for a single year (see Rösemann et. al., 2017, Chapter 9, ³⁾ and submitted to the TERT of the NECD-Review. The TERT confirmed that emissions are below the threshold of significance. For GHG emission reporting the UNFCCC has acknowledged that the emissions from Germany's other animals are negligible. To ensure consistency between UNFCCC and UNECE/NEC reporting, no air pollutants from other animals are reported.

Additional data

Emission calculations in accordance with a Tier 2 or Tier 3 method require data on animal performance (animal weight, weight gain, milk yield, milk protein content, milk fat content, numbers of births, numbers of eggs and weights of eggs) and on the relevant feeding details (phase feeding, feed components, protein and energy content, digestibility and feed efficiency). To subdivide officially recorded total numbers of turkeys into roosters and hens, the respective population percentages need to be known. Details on data requirements for the modelling of emissions from livestock husbandry in the German inventory can be found in Voss et al. (2022), Chapters 4 to 8.

Most of the data mentioned above is not available from official statistics and was obtained from literature, from publications by agricultural associations, from regulations for agricultural consulting in Germany and from expert judgments. For 1991, 1995 and 1999, frequency distributions of feeding strategies, husbandry systems (shares of pasturing/stabling; shares of various housing methods), storage types as well as techniques of farm manure spreading were obtained with the help of the RAUMIS agricultural sector model (Regionalisiertes Agrar- und UmweltInformationssystem für Deutschland/ Regionalised agricultural and environmental information system for Germany). RAUMIS has been developed and is operated by the Institute of Rural Studies of the Thünen Institute (Federal Research Institute for Rural Areas, Forestry and Fisheries). For an introduction to RAUMIS see Weingarten (1995) ⁴⁾; a detailed description is provided in Henrichsmeyer et al. (1996) ⁵⁾.


RAUMIS did not model complete time series but only selected years. RAUMIS data for the years 1991, 1995, and 1999 are used in the inventory for years 1990 – 1993, 1994 – 1997, and 1998 – 1999, respectively. For the year 2010, respective data are used that were derived from the 2010 official agricultural census and the simultaneous survey of agricultural production methods (Landwirtschaftliche Zählung 2010, Statistisches Bundesamt/ Federal Statistical Office) as well as the 2011 survey on manure application practices (Erhebung über Wirtschaftsdüngerausbringung, Statistisches Bundesamt/ Federal Statistical Office).

For the year 2015, data on techniques of farm manure spreading from the 2016 official agricultural census (Agrarstrukturerhebung 2016, Statistisches Bundesamt / Federal Statistical Office) are used. The gaps between the latest

N mass flow and emission assessment

The calculation of the emissions of NH_3 , N_2O , NO_x and N_2 from German animal husbandry is based on the so-called N mass flow approach (e. g. Dämmgen and Hutchings, 2008⁷⁾). This approach differentiates between N excreted with faeces (organic nitrogen Norg, i. e. undigested feed N) and urine (total ammoniacal nitrogen TAN, i. e. fraction of feed N metabolized). The N flow within the manure management system is treated as depicted in the figure below. This method reconciles the requirements of both the Atmospheric Emission Inventory Guidebook for NH_3 emissions (EMEP, 2019), and the IPCC guidelines for greenhouse gas emissions (IPCC (2006)⁸⁾. Reidy et al. (2008),⁹⁾ showed for several European countries (Germany, the Netherlands, Switzerland, United Kingdom) that their N-flow based inventory models yielded, in spite of national peculiarities, comparable results as long as standardised data sets for the input variables were used.

Not explicitly shown in the N mass flow scheme is air scrubbing in housing and anaerobic digestion of manure. These issues are separately described further below. Note that emissions from grazing and application are reported in sector 3.D.



General scheme of N flows in animal husbandry

m: mass from which emissions may occur. Narrow broken arrows: TAN (total ammoniacal nitrogen); narrow continuous arrows: organic N. The horizontal arrows denote the process of immobilisation in systems with bedding occurring in the house, and the process of mineralisation during storage, which occurs in any case. Broad arrows denote N-emissions assigned to manure management (E_{yard} NH_3 emissions from yards; E_{house} NH_3 emissions from house; $E_{storage}$ NH_3 , N_2O , NO_x and N_2 emissions from storage; E_{applic} NH_3 emissions during and after spreading; E_{graz} NH_3 , N_2O , NO_x and N_2 emissions during and after grazing; E_{soil} N_2O , NO_x and N_2 emissions from soil resulting from manure input).

The model allows tracing of the pathways of the two N fractions after excretion. The various locations where excretion may take place are considered. The partial mass flows down to the input to soil are represented. During storage Norg can be transformed into TAN and vice versa. Both, the way and the magnitude of such transformations may be influenced by manure treatment processes like, e. g., anaerobic digestion where a considerable fraction of Norg is mineralized to TAN. For details see Vos et al. (2022), Chapters 3.3.4.3 and 3.3.4.4. Wherever NH_3 is emitted, its formation is related to the amount of the TAN present. N_2O emissions are related to the total amount of N available (Norg + TAN). NO_x emissions (i. e. NO emissions) are calculated proportionally to the N_2O emissions, see section 'Emission factors'. Note that the N_2O , NO_x and N_2 emissions from the various storage systems include the respective emissions from the related housing systems.

Air scrubber systems in swine and poultry housings

For pig and poultry production the inventory model considers the effect of air scrubbing. Data on frequencies of air scrubbing facilities and the removal efficiency are provided by KTBL (Kuratorium für Technik und Bauwesen in der Landwirtschaft / Association for Technology and Structures in Agriculture). The average removal efficiency of NH_3 is 80 % for swine and 70 % for poultry, while for TSP and PM_{10} the rates are set to 90 % and for $\text{PM}_{2.5}$ to 70 % for both animal categories. For swine two types of air scrubbers are distinguished: certified systems that remove both NH_3 and particles, and non-certified systems

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| sheep | 0.83 | 0.82 | 0.84 | 0.83 | 0.84 | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.82 | 0.83 | 0.83 |
| goats | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 | 1.62 |
| swine | 4.60 | 4.50 | 4.37 | 4.30 | 4.13 | 4.07 | 4.03 | 4.00 | 3.98 | 4.02 | 4.00 | 3.98 | 3.97 | 3.94 | 3.92 |
| laying hens | 0.212 | 0.203 | 0.199 | 0.198 | 0.142 | 0.144 | 0.144 | 0.145 | 0.146 | 0.146 | 0.147 | 0.147 | 0.148 | 0.149 | 0.150 |
| broilers | 0.143 | 0.120 | 0.128 | 0.131 | 0.128 | 0.118 | 0.103 | 0.092 | 0.094 | 0.094 | 0.094 | 0.093 | 0.094 | 0.092 | 0.089 |
| turkeys | 0.793 | 0.793 | 0.797 | 0.874 | 0.836 | 0.839 | 0.892 | 0.862 | 0.861 | 0.859 | 0.860 | 0.860 | 0.835 | 0.835 | 0.784 |
| pullets | 0.103 | 0.095 | 0.087 | 0.087 | 0.084 | 0.083 | 0.083 | 0.082 | 0.082 | 0.082 | 0.083 | 0.083 | 0.084 | 0.084 | 0.084 |
| ducks | 0.193 | 0.193 | 0.193 | 0.192 | 0.189 | 0.188 | 0.188 | 0.186 | 0.186 | 0.185 | 0.185 | 0.186 | 0.186 | 0.186 | 0.186 |
| geese | 0.384 | 0.384 | 0.384 | 0.383 | 0.380 | 0.380 | 0.380 | 0.379 | 0.379 | 0.378 | 0.378 | 0.378 | 0.378 | 0.378 | 0.378 |
| IEF in kg NO_x per animal place | | | | | | | | | | | | | | | |
| dairy cattle | 0.106 | 0.114 | 0.125 | 0.130 | 0.125 | 0.123 | 0.119 | 0.117 | 0.117 | 0.118 | 0.120 | 0.120 | 0.123 | 0.126 | 0.129 |
| other cattle | 0.053 | 0.057 | 0.059 | 0.062 | 0.064 | 0.063 | 0.063 | 0.063 | 0.064 | 0.064 | 0.065 | 0.066 | 0.066 | 0.068 | 0.069 |
| horses | 0.084 | 0.084 | 0.086 | 0.086 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.086 | 0.086 | 0.086 | 0.086 |
| sheep | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| goats | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 |
| swine | 0.011 | 0.013 | 0.012 | 0.014 | 0.014 | 0.014 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.013 | 0.012 | 0.012 | 0.012 |
| laying hens | 0.00027 | 0.00026 | 0.00026 | 0.00029 | 0.00035 | 0.00035 | 0.00034 | 0.00034 | 0.00034 | 0.00034 | 0.00034 | 0.00034 | 0.00034 | 0.00034 | 0.00034 |
| broilers | 0.00016 | 0.00014 | 0.00015 | 0.00018 | 0.00020 | 0.00019 | 0.00016 | 0.00015 | 0.00015 | 0.00015 | 0.00015 | 0.00015 | 0.00015 | 0.00015 | 0.00015 |
| turkeys | 0.00067 | 0.00067 | 0.00070 | 0.00084 | 0.00090 | 0.00091 | 0.00092 | 0.00090 | 0.00089 | 0.00089 | 0.00089 | 0.00089 | 0.00086 | 0.00085 | 0.00081 |
| pullets | 0.00011 | 0.00010 | 0.00009 | 0.00010 | 0.00012 | 0.00012 | 0.00011 | 0.00011 | 0.00011 | 0.00011 | 0.00011 | 0.00011 | 0.00011 | 0.00011 | 0.00011 |
| ducks | 0.00024 | 0.00024 | 0.00024 | 0.00025 | 0.00027 | 0.00027 | 0.00026 | 0.00027 | 0.00027 | 0.00027 | 0.00027 | 0.00027 | 0.00027 | 0.00026 | 0.00026 |
| geese | 0.00024 | 0.00024 | 0.00025 | 0.00027 | 0.00030 | 0.00030 | 0.00028 | 0.00029 | 0.00028 | 0.00029 | 0.00029 | 0.00029 | 0.00028 | 0.00028 | 0.00028 |

Trend discussion for Key Sources

Dairy cattle, other cattle and swine are key sources of NH₃ emissions from manure management. The time series of the total NH₃ emissions from all three categories are predominantly driven by the development of the animal numbers, see Table 1. However, the effect of decreasing animal numbers is partly compensated by the continuously increasing animal performance. This leads to increasing N excretions per animal, see Table 2, which, in principle, is reflected by increasing implied emission factors, see Table 3. For swine, as of 2012, the IEF is almost constant over time due to the use of air scrubbing systems that, to a high degree, remove NH₃ from the housings.

For NO_x there are no key categories.

Recalculations

All time series of the emission inventory have completely been recalculated since 1990. Tables REC-1 and REC-2 compare the recalculated time series for NH₃ and NO_x from 3B with the respective data of last year's submission. The total emissions of NH₃ slightly higher than those of submission 2021. The emissions from cattle are higher than in last year's submission mainly due to the incorporation of new housing data from the 2020 agricultural census. The extent of cattle housing in tied systems, which has a lower NH₃ emission factor than loose housing, has decreased in the last years. The NH₃ emissions from swine and poultry are lower than in the 2021 submission due to the use of new data on raw protein content in fattening pig and broiler feed. See main page of the agricultural sector ([Chapter 5 - NFR 3 - Agriculture \(OVERVIEW\)](#)), list of **recalculation reasons, No. 1, 7, and 8**.

The overall NH₃ emissions of other animals increased for the years 2017-2019 due to newly available figures on animal numbers of sheep and horses, see recalculations reasons No. 1 ([Chapter 5 - NFR 3 - Agriculture \(OVERVIEW\)](#)), list of **recalculation reasons, No. 1**. Further details on recalculations are described in Vos et al. (2022), Chapter 3.5.2.

Tables REC-1 and REC-2: Comparison of the NH₃ and NO_x emissions of the submissions (SUB) 2021 and 2022

| NH₃ emissions from manure management, in Gg | | | | | | | | | | | | | | | | |
|---|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | SUB | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Total | 2022 | 307.85 | 257.56 | 256.39 | 256.42 | 251.11 | 252.27 | 257.23 | 259.54 | 262.08 | 261.24 | 259.61 | 259.48 | 254.22 | 251.43 | 249.17 |
| | 2021 | 303.26 | 256.08 | 256.91 | 257.32 | 252.88 | 253.17 | 257.21 | 258.42 | 260.03 | 256.36 | 253.73 | 252.85 | 247.20 | 243.31 | |

| | | | | | | | | | | | | | | | | |
|----------------------|-------------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Dairy cattle | 2022 | 62.19 | 54.13 | 50.82 | 51.39 | 52.87 | 53.48 | 53.79 | 55.15 | 56.27 | 57.48 | 57.79 | 58.21 | 58.22 | 58.88 | 59.09 |
| | 2021 | 62.19 | 54.13 | 50.81 | 51.28 | 52.71 | 52.44 | 51.86 | 52.30 | 52.48 | 52.72 | 52.10 | 51.64 | 50.74 | 50.40 | |
| Other cattle | 2022 | 91.43 | 78.85 | 76.86 | 71.97 | 73.10 | 71.17 | 71.64 | 73.40 | 74.47 | 74.55 | 74.32 | 73.70 | 72.46 | 71.69 | 70.41 |
| | 2021 | 91.39 | 78.88 | 76.79 | 71.73 | 72.48 | 69.75 | 69.39 | 70.26 | 70.48 | 69.85 | 68.90 | 67.62 | 65.85 | 64.47 | |
| Swine | 2022 | 121.81 | 91.84 | 95.23 | 97.70 | 91.92 | 92.81 | 95.19 | 93.59 | 94.09 | 92.36 | 91.07 | 91.20 | 87.44 | 85.01 | 84.75 |
| | 2021 | 117.12 | 89.92 | 95.08 | 98.23 | 93.21 | 94.56 | 97.47 | 96.29 | 97.28 | 94.06 | 93.26 | 93.92 | 90.54 | 88.50 | |
| poultry | 2022 | 22.84 | 21.58 | 24.10 | 25.93 | 24.79 | 26.62 | 28.46 | 29.33 | 29.25 | 28.95 | 28.64 | 28.52 | 28.22 | 27.95 | 27.00 |
| | 2021 | 22.96 | 21.99 | 24.87 | 26.65 | 26.06 | 28.23 | 30.34 | 31.50 | 31.80 | 31.84 | 31.68 | 31.95 | 32.45 | 32.43 | |
| Other animals | 2022 | 9.59 | 11.16 | 9.37 | 9.43 | 8.43 | 8.18 | 8.16 | 8.07 | 8.00 | 7.89 | 7.79 | 7.85 | 7.88 | 7.90 | 7.93 |
| | 2021 | 9.59 | 11.16 | 9.37 | 9.43 | 8.43 | 8.18 | 8.16 | 8.07 | 8.00 | 7.89 | 7.79 | 7.72 | 7.62 | 7.51 | |

NO_x emissions from manure management, in Gg

| | SUB | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|----------------------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Total | 2022 | 1.731 | 1.554 | 1.516 | 1.505 | 1.487 | 1.460 | 1.439 | 1.437 | 1.444 | 1.436 | 1.427 | 1.416 | 1.393 | 1.379 | 1.365 |
| | 2021 | 1.720 | 1.551 | 1.523 | 1.534 | 1.529 | 1.502 | 1.480 | 1.475 | 1.482 | 1.465 | 1.455 | 1.444 | 1.418 | 1.401 | |
| Dairy cattle | 2022 | 0.671 | 0.597 | 0.570 | 0.553 | 0.524 | 0.517 | 0.501 | 0.499 | 0.503 | 0.507 | 0.506 | 0.503 | 0.503 | 0.507 | 0.506 |
| | 2021 | 0.671 | 0.597 | 0.570 | 0.552 | 0.521 | 0.513 | 0.495 | 0.491 | 0.493 | 0.495 | 0.494 | 0.489 | 0.488 | 0.489 | |
| Other cattle | 2022 | 0.690 | 0.604 | 0.587 | 0.550 | 0.548 | 0.529 | 0.525 | 0.534 | 0.539 | 0.538 | 0.535 | 0.530 | 0.521 | 0.515 | 0.506 |
| | 2021 | 0.696 | 0.614 | 0.600 | 0.566 | 0.562 | 0.538 | 0.529 | 0.532 | 0.532 | 0.526 | 0.519 | 0.509 | 0.497 | 0.487 | |
| Swine | 2022 | 0.281 | 0.256 | 0.270 | 0.309 | 0.320 | 0.318 | 0.317 | 0.307 | 0.305 | 0.295 | 0.289 | 0.287 | 0.273 | 0.262 | 0.258 |
| | 2021 | 0.264 | 0.242 | 0.263 | 0.322 | 0.351 | 0.355 | 0.359 | 0.353 | 0.357 | 0.344 | 0.343 | 0.346 | 0.335 | 0.328 | |
| poultry | 2022 | 0.026 | 0.024 | 0.027 | 0.032 | 0.039 | 0.042 | 0.043 | 0.045 | 0.045 | 0.045 | 0.045 | 0.045 | 0.045 | 0.044 | 0.044 |
| | 2021 | 0.026 | 0.024 | 0.028 | 0.033 | 0.040 | 0.043 | 0.045 | 0.047 | 0.048 | 0.048 | 0.048 | 0.048 | 0.049 | 0.049 | |
| Other animals | 2022 | 0.063 | 0.073 | 0.062 | 0.062 | 0.055 | 0.053 | 0.053 | 0.053 | 0.052 | 0.051 | 0.051 | 0.051 | 0.051 | 0.052 | 0.052 |
| | 2021 | 0.063 | 0.073 | 0.062 | 0.062 | 0.055 | 0.053 | 0.053 | 0.053 | 0.052 | 0.051 | 0.051 | 0.050 | 0.050 | 0.049 | |

Planned improvements

No improvements are planned at present.

NMVOG

In 2020, NMVOC emissions from manure management amount to 289.8 which is 96.9 % of total NMVOC emissions from the agricultural sector. 84.7 % originate from cattle, 4.8 % from pigs, and 9.4 % from poultry.

Method

The Tier 2 methodology provided by EMEP (2019)-3B-28 was used to assess the NMVOC emissions from manure management for dairy cattle and other cattle. For all other animals the Tier 1 methodology (EMEP (2019)-3B-17) was used.

Activity data

Animal numbers serve as activity data, see Table 1.

Emission factors

For the Tier 2 methodology applied to dairy cattle and other cattle the following data was used:

- gross feed intake in MJ per year, country specific data from the annual reporting of greenhouse gas emissions, see NIR 2022, Chapter 5.1.3.3,
- proportion x_{house} of the year the animals spend in the livestock building: country specific data, being equal to $1 - x_{\text{graz}}$ with x_{graz} the proportion of the year spent on pasture, see NIR 2022, Chapter 19.3.2,

- $FRAC_{silage}$: 1 as proposed by EMEP (2019)-3B-29, since silage feeding for cattle is considered dominant in Germany
- $FRAC_{silage\ store}$: 0.25 as proposed by EMEP (2019)-3B-30 for European conditions
- $EF_{NMVOC, silage\ feeding}$, $EF_{NMVOC, house}$, $EF_{NMVOC, graz}$ are taken from EMEP (2019)-3B-32, table 3.11 as 0.0002002, 0.0000353 and 0.0000069 kg NMVOC/MJ feed intake, respectively,
- $EF_{NH_3, storage}$, $EF_{NH_3, building}$ and $EF_{NH_3, application}$ are taken from the NH_3 reporting (see above and 3.D).

For all other animal categories the Tier 1 emission factors for NMVOC were used as provided in EMEP (2019)-3B-18, Table 3.4 [10]: For horses the emission factors for feeding with silage was chosen, for all other animals the emission factors for feeding without silage. Due to missing country-specific emission factors or emission factors that do not correspond to the inventory's animal categories, the emission factors provided in EMEP (2019)-3B-18, Table 3.4, were used to define specific emission factors for weaners, boars, lambs, ponies/light horses and pullets, see Vos et al. (2022), Chapter 3.3.4.2. The implied emission factors given in Table 4 relate the overall NMVOC emissions to the number of animals in each animal category. The IEFs for dairy cattle and other cattle are much higher than the EMEP Tier 1 EF, which are 17.937 kg NMVOC for dairy cattle and 8.902 kg NMVOC for other cattle. The only possible explanation for those huge differences is that the EMEP Tier 2 and Tier 1 methods are not consistent.

The IEFs for the other categories provided in Table 4 correspond to the EMEP Tier 1 emission factors, except for horses, sheep, swine and other poultry. These categories comprise subcategories with different emission factors so that their overall IEFs in Table 4 represent subpopulation-weighted national mean values. Note that other poultry in Germany includes not only geese and ducks but also pullets. For pullets no default EF is given in the EMEP guidebook (EMEP, 2019), hence the EF of broilers has been adopted (because of similar housing). This assumption significantly lowers the overall IEF of other poultry in Table 4 (the IEFs are listed separately for each poultry category). The IEF of the sheep category is significantly lower than the EMEP Tier 1 emission factor, because for lambs the EF is assumed to be 40% lower compared to an adult sheep in accordance with the difference in N excretion between lambs and adult sheep.

Table 4: IEF for NMVOC from manure management

| | IEF in kg NMVOC per animal place | | | | | | | | | | | | | | |
|---------------------|----------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| dairy cattle | 30.940 | 32.695 | 35.472 | 36.709 | 37.242 | 37.609 | 37.625 | 37.467 | 37.891 | 38.155 | 38.552 | 38.524 | 39.322 | 40.134 | 40.799 |
| other cattle | 11.738 | 11.729 | 11.852 | 11.720 | 11.769 | 11.694 | 11.638 | 11.617 | 11.540 | 11.533 | 11.483 | 11.475 | 11.481 | 11.533 | 11.574 |
| horses | 6.497 | 6.491 | 6.688 | 6.660 | 6.644 | 6.643 | 6.642 | 6.641 | 6.644 | 6.646 | 6.648 | 6.651 | 6.654 | 6.657 | 6.660 |
| sheep | 0.131 | 0.131 | 0.132 | 0.132 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 | 0.131 |
| goats | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 | 0.542 |
| swine | 0.695 | 0.698 | 0.690 | 0.682 | 0.669 | 0.663 | 0.656 | 0.654 | 0.652 | 0.651 | 0.649 | 0.648 | 0.648 | 0.648 | 0.642 |
| laying hens | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 | 0.165 |
| broilers | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 |
| turkeys | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 |
| pullets | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 | 0.108 |
| ducks | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 |
| geese | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 | 0.489 |

Trend discussion for Key Sources

Dairy cattle and other cattle are key sources of NMVOC emissions from manure management. The total NMVOC emissions from both animal categories strongly correlate with the animal numbers given in Table 1 (dairy cattle: $R^2 = 0.887$; other cattle: $R^2 = 0.998$).

Recalculations

All time series of the emission inventory have completely been recalculated since 1990. Table REC-3 compares the recalculated time series of the NMVOC emissions from 3.B with the respective data of last year's submission. The recalculated total emissions are slightly lower in some years and slightly higher in other years. This is due to improved methodology for the suckler cows (**recalculation reason 4**, see [main page of the agricultural sector](#)), and the use of the data from the official agricultural census of 2020 (recalculation reason 1) which are changing the NH_3 emissions that have impact to the Tier 2 methodology which is applied for cattle NMVOC emissions. For other animals there are differences back to the year 2017. These differences are caused by updated animal numbers (see above). Further details on recalculations

are described in Vos et al. (2022), Chapter 3.5.2.

Table REC-3: Comparison of NMVOC emissions of the submissions (SUB) 2021 and 2022

| NMVOC emissions from manure management, in Gg | | | | | | | | | | | | | | | | |
|---|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | SUB | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Total | 2022 | 391.24 | 332.95 | 318.87 | 298.26 | 297.23 | 297.42 | 299.37 | 304.56 | 306.95 | 305.72 | 302.74 | 299.98 | 296.20 | 293.49 | 289.79 |
| | 2021 | 390.80 | 332.18 | 318.01 | 297.65 | 296.79 | 297.13 | 299.17 | 304.38 | 306.74 | 305.38 | 303.07 | 300.94 | 297.86 | 295.83 | |
| Dairy cattle | 2022 | 196.61 | 170.97 | 162.10 | 155.51 | 155.79 | 157.59 | 157.67 | 159.89 | 162.77 | 163.48 | 162.60 | 161.76 | 161.25 | 161.00 | 159.99 |
| | 2021 | 196.61 | 170.97 | 162.10 | 155.50 | 155.75 | 157.45 | 157.39 | 159.40 | 162.04 | 162.45 | 161.82 | 161.15 | 160.81 | 160.84 | |
| Other cattle | 2022 | 154.16 | 125.04 | 118.15 | 103.14 | 101.55 | 97.54 | 96.82 | 97.80 | 97.48 | 96.31 | 94.72 | 92.75 | 90.10 | 87.97 | 85.42 |
| | 2021 | 153.72 | 124.27 | 117.29 | 102.54 | 101.14 | 97.38 | 96.89 | 98.11 | 98.00 | 97.00 | 95.83 | 94.23 | 91.95 | 90.15 | |
| Other animals | 2022 | 40.46 | 36.94 | 38.62 | 39.61 | 39.89 | 42.29 | 44.89 | 46.87 | 46.70 | 45.93 | 45.42 | 45.47 | 44.85 | 44.52 | 44.38 |
| | 2021 | 40.46 | 36.94 | 38.62 | 39.61 | 39.89 | 42.29 | 44.89 | 46.87 | 46.70 | 45.93 | 45.42 | 45.56 | 45.11 | 44.84 | |

Planned improvements

No improvements are planned at present.

TSP, PM10 and PM2.5

In 2020, TSP emissions from manure management amount to 71.3 % of total emissions from the agricultural sector. Within the emissions from manure management 22.5 % originate from cattle, 39.5 % from pigs, and 37.4 % from poultry. 42.8 % of the PM₁₀ emissions from the agricultural sector are caused by manure management, where 34.4 % originate from cattle, 19.1 % from pigs, and 45.6 % from poultry. PM_{2.5} emissions from the agricultural sector mostly originate from manure management (84.9 %), of which are 78.0 % from cattle, 3.0 % from pigs, and 17.5 % from poultry.

Method

EMEP (2013-3B-26) provided a Tier 2 methodology. In the current Guidebook (EMEP, 2019), this methodology has been replaced by a Tier 1 methodology. However, EF for cattle derived with the EMEP 2013 Tier 2 methodology remained unchanged. Therefore, the EMEP 2013¹⁰⁾ methodology was kept for cattle. For swine the EMEP 2013 methodology was formally kept but the EMEP 2019 Tier 1 EF was used both for slurry and solid based manure management systems. The same was done with the EMEP 2016 EFs for laying hens (used for cages and perchery). In case the EMEP 2019 EFs are simply rounded EMEP 2013 EFs, the unrounded EMEP 2013 EFs were kept. The inventory considers air scrubber systems in swine and poultry husbandry. For animal places equipped with air scrubbing the emission factors are reduced according to the removal efficiency of the air scrubber systems (90 % for TSP and PM₁₀, 70 % for PM_{2.5}). For details see Vos et al. (2022), Chapter 3.3.4.3.3.

Activity data

Animal numbers serve as activity data, see Table 1.

Emission factors

Tier 1 emission factors for TSP, PM₁₀ and PM_{2.5} from livestock husbandry are provided in EMEP (2019-3B-19), Table 3.5 and 55, Table A1.7. For cattle the Tier 2 emission factors provided in EMEP (2013-3B-29), Table 3-11 were used, because they differentiate between slurry and solid manure systems and were also used to develop the EMEP 2019 Tier 1 emissions factors.

The implied emission factors given in Table 5 relate the overall TSP and PM emissions to the number of animals in each animal category. The Guidebook does not indicate whether EFs have considered the condensable component (with or without).

Table 5: IEF for TSP, PM₁₀ & PM_{2.5} from manure management

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| IEF in kg TSP per animal place | | | | | | | | | | | | | | | |
| dairy cattle | 1.2124 | 1.4016 | 1.4544 | 1.4738 | 1.4882 | 1.4953 | 1.5028 | 1.5101 | 1.5189 | 1.5270 | 1.5365 | 1.5451 | 1.5540 | 1.5631 | 1.5721 |
| other cattle | 0.5194 | 0.5107 | 0.5017 | 0.4916 | 0.4804 | 0.4790 | 0.4788 | 0.4788 | 0.4775 | 0.4766 | 0.4763 | 0.4760 | 0.4758 | 0.4751 | 0.4746 |
| horses | 0.3514 | 0.3512 | 0.3558 | 0.3552 | 0.3548 | 0.3548 | 0.3548 | 0.3548 | 0.3548 | 0.3549 | 0.3549 | 0.3550 | 0.3551 | 0.3551 | 0.3552 |
| sheep | 0.0484 | 0.0478 | 0.0489 | 0.0486 | 0.0489 | 0.0485 | 0.0485 | 0.0485 | 0.0483 | 0.0482 | 0.0482 | 0.0482 | 0.0480 | 0.0482 | 0.0482 |
| goats | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 | 0.0914 |
| swine | 0.8260 | 0.8366 | 0.8320 | 0.8247 | 0.8104 | 0.8043 | 0.8086 | 0.8037 | 0.8002 | 0.7910 | 0.7918 | 0.7891 | 0.7865 | 0.7813 | 0.7858 |
| laying hens | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1900 | 0.1899 | 0.1898 | 0.1897 | 0.1894 | 0.1890 | 0.1889 |
| broilers | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0399 | 0.0399 | 0.0397 | 0.0396 | 0.0395 | 0.0394 | 0.0394 |
| turkeys | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 |
| pullets | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 |
| ducks | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 |
| geese | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 |
| IEF in kg PM₁₀ per animal place | | | | | | | | | | | | | | | |
| dairy cattle | 0.5557 | 0.6426 | 0.6668 | 0.6757 | 0.6823 | 0.6855 | 0.6890 | 0.6923 | 0.6964 | 0.7001 | 0.7044 | 0.7084 | 0.7125 | 0.7166 | 0.7208 |
| other cattle | 0.2403 | 0.2363 | 0.2320 | 0.2273 | 0.2221 | 0.2214 | 0.2213 | 0.2213 | 0.2207 | 0.2203 | 0.2202 | 0.2200 | 0.2200 | 0.2196 | 0.2194 |
| horses | 0.1619 | 0.1619 | 0.1639 | 0.1636 | 0.1634 | 0.1634 | 0.1634 | 0.1634 | 0.1634 | 0.1634 | 0.1635 | 0.1635 | 0.1635 | 0.1635 | 0.1636 |
| sheep | 0.0194 | 0.0192 | 0.0196 | 0.0195 | 0.0196 | 0.0195 | 0.0194 | 0.0194 | 0.0194 | 0.0193 | 0.0193 | 0.0193 | 0.0192 | 0.0193 | 0.0193 |
| goats | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 | 0.0368 |
| swine | 0.1241 | 0.1255 | 0.1244 | 0.1230 | 0.1199 | 0.1187 | 0.1185 | 0.1177 | 0.1170 | 0.1158 | 0.1156 | 0.1151 | 0.1147 | 0.1139 | 0.1141 |
| laying hens | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0400 | 0.0399 | 0.0399 | 0.0398 | 0.0398 |
| broilers | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0199 | 0.0198 | 0.0198 | 0.0198 | 0.0197 | 0.0197 |
| turkeys | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 | 0.1100 |
| pullets | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 |
| ducks | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 | 0.1400 |
| geese | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 | 0.2400 |
| IEF in kg PM_{2.5} per animal place | | | | | | | | | | | | | | | |
| dairy cattle | 0.3616 | 0.4181 | 0.4338 | 0.4396 | 0.4439 | 0.4460 | 0.4483 | 0.4505 | 0.4531 | 0.4555 | 0.4583 | 0.4609 | 0.4636 | 0.4663 | 0.4690 |
| other cattle | 0.1574 | 0.1548 | 0.1520 | 0.1490 | 0.1457 | 0.1453 | 0.1452 | 0.1452 | 0.1448 | 0.1445 | 0.1445 | 0.1444 | 0.1443 | 0.1441 | 0.1439 |
| horses | 0.1027 | 0.1026 | 0.1039 | 0.1038 | 0.1036 | 0.1036 | 0.1036 | 0.1036 | 0.1036 | 0.1037 | 0.1037 | 0.1037 | 0.1037 | 0.1037 | 0.1038 |
| sheep | 0.0059 | 0.0059 | 0.0060 | 0.0060 | 0.0060 | 0.0060 | 0.0060 | 0.0060 | 0.0059 | 0.0059 | 0.0059 | 0.0059 | 0.0059 | 0.0059 | 0.0059 |
| goats | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 | 0.0112 |
| swine | 0.0056 | 0.0057 | 0.0056 | 0.0056 | 0.0054 | 0.0054 | 0.0053 | 0.0053 | 0.0053 | 0.0052 | 0.0052 | 0.0052 | 0.0052 | 0.0052 | 0.0051 |
| laying hens | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 | 0.0030 |
| broilers | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 |
| turkeys | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 | 0.0200 |
| pullets | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 | 0.0020 |
| ducks | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 | 0.0180 |
| geese | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 | 0.0320 |

Trend discussion for Key Sources

Swine and laying hens are key sources of TSP emissions from manure management. The total TSP emissions from swine mainly follow the animal numbers given in Table 1. However, due to air scrubbing and different emission factors of the different housing systems of the four swine subcategories (sows with piglets, weaners, fattening pigs, boars) and the varying population shares in those housing systems the R² of the linear regression is lower than 1 (0.78). For laying hens and

broilers, due to the low prevalence of air scrubbing systems. TSP emissions almost perfectly correlate with the animal numbers provided in Table 1 ($R^2 = 1$).

Recalculations

Table REC-4 shows the effects of recalculations on emissions of particulate matter. Changes in the years 2000 through 2019 are a consequence of the use of the data of the official agricultural census 2020 as well as new animal population figures for the years 2017-2019 (**recalculation reason 1**, see [main page of the agricultural sector](#)). Further details on recalculations are described in Vos et al. (2022), Chapter 3.5.2.

Table REC-4: Comparison of particle emissions (TSP, PM_{10} & $PM_{2.5}$) of the submissions (SUB) 2021 and 2022

| TSP, PM10, PM2.5 emissions from manure management, in Gg | | | | | | | | | | | | | | | | |
|--|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | SUB | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| TSP | 2022 | 50.04 | 42.24 | 42.43 | 41.25 | 40.30 | 41.80 | 43.95 | 45.14 | 45.45 | 44.74 | 44.54 | 44.59 | 43.65 | 43.04 | 42.99 |
| TSP | 2021 | 50.04 | 42.24 | 42.44 | 41.26 | 40.32 | 41.79 | 43.90 | 45.06 | 45.33 | 44.58 | 44.35 | 44.40 | 43.55 | 42.90 | |
| PM₁₀ | 2022 | 14.34 | 12.71 | 12.63 | 12.29 | 12.31 | 12.76 | 13.33 | 13.84 | 13.82 | 13.63 | 13.48 | 13.43 | 13.20 | 13.02 | 12.91 |
| PM₁₀ | 2021 | 14.34 | 12.71 | 12.63 | 12.29 | 12.32 | 12.75 | 13.31 | 13.80 | 13.77 | 13.56 | 13.39 | 13.36 | 13.17 | 13.00 | |
| PM_{2.5} | 2022 | 5.01 | 4.47 | 4.18 | 3.89 | 3.85 | 3.87 | 3.93 | 4.03 | 4.04 | 4.02 | 3.97 | 3.94 | 3.86 | 3.79 | 3.72 |
| PM_{2.5} | 2021 | 5.01 | 4.47 | 4.18 | 3.89 | 3.86 | 3.86 | 3.91 | 4.01 | 4.01 | 3.97 | 3.91 | 3.88 | 3.80 | 3.72 | |

Planned improvements

No improvements are planned at present.

Uncertainty

Details will be described in [chapter 1.7](#).

¹⁾

Vos, C., Rösemann C., Haenel H-D., Dämmgen U., Döring U., Wulf S., Eurich-Menden B., Freibauer A., Döhler H., Schreiner C., Osterburg B. & Fuß, R. (2022): Calculations of gaseous and particulate emissions from German Agriculture 1990 –2020. Report on methods and data (RMD), Submission 2022. Thünen Report (in preparation).

<https://www.thuenen.de/de/ak/arbeitsbereiche/emissionsinventare>

²⁾

NIR, National Inventory Report 2022 for the German Greenhouse Gas Inventory 1990-2020. Available in April 2022

³⁾

Rösemann C, Haenel H-D, Dämmgen U, Freibauer A, Döring, U, Wulf S, Eurich-Menden B, Döhler H, Schreiner C, and Osterburg B, 2017, Calculations of gaseous and particulate emissions from German Agriculture 1990 – 2015. Report on methods and data (RMD), Submission 2017. Thünen Report 46, 423 p.

⁴⁾

Weingarten, 1995: Das „Regionalisierte Agrar- und Umweltinformationssystem für die Bundesrepublik Deutschland“ (RAUMIS). Berichte über die Landwirtschaft Band 73, 272-302.

⁵⁾

Henrichsmeyer, W.; Cypris, Ch.; Löhe, W.; Meuth, M.; Isermeyer F; Heinrich, I.; Schefski, A.; Neander, E.; Fasterding, F.; Neumann, M.; Nieberg, H., 1996: Entwicklung des gesamtdeutschen Agrarsektormodells RAUMIS96. Endbericht zum Kooperationsprojekt. Forschungsbericht für das BMELF (94 HS 021), Bonn, Braunschweig.

⁶⁾

EMEP/EEA air pollutant emission inventory guidebook – 2019, EEA Report No 13/2019,

<https://www.eea.europa.eu/publications/emep-eea-guidebook-2019>.

⁷⁾

Dämmgen U., Hutchings N.J. (2008): Emissions of gaseous nitrogen species from manure management - a new approach. Environmental Pollution 154, 488-497.

⁸⁾

IPCC – Intergovernmental Panel on Climate Change (2006): IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4 Agriculture, Forestry and Other Land Use.

⁹⁾

Reidy B. et al. (2008): Reidy B., Dämmgen U., Döhler H., Eurich-Menden B., Hutchings N.J., Luesink H.H., Menzi H., Misselbrook T.H., Monteny G.-J., Webb J. (2008): Comparison of models used for the calculation of national NH₃ emission

inventories from agriculture: liquid manure systems. Atmospheric Environment 42, 3452-3467.

¹⁰⁾

EMEP (2013): EMEP/EEA air pollutant emission inventory guidebook – 2013