

## 1.A.5.b ii - Military Aviation

### Short description

In sub-category *1.A.5.b ii - Other, Mobile (including Military)* emissions from military aviation are reported.

Method	AD	EF	Key Category Analysis
T1	NS	CS, D	see <a href="#">superordinate chapter</a>

### Methodology

#### Activity data

The Energy Balance of the Federal Republic of Germany (AGEB) provides the basis for the activity data used. Since the Energy Balance does not provide separate listings of military agencies' final energy consumption as of 1995 – and includes this consumption in line 67, under “commerce, trade, services and other consumers” – additional sources of energy statistics had to be found for source category 1.A.5.

For source category 1.A.5.b, consumption data for **kerosene**, until 1995, were drawn from a special analysis of the Working Group on Energy Balances (AGEB).

For the years as of 1995, the official mineral-oil data of the Federal Republic of Germany (Amtliche Mineralölstatistik der Bundesrepublik Deutschland 2012), prepared by the Federal Office of Economics and Export Control (BAFA), are used (BAFA, 2021) <sup>1)</sup>. Provided in units of 1,000 tonnes [kt], these amounts have to be converted into terajoules [TJ] on the basis of the relevant net calorific values given by (AGEB, 2021) <sup>2)</sup>.

As there is no consistent AGEB data available for **aviation gasoline**, delivery data from BAFA is used.


Table 1: Sources for consumption data in 1.A.5.b

Relevant years	Data Source
through 1994	<b>AGEB</b> - Special evaluation 1990-1994
since 1995	<b>BAFA</b> - Official oil data, table 7j, column: 'An das Militär'

Table 2: Annual fuel consumption in military aviation, in terajoules

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>Jet Kerosene</b>	38,385	16,143	9,862	2,200	2,441	2,554	3,597	4,396	3,286	4,114	1,171	2,049	3,060	3,726	3,845	1,507	1,025	3,746	2,904
<b>Aviation Gasoline</b>	15.2	6.35	1.09	0.26	2.48	14.8	0.00	0.04	0.17	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.17	0.04	0.00
<b>Σ 1.A.5.b ii</b>	<b>38,400</b>	<b>16,149</b>	<b>9,863</b>	<b>2,200</b>	<b>2,443</b>	<b>2,569</b>	<b>3,597</b>	<b>4,396</b>	<b>3,286</b>	<b>4,114</b>	<b>1,171</b>	<b>2,049</b>	<b>3,060</b>	<b>3,726</b>	<b>3,845</b>	<b>1,507</b>	<b>1,025</b>	<b>3,746</b>	<b>2,904</b>

<sup>1)</sup> possible reason for jumps in delivered amounts: storage (resulting in no (2008, 2011+) or very small deliveries (2009) (see also: FAQs)



## Emission factors

Without better information, constant tier1 values are used mainly (see table below).

NOTE: As the aircraft used for military purposes differ strongly from those used in civil aviation, the country specific EF used for kerosene in 1.A.3.a could not be used for reporting emissions from 1.A.5.b as well. Therefore, and due to missing

information on the technical developments within the military aircraft fleet, the EF values applied show no trend.

Table 3: Country-specific emission factors, in kg/TJ

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>JET KEROSENE</b>																			
<b>NH<sub>3</sub></b>	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00
<b>NMVO</b>	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0	98.0
<b>NO<sub>x</sub></b>	205	205	205	205	205	205	205	205	205	205	205	205	205	205	205	205	205	205	205
<b>SO<sub>x</sub></b>	25.1	15.2	8.46	6.34	5.92	5.50	5.07	4.65	4.65	4.65	4.65	4.65	4.65	4.65	4.65	4.65	4.65	4.65	4.65
<b>BC<sup>1</sup></b>	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76	5.76
<b>PM<sup>2</sup></b>	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0
<b>CO</b>	485	485	485	485	485	485	485	485	485	485	485	485	485	485	485	485	485	485	485
<b>AVIATION GASOLINE</b>																			
<b>NH<sub>3</sub></b>	NE																		
<b>NMVO</b>	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300
<b>NO<sub>x</sub></b>	302	302	302	302	302	302	302	302	302	302	302	302	302	302	302	302	302	302	302
<b>SO<sub>x</sub></b>	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51
<b>BC<sup>1</sup></b>	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10
<b>PM<sup>2</sup></b>	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50	7.50
<b>TSP<sup>3</sup></b>	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7	22.7
<b>CO</b>	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000	15,000

<sup>1</sup> estimated via a f-BCs (avgas: 0.15, jet kerosene: 0.48) as provided in <sup>3)</sup>

<sup>2</sup> EF(PM<sub>2.5</sub>) also applied for PM<sub>10</sub>, and TSP (assumption: > 99% of TSP from diesel oil combustion consists of PM<sub>2.5</sub>)

<sup>3</sup> TSP from leaded aviation gasoline = EF(Pb) x 1.6 (see also: FAQs)



For the country-specific emission factors applied for particulate matter, no clear indication is available, whether or not condensables are included.




For information on the **emission factors for heavy-metal and POP exhaust emissions**, please refer to Appendix 2.3 - Heavy Metal (HM) exhaust emissions from mobile sources and Appendix 2.4 - Persistent Organic Pollutant (POP) exhaust emissions from mobile sources. - Here, regarding lead and TSP from leaded avgas, constant tier1 EFs based on the average lead content of AvGas 100 LL are used.

## Discussion of emission trends




As only NFR 1.A.5.b as a whole is taken into account within the key category analysis, this country-specific sub-sector is not considered separately.

Due to the application of very several tier1 emission factors, most emission trends reported for this sub-category only reflect the trend in fuel deliveries. Therefore, the fuel-consumption dependend trends in emission estimates are only influenced by the annual fuel mix.



Here, as the EF(BC) are estimated via fractions provided in <sup>4)</sup>, black carbon emissions follow the corresponding emissions of PM<sub>2.5</sub>.

Nonetheless, this NFR category shows interesting trends for emissions of **Lead (Pb)** from leaded gasoline (until 1997) and aviation gasoline:



Until 1997, lead emissions were dominated by the combustion of leaded gasoline in military ground-based vehicles. Therefore, the over-all trend for lead emissions from military vehicles and aircraft is driven mostly by the abolition of leaded gasoline in 1997. Towards this date, the amount of leaded gasoline decreased significantly. After 1997, the only source for lead from mobile fuel combustion is avgas used in military aircraft. As for avgas, the trend of consumption is more or less decreasing steadily until 2005 but then shows a strong increase for 2006 and '07 (!), followed by no (2008 and 2011) or very small deliveries (2009, 2010). As mentioned above, there are no real consumption data available: AD is based on fuel deliveries to the military only. Thus, especially the trends for the use of aviation gasoline and the resulting emissions show this significant jumps in 2006 and 07, falling back to zero in 2008 and 2011ff. The party is aware of this issue and will try to solve it as soon as data allows. (see also: FAQ)

## Recalculations

With both **activity data** and **emission factors** remaining unrevised, no recalculations took place with this submission.



For pollutant-specific information on recalculated emission estimates for Base Year and 2019, please see the pollutant specific recalculation tables following chapter [8.1 - Recalculations](#).

## Uncertainties

Uncertainty estimates for **activity data** of mobile sources derive from research project FKZ 360 16 023: "Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland". For detailed information, please refer to the project's final report [<https://www.umweltbundesamt.de/publikationen/ermittlung-unsicherheiten-den-modellen-tremod> here] (German version only!).

Uncertainty estimates for **emission factors** were compiled during the PAREST research project. Here, the final report has not yet been published.

## Planned improvements

There are no specific improvements planned at the moment.

## FAQs

### ***What is the reason for the big jumps in the consumption of aviation gasoline in 2006 & '07 and the zero-consumption in 2008?***

As mentioned above, consumption is deducted from AGEB and BAFA data on the amounts of fuels delivered to the sector. Therefore, the big jumps reported for 2006 & '07 might result from the storage of aviation gasoline in military stocks. Consequentially, the 0.00 TJ reported for 2008 show the missing of any deliveries to the military and should not be misunderstood as a non-use. The party is aware of this issue and will try to solve it as soon as data allows.

### ***On which basis does the party estimate the reported lead emissions from aviation gasoline?***

assumption by party: aviation gasoline = AvGas 100 LL (AvGas 100 LL is the predominant sort of aviation gasoline in Western Europe) <sup>1)</sup> lead content of AvGas 100 LL: 0.56 g lead/liter (as tetra ethyl lead) <sup>2)</sup>

The applied procedure is similar to the one used for calculating lead emissions from leaded gasoline used in road transport. (There, in contrast to aviation gasoline, the lead content constantly declined resulting in a ban of leaded gasoline in 1997.)

### ***What is the country-specific methodology for estimating the reported TSP emissions from aviation gasoline?***

The TSP emissions calculated depend directly on the reported lead emissions: The emission factor for TSP is 1.6 times the emission factor used for lead:  $EF(TSP) = 1.6 \times EF(Pb)$ . - The applied procedure is similar to the one used for calculating TSP emissions from leaded gasoline used in road transport.

### ***Why does the party report TSP emissions from leaded avgas, but no such $PM_{2.5}$ or $PM_{10}$ emissions?***

The  $EF(TSP)$  is estimated from the  $EF(Pb)$  which has been calculated from the lead content of Avgas 100 LL. There is no information on the per centual shares of  $PM_{2.5}$  &  $PM_{10}$  in the reported TSP and therefore no  $EF(PM_{2.5})$  &  $EF(PM_{10})$  were deducted.

### ***Why are similar EF applied for estimating exhaust heavy metal emissions from both fossil and biofuels?***

The EF provided in <sup>5)</sup> represent summatory values for (i) the fuel's and (ii) the lubricant's heavy-metal content as well as (iii) engine wear. Here, there might be no heavy metal contained the biofuels. But since the specific shares of (i), (ii) and (iii) cannot be separated, and since the contributions of lubricant and engine wear might be dominant, the same emission factors are applied to biodiesel and bioethanol.

<sup>1)</sup> BAFA, 2021: Federal Office of Economics and Export Control (Bundesamt für Wirtschaft und Ausfuhrkontrolle, BAFA): Amtliche Mineralöl-daten für die Bundesrepublik Deutschland; URL: [https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel\\_amtliche\\_daten\\_2019\\_dezember.html](https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel_amtliche_daten_2019_dezember.html), Eschborn, 2021.

<sup>2)</sup> AGEB, 2021: Working Group on Energy Balances (Arbeitsgemeinschaft Energiebilanzen (Hrsg.), AGEB): Energiebilanz für die Bundesrepublik Deutschland; URL: <http://www.ag-energiebilanzen.de/7-0-Bilanzen-1990-2019.html>, (Aufruf: 29.11.2021), Köln & Berlin, 2021.

<sup>3), 4), 5)</sup> EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook – 2019, Copenhagen, 2019.

<sup>1)</sup> <https://en.wikipedia.org/wiki/Avgas> : "...Common in North America and western Europe, limited availability elsewhere worldwide."

<sup>2)</sup> EMEP/EEA GB 2016: "Thus, general emission factors for the stationary combustion of kerosene and the combustion of gasoline in cars may be applied. The only exception is lead. Lead is added to aviation gasoline to increase the octane number. The lead content is higher than in leaded car gasoline, and the maximum permitted levels in the UK are shown below. A value of 0.6 g of lead per litre of gasoline should be used as the default value if there is an absence of more accurate information. Actual data may be obtained from oil companies."