

# 1.A.3.d ii - National Navigation

## Short description

Under category 1.A.3.d ii - *National Navigation* emissions from national navigation (both inland and maritime) are reported.

| Category Code | Method          |                 |                 |        |     | AD    |     |     |     |      | EF       |     |     |                  |                   |
|---------------|-----------------|-----------------|-----------------|--------|-----|-------|-----|-----|-----|------|----------|-----|-----|------------------|-------------------|
| 1.A.3.d ii    | T1, T2, T3      |                 |                 |        |     | NS, M |     |     |     |      | CS, D, M |     |     |                  |                   |
| Key Category  | SO <sub>2</sub> | NO <sub>x</sub> | NH <sub>3</sub> | NM VOC | CO  | BC    | Pb  | Hg  | Cd  | Diox | PAH      | HCB | TSP | PM <sub>10</sub> | PM <sub>2.5</sub> |
| 1.A.3.d ii    | -/-             | L/-             | -/-             | -/-    | -/- | -/-   | -/- | -/- | -/- | -/-  | -/-      | -/- | -/- | L/T              | L/T               |

**T** = key source by Trend **L** = key source by Level

| Methods   |                                 |
|-----------|---------------------------------|
| <b>D</b>  | Default                         |
| <b>RA</b> | Reference Approach              |
| <b>T1</b> | Tier 1 / Simple Methodology *   |
| <b>T2</b> | Tier 2*                         |
| <b>T3</b> | Tier 3 / Detailed Methodology * |
| <b>C</b>  | CORINAIR                        |
| <b>CS</b> | Country Specific                |
| <b>M</b>  | Model                           |

\* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

| AD - Data Source for Activity Data |                                      |
|------------------------------------|--------------------------------------|
| <b>NS</b>                          | National Statistics                  |
| <b>RS</b>                          | Regional Statistics                  |
| <b>IS</b>                          | International Statistics             |
| <b>PS</b>                          | Plant Specific data                  |
| <b>AS</b>                          | Associations, business organisations |
| <b>Q</b>                           | specific questionnaires, surveys     |
| EF - Emission Factors              |                                      |
| <b>D</b>                           | Default (EMEP Guidebook)             |
| <b>C</b>                           | Confidential                         |
| <b>CS</b>                          | Country Specific                     |
| <b>PS</b>                          | Plant Specific data                  |

## Methodology

### Activity data

As described for the over-all sector 1.A.3.d and all other navigational activities [here](#) ], specific fuel consumption data for NFR 1.A.3.d ii is included in the primary fuel deliveries data provided in NEB lines 6 ('International Maritime Bunkers') and 64 ('Coastal and Inland Navigation') <sup>1)</sup>.

Here, the annual fuel consumption for domestic *maritime* navigation are modelled within <sup>2)</sup> based on AIS data and deduced from NEB lines 6 and 64 respectively, depending on whether or not a certain ship is registered by the International Maritime Organization (IMO). Here, fuels consumed by large, IMO-registered and sea-going ships and vessels are included in NEB line 6 whereas fuels consumed by smaller ships without IMO-registration are included in NEB line 64. After these deductions, the amounts of fuels remaining in NEB 64 are allocated to domestic *inland* navigation.

Table 1: Annual over-all fuel consumption for domestic navigation, in terajoule

|                   | 1990   | 1995   | 2000   | 2005   | 2010   | 2011   | 2012   | 2013   | 2014   | 2015   | 2016   | 2017   | 2018   | 2019   |
|-------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| <b>Diesel Oil</b> | 36,604 | 29,855 | 18,648 | 18,596 | 16,183 | 16,954 | 16,601 | 16,824 | 18,532 | 22,781 | 24,167 | 22,400 | 23,847 | 21,556 |

|                       |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Heavy fuel oil</b> | 11,723        | 8,041         | 8,577         | 7,172         | 6,114         | 5,961         | 6,410         | 6,376         | 6,046         | 50,0          | 7,05          | 7,01          | 190           | 358           |
| <b>Σ</b>              |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>1.A.3.d ii</b>     | <b>48,326</b> | <b>37,896</b> | <b>27,224</b> | <b>25,768</b> | <b>22,297</b> | <b>22,916</b> | <b>23,011</b> | <b>23,200</b> | <b>24,578</b> | <b>22,831</b> | <b>24,174</b> | <b>22,407</b> | <b>24,037</b> | <b>21,914</b> |

gallery size="medium" : 1A3dii\_AD.png gallery

Table 2: Specific fuel consumption data for domestic maritime and inland navigation, in terajoule

|                                     | 1990          | 1995          | 2000          | 2005          | 2010          | 2011          | 2012          | 2013          | 2014          | 2015          | 2016          | 2017          | 2018          | 2019          |
|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>NATIONAL MARITIME NAVIGATION</b> |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Diesel Oil</b>                   | 15940         | 11258         | 11860         | 9962          | 8685          | 8489          | 9046          | 9047          | 9965          | 13359         | 16295         | 15221         | 16336         | 13961         |
| <b>Heavy fuel oil</b>               | 11723         | 8041          | 8577          | 7172          | 6114          | 5961          | 6410          | 6376          | 6046          | 50,0          | 7,05          | 7,01          | 190           | 358           |
| <b>NATIONAL INLAND NAVIGATION</b>   |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>Diesel Oil</b>                   | 20.664        | 18.597        | 6.788         | 8.634         | 7.497         | 8.466         | 7.556         | 7.777         | 8.567         | 9.422         | 7.873         | 7.179         | 7.511         | 7.595         |
| <b>Σ</b>                            |               |               |               |               |               |               |               |               |               |               |               |               |               |               |
| <b>1.A.3.d ii</b>                   | <b>48.326</b> | <b>37.896</b> | <b>27.224</b> | <b>25.768</b> | <b>22.297</b> | <b>22.916</b> | <b>23.011</b> | <b>23.200</b> | <b>24.578</b> | <b>22.831</b> | <b>24.174</b> | <b>22.407</b> | <b>24.037</b> | <b>21.914</b> |

The emission factors applied for **national maritime navigation** are derived from different sources and therefore are of very different quality.

For the main pollutants, country-specific implied values are used, that are based on tier3 EF included in the BSH model <sup>3)</sup> which mainly relate on values from the EMEP/EEA guidebook 2019 <sup>4)</sup>. These modelled IEFs take into account the ship specific information derived from AIS data as well as the mix of fuel-qualities applied depending on the type of ship and the current state of activity.

Here, for **sulphur dioxide** and **particulate matter**, annual values are available representing the impact of fuel sulphur legislation. In addition, regarding SO<sub>x</sub>, the increasing operation of so-called scrubbers in order to fulfil emission limits especially within SECA areas is reflected for heavy fuel oil.

Table 3: Country-specific emission factors applied for fuels used in domestic maritime navigation, in [kg/TJ]

|                         | 1990  | 1995  | 2000  | 2005  | 2010  | 2011  | 2012  | 2013  | 2014  | 2015  | 2016  | 2017  | 2018  | 2019  |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>DIESEL OIL</b>       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| <b>NH<sub>3</sub></b>   | 0,32  | 0,32  | 0,32  | 0,32  | 0,32  | 0,32  | 0,32  | 0,32  | 0,32  | 0,33  | 0,32  | 0,32  | 0,32  | 0,32  |
| <b>NM VOC</b>           | 48,7  | 48,7  | 48,7  | 48,7  | 48,7  | 48,7  | 48,7  | 48,4  | 48,0  | 44,8  | 44,7  | 45,0  | 45,2  | 45,2  |
| <b>NO<sub>x</sub></b>   | 1.070 | 1.070 | 1.070 | 1.070 | 1.069 | 1.069 | 1.069 | 1.073 | 1.077 | 1.151 | 1.132 | 1.157 | 1.128 | 1.128 |
| <b>SO<sub>x</sub></b>   | 465,5 | 419,0 | 232,8 | 186,2 | 69,8  | 65,18 | 54,53 | 52,56 | 50,48 | 40,74 | 40,61 | 40,74 | 40,76 | 40,76 |
| <b>BC<sup>1</sup></b>   | 109,2 | 98,6  | 54,7  | 43,8  | 16,4  | 15,4  | 15,3  | 15,2  | 15,2  | 16,3  | 16,9  | 16,9  | 16,5  | 16,5  |
| <b>PM<sub>2,5</sub></b> | 352,4 | 318,0 | 176,5 | 141,2 | 53,0  | 49,6  | 49,5  | 49,1  | 49,1  | 52,7  | 54,5  | 54,5  | 53,1  | 53,1  |
| <b>PM<sub>10</sub></b>  | 377,1 | 340,3 | 188,9 | 151,0 | 56,7  | 53,0  | 53,0  | 52,5  | 52,6  | 56,4  | 58,3  | 58,3  | 56,9  | 56,9  |
| <b>TSP<sup>2</sup></b>  | 377,1 | 340,3 | 188,9 | 151,0 | 56,7  | 53,0  | 53,0  | 52,5  | 52,6  | 56,4  | 58,3  | 58,3  | 56,9  | 56,9  |
| <b>CO</b>               | 127   | 128   | 128   | 128   | 128   | 128   | 128   | 127   | 128   | 134   | 139   | 138   | 136   | 136   |
| <b>HEAVY FUEL OIL</b>   |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| <b>NH<sub>3</sub></b>   | 0,34  | 0,34  | 0,34  | 0,34  | 0,34  | 0,34  | 0,34  | 0,34  | 0,35  | 0,33  | 0,33  | 0,34  | 0,35  | 0,35  |
| <b>NM VOC</b>           | 36,6  | 36,6  | 36,6  | 36,6  | 36,6  | 36,6  | 36,6  | 37,6  | 37,8  | 30,0  | 36,8  | 30,4  | 28,3  | 28,3  |
| <b>NO<sub>x</sub></b>   | 1.379 | 1.378 | 1.378 | 1.378 | 1.378 | 1.377 | 1.379 | 1.382 | 1.393 | 1.348 | 1.245 | 1.360 | 1.503 | 1.503 |
| <b>SO<sub>x</sub></b>   | 1.319 | 1.332 | 1.323 | 1.336 | 496   | 496   | 496   | 496   | 506   | 47,5  | 49,3  | 46,4  | 49,8  | 49,8  |
| <b>BC<sup>1</sup></b>   | 57,4  | 58,0  | 57,6  | 58,2  | 21,6  | 21,6  | 21,6  | 22,1  | 22,4  | 18,1  | 24,7  | 18,3  | 14,7  | 14,7  |
| <b>PM<sub>2,5</sub></b> | 479   | 483   | 480   | 485   | 180   | 180   | 180   | 184   | 187   | 151   | 205   | 153   | 123   | 123   |
| <b>PM<sub>10</sub></b>  | 526   | 532   | 528   | 533   | 198   | 198   | 198   | 203   | 206   | 166   | 226   | 168   | 135   | 135   |
| <b>TSP<sup>2</sup></b>  | 526   | 532   | 528   | 533   | 198   | 198   | 198   | 203   | 206   | 166   | 226   | 168   | 135   | 135   |
| <b>CO</b>               | 162   | 162   | 162   | 162   | 162   | 162   | 162   | 162   | 167   | 165   | 198   | 167   | 134   | 134   |

<sup>1</sup> estimated from f-BCs as provided in <sup>5)</sup>: f-BC (HFO) = 0.12, f-BC (MDO/MGO) = 0.31 as provided in <sup>6)</sup>, chapter: 1.A.3.d.i,

1.A.3.d.ii, 1.A.4.c.iii Navigation, Tables 3-1 & 3-2

<sup>2</sup> ratios PM<sub>2.5</sub> : PM<sub>10</sub> : TSP derived from the tier1 default EF as provided in <sup>7)</sup>, chapter: 1.A.3.d.i, 1.A.3.d.ii, 1.A.4.c.iii Navigation, Tables 3-1 & 3-2



For the country-specific emission factors applied for particulate matter, no clear indication is available, whether or not condensables are included.

For main pollutants and particulate matter from **national inland navigation**, modelled emission factors are available from TREMOD (Knörr et al. (2019a)) <sup>8)</sup>. Here, for SO<sub>2</sub>, and PM, annual values reflect the impact of fuel-sulphur legislation.

Table 4: Country-specific emission factors for diesel fuels used in domestic inland navigation, in [kg/TJ]

|                       | 1990  | 1995  | 2000  | 2005  | 2010  | 2011  | 2012  | 2013  | 2014  | 2015  | 2016  | 2017  | 2018  | 2019  |
|-----------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>NH<sub>3</sub></b> | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  | 0,23  |
| <b>NMVOC</b>          | 96,4  | 87,9  | 77,7  | 73,4  | 69,3  | 68,2  | 66,9  | 65,9  | 64,6  | 63,4  | 62,2  | 61,0  | 59,8  | 59,8  |
| <b>NO<sub>x</sub></b> | 1.327 | 1.331 | 1.336 | 1.294 | 1.245 | 1.236 | 1.224 | 1.213 | 1.198 | 1.184 | 1.170 | 1.155 | 1.141 | 1.141 |
| <b>SO<sub>x</sub></b> | 86,4  | 60,5  | 60,5  | 60,5  | 60,5  | 0,37  | 0,37  | 0,37  | 0,38  | 0,37  | 0,38  | 0,38  | 0,38  | 0,38  |
| <b>BC<sup>1</sup></b> | 17,5  | 16,0  | 14,1  | 12,2  | 10,07 | 9,86  | 9,61  | 9,44  | 9,23  | 9,04  | 8,87  | 8,69  | 8,53  | 8,53  |
| <b>PM<sup>2</sup></b> | 56,5  | 51,7  | 45,6  | 39,3  | 32,5  | 31,8  | 31,0  | 30,5  | 29,8  | 29,2  | 28,6  | 28,0  | 27,5  | 27,5  |
| <b>CO</b>             | 417   | 387   | 337   | 307   | 277   | 272   | 266   | 261   | 256   | 251   | 246   | 241   | 236   | 236   |

<sup>1</sup> calculated from f-BC as provided in <sup>9)</sup>, Chapter: 1.A.3.d.i, 1.A.3.d.ii, 1.A.4.c.iii, Table 3-2: f-BC (MDO/MGO) = 0.31

<sup>2</sup> EF(PM<sub>2.5</sub>) also applied for PM<sub>10</sub>, and TSP (assumption: > 99% of TSP from diesel oil combustion consists of PM<sub>2.5</sub>)



With respect to the emission factors applied for particulate matter, given the circumstances during test-bench measurements, condensables are most likely included at least partly. <sup>1)</sup>



For information on the **emission factors for heavy-metal and POP exhaust emissions**, please refer to [Appendix 2.3 - Heavy Metal \(HM\) exhaust emissions from mobile sources](#) and [Appendix 2.4 - Persistent Organic Pollutant \(POP\) exhaust emissions from mobile sources](#)].

## Discussion of emission trends

Table: Outcome of Key Category Analysis

|      |                        |                         |
|------|------------------------|-------------------------|
| for: | <b>PM<sub>10</sub></b> | <b>PM<sub>2.5</sub></b> |
| by:  | L/T                    | L/T                     |

For **ammonia**, **NMVOC**, and **nitrogen oxides** as well as **carbon monoxide**, emission trends more or less represent the trend in over-all fuel consumption.

Nonetheless, for these pollutants, annual emission factors from BSH <sup>10)</sup> and TREMOD <sup>11)</sup> have been applied for national *maritime* and *inland* navigation, respectively, reflecting the technical development of the German inland navigation fleet.

Here, the trends in **sulphur dioxide** and **particulate matter** emissions reflect the impact of ongoing fuel-sulphur legislation especially in maritime navigation.

## Recalculations

Table 5: Revised fuel consumption data for national maritime navigation, in terajoules

Furthermore, the country-specific **emission factors** applied for diesel fuels used in **domestic inland navigation** have been revised within TREMOD <sup>12)</sup>:

Table 6: Revised country-specific emission factors for diesel fuels used in domestic inland navigation, in [kg/T]

[illegible]

|                             |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-----------------------------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
| <b>relative change</b>      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| <b>CARBON MONOXIDE - CO</b> |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| <b>Submission 2021</b>      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| <b>Submission 2020</b>      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| <b>absolute change</b>      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| <b>relative change</b>      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

In contrast, the country-specific **emission factors** applied for fuels used in **national maritime navigation** remain unaltered.



For more information on recalculated emission estimates for Base Year and 2018, please see the pollutant-specific recalculation tables following chapter [8.1 - Recalculations](#).

## Uncertainties

Uncertainty estimates for **activity data** of mobile sources derive from research project FKZ 360 16 023: "Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland" by Knörr et al. (2009) <sup>13)</sup>.

## Planned improvements

Besides the **routine revisions of the models** used for maritime and inland navigation, no specific improvements are scheduled.

<sup>14)</sup> : 2 : BAFA (2019): Federal Office of Economics and Export Control (Bundesamt für Wirtschaft und Ausfuhrkontrolle, BAFA): Amtliche Mineralöl-daten für die Bundesrepublik Deutschland; URL: [https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel\\_amtliche\\_daten\\_2018\\_dezember.xlsx?\\_\\_blob=publicationFile&v=4](https://www.bafa.de/SharedDocs/Downloads/DE/Energie/Mineraloel/moel_amtliche_daten_2018_dezember.xlsx?__blob=publicationFile&v=4), (Aufruf: 29.11.2019), Eschborn, 2019. : 3 : MWV, 2019: German Petroleum Industry Association (Mineralölwirtschaftsverband, MWV): MWV Jahresberichte: URL: <https://www.mwv.de/publikationen/jahresberichte/>, Berlin, 2019. [DEICHNIK2020> Deichnik (2020): Deichnik, K.: Aktualisierung und Revision des Modells zur Berechnung der spezifischen Verbräuche und Emissionen des von Deutschland ausgehenden Seeverkehrs. from Bundesamts für Seeschifffahrt und Hydrographie (BSH); Hamburg, 2020.)] : 5 : Knörr et al. (2019a): Knörr, W., Heidt, C., Gores, S., & Bergk, F.: ifeu Institute for Energy and Environmental Research (Institut für Energie- und Umweltforschung Heidelberg gGmbH, ifeu): Fortschreibung des Daten- und Rechenmodells: Energieverbrauch und Schadstoffemissionen des motorisierten Verkehrs in Deutschland 1960-2035, sowie TREMOD, im Auftrag des Umweltbundesamtes, Heidelberg & Berlin, 2019. : 6 : EMEP/EEA (2019): EMEP/EEA air pollutant emission inventory guidebook 2019, Copenhagen, 2019. <sup>15)</sup>

<sup>1)</sup> (bibcite 1)

<sup>2)</sup> (bibcite 4)

<sup>3)</sup> (bibcite 4)

<sup>4)</sup> (bibcite 2)

<sup>5)</sup> (bibcite 2)

<sup>6)</sup> (bibcite 2)

<sup>7)</sup> (bibcite 2)

<sup>8)</sup> (bibcite 5)

<sup>9)</sup> (bibcite 3)

<sup>12)</sup> (bibcite 5)

<sup>13)</sup> (bibcite 7)

<sup>14)</sup> AGEb, 2020: Working Group on Energy Balances (Arbeitsgemeinschaft Energiebilanzen (Hrsg.), AGEb): Energiebilanz für die Bundesrepublik Deutschland; URL: <http://www.ag-energiebilanzen.de/7-0-Bilanzen-1990-2018.html>, Köln & Berlin, 2020.

<sup>15)</sup> Knörr et al. (2009): Knörr, W., Heldstab, J., & Kasser, F.: Ermittlung der Unsicherheiten der mit den Modellen TREMOD und TREMOD-MM berechneten Luftschadstoffemissionen des landgebundenen Verkehrs in Deutschland; final report; URL: <https://www.umweltbundesamt.de/sites/default/files/medien/461/publikationen/3937.pdf>, FKZ 360 16 023, Heidelberg &

Zürich, 2009.

<sup>1)</sup>

During test-bench measurements, temperatures are likely to be significantly higher than under real-world conditions, thus reducing condensation. On the contrary, smaller dilution (higher number of primary particles acting as condensation germs) together with higher pressures increase the likeliness of condensation. So over-all condensables are very likely to occur but different to real-world conditions.