2.C.7.b - Nickel Production 1/2

2.C.7.b - Nickel Production

In subcategory *NFR 2.C.7.b - Nickel production* the TSP, SO₂ and Ni emissions from nickel mining are reported. Reporting only covers the year 1990 because in 1991 nickel mining stopped in Germany.

Category Code		Method			AD				EF						
2.C.7.b		T1			AS				D						
	NO _x	NMVOC	SO ₂	NH ₃	PM _{2.5}	PM ₁₀	TSP	вс	СО	Pb	Cd	Hg	Diox	PAH	нсв
Key Category:	-	-	-/-	-	-	-	-/-	-	-	-	-	-	-	-	-

T = key source by Trend L = key source by Level

Methods	
D	Default
RA	Reference Approach
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
Т3	Tier 3 / Detailed Methodology *
С	CORINAIR
CS	Country Specific
M	Model
·	· · · · · · · · · · · · · · · · · · ·

^{*} as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

AD - Data Source for Activity Data						
NS	National Statistics					
RS	Regional Statistics					
IS	International Statistics					
PS	Plant Specific data					
AS	Associations, business organisations					
Q	specific questionnaires, surveys					

EF	- Emission Factors
D	Default (EMEP Guidebook)
С	Confidential
CS	Country Specific
PS	Plant Specific data

Method

Activity data

The yearly production figure was taken from the annual statistical report of the US Geological Survey (USGS) for non-ferrous metals [Lit. 1].

Emission factors

The emission factors are the default values according to the 2019 EMEP/EEA air pollutant emission inventory guidebook [Lit. 2].

Table 1: applied Tier1 emission factors

Pollutant	EF	Unit
SO ₂	18	kg/t
TSP	0.3	kg/t
Ni	0.025	kg/t

2.C.7.b - Nickel Production 2/2

Uncertainties

Recalculations

No recalculations were carried out with this submission.

Planned improvements

No category specific improvements necessary.

Bibliography

Lit. 1: US Geological Survey https://www.usgs.gov

Lit. 2: EMEP/EEA air pollutant emission inventory guidebook 2019 https://www.eea.europa.eu