## 2.G.4(c) - Charcoal

In sub-category NFR 2.G.4( c) - Charcoal TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions from charcoal used for barbecue are reported.

Method	AD	EF	Key Category
T1	NS	D	For <b><u>2.G.</u> L:</b> TSP, Cd, PM <sub>10</sub> / <b>L &amp; T:</b> PM <sub>2.5</sub> , Pb

 $\mathbf{T}$  = key source by Trend  $\mathbf{L}$  = key source by Level

Methods	
D	Default
RA	Reference Approach
T1	Tier 1 / Simple Methodology *
Т2	Tier 2*
Т3	Tier 3 / Detailed Methodology *
С	CORINAIR
CS	Country Specific
М	Model
* as described in the EMEP/COR	NAIR Emission Inventory Guidebook - 2007, in the group specific chapters
AD - Data Source for Activity	Data
NS National Statistics	
<b>RS</b> Regional Statistics	
IS International Statistics	
PS Plant Specific data	
AS Associations, business orgar	isations
<b>Q</b> specific questionnaires, surv	eys
EF - Emission Factors	
<b>D</b> Default (EMEP Guidebook)	
C Confidential	
CS Country Specific	
PS Plant Specific data	