

2.H.2 - Food & Beverages Industry

| Category Code | Method | | | | AD | | | | EF | | | | | | |
|---------------|-----------------|-----------------|-----------------|-------|----|----|----|----|----|------|-----|-----|-----|------------------|-------------------|
| 2.H.2 | T1 | | | | NS | | | | CS | | | | | | |
| Key Category | SO ₂ | NO _x | NH ₃ | NMVOC | CO | BC | Pb | Hg | Cd | Diox | PAH | HCB | TSP | PM ₁₀ | PM _{2.5} |
| 2.H.2 | - | - | - | -/- | - | - | - | - | - | - | - | - | -/- | -/- | -/- |

T = key source by Trend **L** = key source by Level

| Methods | |
|----------------|---------------------------------|
| D | Default |
| RA | Reference Approach |
| T1 | Tier 1 / Simple Methodology * |
| T2 | Tier 2* |
| T3 | Tier 3 / Detailed Methodology * |
| C | CORINAIR |
| CS | Country Specific |
| M | Model |

* as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2007, in the group specific chapters.

AD - Data Source for Activity Data

| | |
|-----------|--------------------------------------|
| NS | National Statistics |
| RS | Regional Statistics |
| IS | International Statistics |
| PS | Plant Specific data |
| AS | Associations, business organisations |
| Q | specific questionnaires, surveys |

EF - Emission Factors

| | |
|-----------|--------------------------|
| D | Default (EMEP Guidebook) |
| C | Confidential |
| CS | Country Specific |
| PS | Plant Specific data |

xyz

Methodology

Activity data

xx

Emission factors

Table 1: Overview of applied emission factors

| Pollutant | Name of Category | EF | Unit | Trend |
|-----------------|------------------|-------|------|----------|
| NO _x | clinker burning | 0.5 | kg/t | falling |
| SO ₂ | clinker burning | 0.25 | kg/t | falling |
| NMVOC | clinker burning | 0.046 | kg/t | constant |
| NH ₃ | clinker burning | 0.044 | kg/t | falling |
| Hg | clinker burning | 0.022 | g/t | falling |
| Pb | clinker burning | 0.016 | g/t | falling |
| Cd | clinker burning | 0.004 | g/t | falling |
| PCB | clinker burning | 28.0 | µg/t | constant |

| | | | | |
|-------------------------|-------------------------|-------|------|----------------|
| PCDD | clinker burning | 0.040 | µg/t | constant |
| B(a)P | clinker burning | 1.0 | mg/t | constant |
| PAH | clinker burning | 240 | mg/t | constant |
| TSP | <i>clinker grinding</i> | 0.046 | kg/t | <i>falling</i> |
| PM₁₀ | <i>clinker grinding</i> | 0.041 | kg/t | <i>falling</i> |
| PM_{2.5} | <i>clinker grinding</i> | 0.037 | kg/t | <i>falling</i> |

Trends in emissions

All trends in emissions correspond to trends of emission factors in table above. No rising trends are to identify.

Invalid Link

Emission trends in NFR 2.H.2

Recalculations

With **activity data** and **emission factors** remaining unrevised, no recalculations have been carried out compared to last year's submission.



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following [chapter 8.1 - Recalculations](#).

Planned improvements

For purposes of updating the EF project has started in 2020, but results are planned not before 2021 ¹⁾.

¹⁾ ReFoPlan FKZ - xx: „yy“