# 5.D.1 - Domestic & Commercial Wastewater Handling

## **Short description**

Category Code		Method				AD					EF					]
5.D.1			Т1				Ν	IS					D			
	NO <sub>x</sub>	NMVOC	SO <sub>2</sub>	NH3	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC	CO	PbC	dŀ	lg [	Diox	PAH	HCB	
Key Category:	-	-/-	-	-	-	-	-	-	-	-   -		-	-	-	-	
<b>T</b> = key source b	by Tre	end $\mathbf{L} = \mathbf{k}$	key s	ourc	e by L	evel										
Methods																
	<b>D</b>			Defau	Default											
RA			Refer	Reference Approach												
					Tier 1 / Simple Methodology *											
	Т2				Tier 2											
	Т3				Tier 3	/ Det	ailed	Me	thod	ology	/*					
					CORINAIR											
	CS				Count		ecific	2								
	Μ				Mode											
* as described in	n the	EMEP/CC	DRINA	AIR E	missic	n Inv	entor	y Gi	uidel	book	- 2	007	, in t	he g	roup	specific chapters
AD - Data Soui			ity D	ata												
NS National Stat																
RS Regional Sta																
IS International																
PS Plant Specifi																
AS Associations					IS											
<b>Q</b> specific ques			irvey	'S												
EF - Emission I																
Default (EME	P Gu	idebook)														
C Confidential																
CS Country Spe																
Plant Specific	c dat	а														

In category **5.D.1**, <u>NMVOC emissions</u> from domestic and commercial wastewater handling are reported. The domestic section is covered by wastewaters of municipal origin (large centralised plants; ranging from 1000 up to >100.000 resident values). The commercial section is covered by industrial and commercial wastewaters, co-treated in municipal wwt-plants.

## Method

Emissions reported under this category are calculated using the Tier 1 approach of the EMEP/EEA Guidebook 2019, where the emission factor (EF) is 15 mg/m<sup>3</sup> wastewater (Part B, 5.D, chap. 3.2.2, Table 3-1, p. 7<sup>11</sup>). This EF is multiplied with the total amount of wastewater (AD) treated in domestic and commercial wwt-plants, following the equation:



#### Activity data

Total volumes of treated municipal wastewater are derived by the German statistical agency (Statistisches Bundesamt, Fachserie 19, Reihe 2.1.2<sup>2)</sup>). The data source is published on a three-year basis with new data only for the respective year of the update. The availability of the data starts in 1991 with an exception for the following update, which was for 1995. Missing data are inter- or extrapolated

#### **Emisson factors**

See method

## Uncertainties

The AD from Statistisches Bundesamt have an uncertainty of  $\pm 3\%$  (normal distribution) whereas the uncertainty for the EF, due to its range (5/50 mg/m<sup>3</sup>), is -70 / +210 % and the distribution lognormal.

### Recalculations

Recalculations were not necessary.



For **pollutant-specific information on recalculated emission estimates for Base Year and 2019**, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

## **Planned improvements**

Currently no improvements are planned.

<sup>1)</sup> EMEP/EEA, 2019: EMEP/EEA air pollutant emission inventory guidebook 2019, Copenhagen, 2019 <sup>2)</sup> Statistisches Bundesamt, Fachserie 19, Reihe 2.1.2