

# Explanation of Key Trends - Ammonia

## Obligations

Germany has made a commitment under the Gothenburg Protocol to reduce ammonia emissions. Since 2010, it is no longer permissible to exceed a National Emission Ceiling of 550 kt NH<sub>3</sub> for Germany as whole. The revised Gothenburg Protocol and the revised NEC Directive both define emission reduction targets relative to a 2005 base year, mandating 5% (2020) and 29% (2030) reductions respectively.

While Germany's compliance with these obligations is not discussed here, further information on this subject can be found in [Chapter 9 - Projections](#) and [Chapter 11 - Adjustments and Emission Ceiling Exceedance](#).

## Main drivers

The Main Drivers for NH<sub>3</sub> emissions are agricultural emissions from **Manure Management (NFR 4.B)** with 43% of total 1990 emissions and a 26% reduction between 1990-2021 and **Agricultural Soils (NFR 4.D)** with even 53% of total 1990 emissions and a 32% decrease between 1990 and 2021.


The overall emission trend follows the agricultural emissions closely with a **total reduction of 29% between 1990 and 2021**. The decrease of NH<sub>3</sub> emission in the year 1991 is due to a reduced livestock population that followed after the German reunification, while no explicit trend is discernible for the years up to 2016. Between 2016 and 2021 the emissions are dropping every year adding up to a 15% drop. Between 2019 and 2020, emissions dropped by 6.6%, only topped by the reduction between 1990 and 1991.

Table: NH<sub>3</sub> Emissions 1990-2021

Total Emissions (kt)																Trend: latest compared to	
1990	1995	2000	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	1990	last years
726	620	633	612	625	628	633	639	647	644	637	619	591	570	530	516	↓ -28.9%	↓



NH<sub>3</sub> trend by sector



NH<sub>3</sub> trend by sector, from 2005