

# 1.A.1.c - Manufacture Of Solid Fuels And Other Energy Industries

## Short description

Source category 1.A.1.c - *Manufacture Of Solid Fuels And Other Energy Industries* includes hard-coal and lignite mining, coking and briquetting plants and extraction of crude oil and natural gas. Used-oil processing plants are also included. Here, CO emissions from coking plants are reported in NFR sub-category 1.B.1.b.

Category Code	Method						AD						EF			
1.A.1.c	T2						NS						CS			
	NO <sub>x</sub>	NM VOC	SO <sub>2</sub>	NH <sub>3</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC	CO	PB	Cd	Hg	Diox	PAH	HCB	
Key Category:	L/T	-/-	L/T	-/-	-/T	-/-	L/T	-/-	-/-	-/-	L/T	L/T	-/-	-/-	-/-	

**T** = key source by Trend **L** = key source by Level

Methods	
<b>D</b>	Default
<b>T1</b>	Tier 1 / Simple Methodology *
<b>T2</b>	Tier 2*
<b>T3</b>	Tier 3 / Detailed Methodology *
<b>C</b>	CORINAIR
<b>CS</b>	Country Specific
<b>M</b>	Model

\* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in the group specific chapters.

AD - Data Source for Activity Data	
<b>NS</b>	National Statistics
<b>RS</b>	Regional Statistics
<b>IS</b>	International Statistics
<b>PS</b>	Plant Specific data
<b>As</b>	Associations, business organisations
<b>Q</b>	specific Questionnaires (or surveys)
<b>M</b>	Model / Modelled
<b>C</b>	Confidential
EF - Emission Factors	
<b>D</b>	Default (EMEP Guidebook)
<b>C</b>	Confidential
<b>CS</b>	Country Specific
<b>PS</b>	Plant Specific data
<b>M</b>	Model / Modelled



## Method

### Activity data

Fuel inputs for electricity production in power plants of the hard-coal and lignite mining sector are listed in Energy Balance line 12 “Industrial thermal power stations”. Fuel inputs for heat production in the transformation sector are listed in Energy Balance lines 33-39 <sup>1)</sup>.

Fuel inputs for electricity production in power stations of the hard-coal mining sector are determined with the help of figures of the Federal Statistical Office (DESTATIS). The activity rates for heat production in power stations of the hard-coal mining sector correspond to energy Balance line 34 “Energy input in pit and briquette plants of the hard-coal mining sector”. The listed fuel input for electricity and heat production in pit power plants is based on association information from DEBRIV (the federal German association of all lignite producing companies and their affiliated organisations). Inputs for heat production, especially for lignite drying for production of lignite products, are not shown in the Energy Balance. Those are calculated from figures for production of lignite products (STATISTIK DER KOHLENWIRTSCHAFT)<sup>2)</sup> and from the specific fuel inputs required for drying from DEBRIV.

Energy consumption data for hard-coal coke production are provided by the Energy Balance line 33.

The fuel input for heat production in the remaining transformation sector is obtained by combining the energy consumption figures in Energy Balance lines 33 to 39 (total energy consumption in the transformation sector). These figures include the pits' own consumption, facilities for petroleum and natural gas production and for processing of old oil; plants that produce coal products; plants for production and processing of fissile and fertile materials; and wastewater-treatment facilities.

### Emission factors

The emission factors for power stations and other boiler combustion for production of steam and hot water, in source category 1.A.1.c, have been taken from the research project “Determination and evaluation of emission factors for combustion systems in Germany for the years 1995, 2000 and 2010” (Rentz et al., 2002)<sup>3)</sup>. A detailed description of the procedure is presented in Chapter: [1.A.1.a - Public Electricity And Heat Production](#). In 2018 all emission factors for large combustion plants were revised (UBA, 2019)<sup>4)</sup>.

Table 1: Implied emission factors for manufacture of solid fuels and other energy industry

	SO <sub>x</sub>	NO <sub>x</sub>	TSP	CO	Pb	Hg	Cd
	[kg/TJ]				[g/TJ]		
<b>Hard Coal</b>	83.2	72.0	2.4	7.7	1.53	2.10	0.50
<b>Lignite</b>	104.1	78.6	3.1	25.1	1.49	3.19	0.16
<b>Pit gas</b>	2.0	118.0	0.3	72.0	NE	NE	NE
<b>Coke oven gas</b>	89.0	78.0	0.3	2.6	NE	NE	NE

	SO <sub>x</sub>	NO <sub>x</sub>	TSP	CO	Pb	Hg	Cd
<b>Sewage sludge</b>	2.0	29.0	0.5	2.2	3.20	2.40	0.34

The table gives an overview of the implied emission factors. In reality the German inventory compiling process is very complex and includes the use of a considerable number of emission factors, which cannot be published completely in the IIR. Actually there are different emission factors available for diverse fuel types, various techniques and licensing requirements. However, the implied emission factor may give an impression about the order of magnitude. PM<sub>10</sub> and PM<sub>2.5</sub> emission factors are calculated as a fraction of TSP. The share of PM<sub>10</sub> is 90 % and the share of PM<sub>2.5</sub> is 80 %. This is a simple but also conservative approach, knowing that, in reality, PM emissions depend on fuel, combustion and abatement technologies. PM emission reporting starts in 1995, since no sufficient information about the dust composition of the early 1990th is available. Emission factors of sewage sludge refer to mono-incineration, using fluidized-bed combustion. Emission factors of coke oven gas does not include underfiring systems of coking plants. The determination of emission factors of coking plants is described in the study: "Emissionsfaktoren zur Eisen und Stahlindustrie für die Emissionsberichterstattung" (VDEh, 2010)<sup>5)</sup>. Emission factors refer to the produced amount of coke, distinction is drawn between diffuse and channelled sources. The following graph gives an overview of the methodology:





Table 2: emission factors for coking plants (solely channelled sources)

SO <sub>x</sub>	NO <sub>x</sub>	CO	TSP	PM <sub>10</sub>	NH <sub>3</sub>	B[a]P	Benzene
[g/t product]						[mg/t product]	Unit
220.5	529.9	828.2	25.9	12.1	1.9	7.2	36.2

## Trend Discussion for Key Sources


The following diagrams give an overview and assistance for explaining dominant emission trends of selected pollutants.

### Sulfur Oxides - SO<sub>x</sub>



The graph shows sharp declining SO<sub>x</sub> emissions between 1990 and 1995 due to decreasing lignite consumption as well as the implementation of stricter regulations in eastern Germany. In the former GDR lignite industry was of prime importance for the economy. After the reunification lignite briquette production in eastern Germany collapsed. The remaining factories had to install flue gas desulphurisation plants. The strong decline of SO<sub>2</sub> emissions in 2012 can be explained by the change of some power plants from the industrial to the public sector as a result of the closure of hard coal mines. A further reduction of SO<sub>2</sub> emission factors followed in 2016 when the reduction efficiency of desulfurization plants increased from 95 to 96%. However, this effect is counterbalanced by the increased use of lignite.



## Nitrogen Oxides - NO<sub>x</sub>




NO<sub>x</sub> emissions decrease gradually from 1990 to 2001. The main reasons are the minor fuel use of lignite and of hard coal in


this sector and the adaptation of regulations in eastern Germany to the western standard. Besides German hard coal production decreased considerably since 1990. Therefore some hard coal fired industrial power plants changed from sector 1.A.1.c to the public sector. This is also the reason for the significant emission reduction in 2012.


### Total Suspended Matter - TSP & PM2.5





**Priority Heavy Metal - Hg & Cd**





Similar to SO<sub>x</sub> emissions, TSP and Priority Heavy Metal emission trends show a high dominance of emissions from lignite combustion. Like already discussed for other pollutant, the main reason for sharp declining emissions in this sector is the complete restructuring of the east German lignite industry. The low standard of dust abatement in eastern Germany in the early 1990s involved high heavy metal emissions too. The closing of briquette factories and the implementation of stricter regulations resulted in a considerably improvement of the air quality especially in the New German Länder.

## Recalculations

Recalculations were necessary for 2020 due to the implementation of the now finalised National Energy Balance.



For **pollutant-specific information on recalculated emission estimates for Base Year and 2020**, please see the recalculation tables following [chapter 8.1 - Recalculations](#).

## Planned improvements

At the moment, no category-specific improvements are planned.

---

<sup>1)</sup> AGEB, 2022: National energy balance and Satellite balance for renewable energy:

<https://ag-energiebilanzen.de/en/data-and-facts/energy-balance-2000-to-2030/>

<sup>2)</sup> Statistik der Kohlenwirtschaft, 2021 - URL: <https://www.kohlenstatistik.de>

<sup>3)</sup> Rentz, O. ; Karl, U. ; Peter, H.: Ermittlung und Evaluierung von Emissionsfaktoren für Feuerungsanlagen in Deutschland für die Jahre 1995, 2000 und 2010: Forschungsbericht 299 43 142; Forschungsvorhaben im Auftrag des Umweltbundesamt; Endbericht; Karlsruhe: Deutsch-Französisches Inst. f. Umweltforschung, Univ. (TH); 2002

<sup>4)</sup> Umweltbundesamt, 2019: Kristina Juhrich, Rolf Beckers: "Updating the Emission Factors for Large Combustion Plants":

<https://www.umweltbundesamt.de/publikationen/updating-emission-factors-large-combustion-plants>

<sup>5)</sup> Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung; Michael Hensmann, Sebastian Haardt, Dominik Ebert (VDEh-Betriebsforschungsinstitut GmbH, Düsseldorf, Juli 2010), FKZ: 3707 42 301/01 und 3707 41 111/2;

<https://www.umweltbundesamt.de/publikationen/emissionsfaktoren-zur-eisen-stahlindustrie-fuer>