1.A.2.b - Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals

Short description

Sub-category 1.A.2.b - Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals includes aluminium production (sub-divided into primary and resmelted aluminium) as well as lead production, thermal galvanisation, copper and zinc production.

In Germany, aluminium is produced at four foundries, in electrolytic furnaces with pre-burnt anodes. The principal emission sources are resulting from fuel provided in the energy related processes.

Category Code		Method				AD				EF					
1.A.2.b		T2				NS				?					
	NO _x	NMVOC	SO ₂	NΗ₃	PM _{2.5}	PM ₁₀	TSP	вс	СО	Pb	Cd	Hg	Diox	PAH	нсв
Key Category:	-/-	-/-	-/-	-/-	-	-	-/-	-	-/-	-	-	-	-	-	-

T = key source by Trend L = key source by Level

Methods	
D	Default
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
Т3	Tier 3 / Detailed Methodology *
С	CORINAIR
CS	Country Specific
М	Model

* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in the group specific chapters.

AD - Data Source for Activity Data					
NS	National Statistics				
RS	Regional Statistics				
IS	International Statistics				
PS	Plant Specific data				
As	Associations, business organisations				
Q	specific Questionnaires (or surveys)				
М	Model / Modelled				
С	Confidential				

EF	- Emission Factors
D	Default (EMEP Guidebook)
С	Confidential
CS	Country Specific
PS	Plant Specific data
М	Model / Modelled

Method

Activity data

The source of the fuel inputs consists of the statistics for the manufacturing sector (Statistik 060 - Energieverwendung des produzierenden Gewerbes / energy use in the manufacturing sector), DESTATIS, reporting number 27.43 and 27.44, production and initial processing of lead, zinc and tin, production and initial processing of copper - and, for differentiations relative to heat and electricity production, Statistik 067 (DESTATIS).

Data for fuel consumption for production and initial processing of precious metals are also provided by these statistics.

Emission factors

Reported pollutants are NOx, NMVOC, SO₂, NH₃ and CO. Instead, all particulate matter emissions are reported as process emissions under 2.C.

The underlying data for the emission factors used is provided by the report on the research project "Ermittlung und Evaluierung von Emissionsfaktoren für Feuerungsanlagen in Deutschland für die Jahre 1995, 2000 und 2010" (Determination and evaluation of emission factors for combustion systems in Germany for the years 1995, 2000 and 2010"; RENTZ et al, 2002)¹⁾. The values for the intermediate years 1996 - 1999 and 2001 - 2010 are obtained via linear interpolation; adjusted values for the following years.

Recalculations

Recalculations were necessary for 2019 due to the implementation of the now finalised National Energy Balance.



For pollutant-specific information on recalculated emission estimates for Base Year and 2019, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

Planned improvements

At the moment, no category specific improvements are planned.

¹⁾ RENTZ et al., 2002: Rentz, O.; Karl, U.; Peter, H.: Ermittlung und Evaluierung von Emissionsfaktoren für Feuerungsanlagen in Deutschland für die Jahre 1995, 2000 und 2010: Forschungsbericht 299 43 142; Forschungsvorhaben im Auftrag des Umweltbundesamt; Endbericht; Karlsruhe: Deutsch-Französisches Inst. f. Umweltforschung, Univ. (TH); 2002