2.A.3 - Glass Production 1/3

# 2.A.3 - Glass Production

### **Short description**

Method	ΑD	EF	<b>Key Category</b>
T2	AS	CS	<b>T:</b> SO <sub>2</sub> , TSP

T = key source by Trend L = key source by Level

Methods	
D	Default
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
Т3	Tier 3 / Detailed Methodology *
С	CORINAIR
CS	Country Specific
M	Model
	Model State of the Action of t

\* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in the group specific chapters.

	•				
AD - Data Source for Activity Data					
NS	National Statistics				
RS	RS Regional Statistics				
IS	International Statistics				
PS	Plant Specific data				
As	Associations, business organisations				
Q	specific Questionnaires (or surveys)				
М	Model / Modelled				
С	Confidential				

EF - Emission Factors						
D	Default (EMEP Guidebook)					
С	Confidential					
CS	Country Specific					
PS	Plant Specific data					
М	Model / Modelled					

Germany's glass industry produces a wide range of different glass types that differ in their chemical composition. Germany's glass sector comprises the following sub-sectors: container glass, flat glass, domestic glass, special glass and mineral fibres (glass and stone wool). The largest production quantities are found in the sectors of container glass and flat glass. Further processing and treatment of glass and glass objects are not considered. Information about the key source relevance can be found in 2.A - Mineral Industry.

# Methodology

The emissions are calculated via a higher Tier method resembling a Tier 2 method, as the activity rates are tied to specific emission factors for different glass types.

### **Activity data**

The production figures are taken from the regularly appearing annual reports of the Federal Association of the German Glass Industry (Bundesverband Glasindustrie; BV Glas). "Production" refers to the amount of glass produced, which is considered to be equivalent to the amount of glass melted down.

2.A.3 - Glass Production 2/3

#### **Emission factors**

The procedure used to determine emission factors for the various glass types involved and the pertinent emissions is described in detail in reports of research projects (Report-No. 001264, search "UBA-FB 001264" in [https://doku.uba.de]  $\Rightarrow$  OPAC  $\Rightarrow$  Signatur). The emission factors were calculated for the various industry sectors. The factors vary annually in keeping with industry monitoring, not only as steady trends, but as time ranges. Ranges below are given as averages over all glass types for main pollutants, but as averages over time for heavy metals:

Table 1: Overview of applied emission factors

~ pollutant	~ products	~ EF	~ unit	~ current Trend
NO,,x,,	all glass types	> 1.0-3.4	kg/t	constant
SO,,2,,	all glass types	> 0.39-1.9	kg/t	constant
NMVOC	all glass types	> 0.96	kg/t	constant
NH,,3,,	all glass types	> 0.03-0.7	kg/t	constant
TSP	all glass types	> 0.02-0.04	kg/t	constant
PM,,10,,	all glass types	> 0.01-0.03	kg/t	constant
PM,,2.5,,	all glass types	> 0.01-0.02	kg/t	constant
As	container glass	> 0.04-0.13	g/t	rising
Pb	container glass	> 0.22-0.41	g/t	rising
Cd	container glass	> 0.01-0.04	g/t	rising
Cr	container glass	> 0.03-0.07	g/t	falling
Cu	container glass	> 0.10-0.23	g/t	falling
Ni	container glass	> 0.01-0.02	g/t	erratic
Se	container glass	> 0.8-1.9	g/t	erratic

### **Trends in emissions**

Trends in emissions correspond to trends of emission factors and of activity data. The resulting trends are not constant, but a complex result of different EF for various glass types. So emissions of SO,,2,, could rising due to increased production Level of a relevant product. gallery size="medium" viewer="yes": EM 2A3 since 1990.PNG gallery

### Recalculations

Recalculations were necessary due to updated activity data for the last reported year.

[!- With **activity data** and **emission factors** remaining unrevised, no recalculations have been carried out compared to last year's submission. -]



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

## **Planned improvements**

For purposes of updating the EF project has started in 2019, results are planned in 2020 1.

[!- At the moment, no category-specific improvements are planned.-]

[!- FAQ No comments!-]

2.A.3 - Glass Production 3/3

 $\frac{\text{bibliography}}{\text{Embissions faktoren für Luftschadstoffe in den Branchen}}{\text{Zementklinker produktion und Glasherstellung}}^{\text{Hilbertand FKZ}}$ 

bibliography

1) (bibcite 1)