# 2.A.3 - Glass Production

### **Short description**

Category Code	•	Method				AD						EF				
2.A.3 T			T	2	AS					CS						
Key Category	SO2	NO×	NH₃	NMVO	CO	BC	Pb	Hg	Cd	Diox	PAH	HCB	TSP	PM10	PM2 5	
2.A.3	L/-	-/-	-/-	-/-	-	-	-/-	-	-/-	-	-	-	-/-	-/-	-/-	
<b>T</b> = key source	by Tr	end <b>I</b>	. = k	ey sourc	e by	Lev	el									
Methods																
				efault												
Т1				Tier 1 / Simple Methodology *												
T2				Tier 2*												
Т3						tail	ed N	1eth	nodolo	ogy *						
C				CORINAIR												
				Country Specific												
	М				odel											
* as described i					on In	ven	tor	y Gu	ide	book	- 2019	9, in t	he gi	roup s	specific	
AD - Data Sou			ctivi	ty Data												
NS National Sta																
	Regional Statistics															
IS International Statistics																
PS Plant Specific data																
As Association:			-		_											
<b>Q</b> specific Que			s (or	surveys	)											
	M Model / Modelled			_												
C Confidentia																
EF - Emission																
<b>D</b> Default (EM		ideb	ook)													
<b>C</b> Confidential																
CS Country Spe																
PS Plant Specif		а														
Model / Mod	elled															

Germany's glass industry produces a wide range of different glass types that differ in their chemical composition. Germany's glass sector comprises the following sub-sectors: container glass, flat glass, domestic glass, special glass and mineral fibres (glass and stone wool). The largest production quantities are found in the sectors of container glass and flat glass. Further processing and treatment of glass and glass objects are not considered. Information about the key source relevance can be found in 2.A - Mineral Industry.

## Methodology

The emissions are calculated via a higher Tier method resembling a Tier 2 method, as the activity rates are tied to specific emission factors for different glass types.

### Activity data

The production figures are taken from the regularly appearing annual reports of the Federal Association of the German Glass Industry (Bundesverband Glasindustrie; BV Glas). "Production" refers to the amount of glass produced, which is considered to be equivalent to the amount of glass melted down.

#### **Emission factors**

The procedure used to determine emission factors for the various glass types involved and the pertinent emissions is described in detail in reports of research projects (Report-No. 001264, search "UBA-FB 001264" in (https://doku.uba.de  $\Rightarrow$  OPAC  $\Rightarrow$  Signatur). The emission factors were calculated for the various industry sectors. The factors vary annually in keeping with industry monitoring, not only as steady trends, but as time ranges. Ranges below are given as averages over all glass types for main pollutants, but as averages over time for heavy metals:

Table 1: Overview of applied emission factors

Pollutant	Products	EF	Unit	Current trend
NOx	all glass types	1.0-3.4	kg/t	constant
<b>SO</b> <sub>2</sub>	all glass types	0.39-1.9	kg/t	constant
NMVOC	all glass types	0.96	kg/t	constant
NH₃	all glass types	0.03-0.7	kg/t	constant
TSP	all glass types	0.02-0.04	kg/t	constant
<b>PM</b> 10	all glass types	0.01-0.03	kg/t	constant
PM2.5	all glass types	0.01-0.02	kg/t	constant
As	container glass	0.04-0.13	g/t	rising
Pb	container glass	0.22-0.41	g/t	rising
Cd	container glass	0.01-0.04	g/t	rising
Cr	container glass	0.03-0.07	g/t	falling
Cu	container glass	0.10-0.23	g/t	falling
Ni	container glass	0.01-0.02	g/t	erratic
Se	container glass	0.8-1.9	g/t	erratic

### **Trends in emissions**

Trends in emissions correspond to trends of emission factors and of activity data. The resulting trends are not constant, but a complex result of different EF for various glass types. So emissions of SO<sub>2</sub> could rising due to increased production Level of a relevant product.

Invalid Link Emission trends in NFR 2.A.3

## Recalculations

Recalculations were necessary due to updated activity data for the last reported year.



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

## **Planned improvements**

For purposes of updating the EF project has started in 2019, results are planned in 2020<sup>1</sup>).

<sup>1)</sup> ReFoPlan FKZ – 3719 52 1010: "Überarbeitung der Emissionsfaktoren für Luftschadstoffe in den Branchen Zementklinkerproduktion und Glasherstellung"