2.A.2 - Lime Production 1/3

2.A.2 - Lime Production

Short description

Category Code		Method				AD				EF					
2.A.2			T1	L		AS CS									
Key Category	SO ₂	NO×	ΝН₃	NMVOC	СО	ВС	Pb	Hg	Cd	Diox	PAH	нсв	TSP	PM ₁₀	PM ₂ 5
2.A.2	-/-	-/-	-	-/-	-	-	-	-/-	-	-	-	-	-/-	-/-	-/-

T = key source by Trend L = key source by Level

Default
Tier 1 / Simple Methodology *
Tier 2*
Tier 3 / Detailed Methodology *
CORINAIR
Country Specific
Model

* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in the group specific chapters.

ш		
	ΑD	- Data Source for Activity Data
	NS	National Statistics
	RS	Regional Statistics
	IS	International Statistics
	PS	Plant Specific data
	As	Associations, business organisation
	Q	specific Questionnaires (or surveys)
I	М	Model / Modelled
	С	Confidential

EF	- Emission Factors
D	Default (EMEP Guidebook)
С	Confidential
CS	Country Specific
PS	Plant Specific data
М	Model / Modelled

The statements made below regarding source category 2.A.2 refer solely to the amounts of burnt lime and dolomite lime produced in German lime works. Other lime-producing processes are included in NFR 2.C.1 and 2.H.2.

Because of the wide range of applications covered by the sector's products, lime production is normally more isolated from economic fluctuations than is production of other mineral products such as cement. Production has fluctuated relatively little since the end of the 1990s. Dolomite-lime production, of which significantly smaller amounts are produced, basically exhibits similar fluctuations.

Methodology

The pertinent emissions level is obtained by multiplying the amount of product in question (quick lime or dolomite lime) and the relevant emission factor.

Activity data

The German Lime Association (BVK) collects the production data for the entire time series on a plant-specific basis, and makes it available for reporting purposes. Production amounts are determined via several different concurrent procedures; their quality is thus adequately assured (Tier 2). Most companies are also required to report lime-production data within the

2.A.2 - Lime Production 2/3

framework of CO_2 -emissions trading. The EU monitoring guidelines for emissions trading specify a maximum accuracy of 2.5%. It is additionally assumed that 2% of the burnt lime is separated as dust in all years of the reporting period from 1990 onwards via appropriate exhaust gas purification systems and is not returned to the production process. This is taken into account by a potential 2% increase in activity rates.

Emission factors

Due to recommendation during NEC-Review 2021 the calculation of CO emissions from lime production is allocated to process emissions based on default-EF. The other EF are country-specific values.

Table 1: Emission factors for quick-lime production

pollutant	Name of Category	EF	unit	Trend
NO _x	quicklime	0.59	kg/t	falling
SO ₂	quicklime	0.12	kg/t	falling
NMVOC	quicklime	0.041	kg/t	constant
СО	quicklime	1.940	kg/t	default
TSP	quicklime	0.050	kg/t	falling
PM ₁₀	quicklime	0.038	kg/t	falling
PM _{2.5}	quicklime	0.023	kg/t	falling
Hg	quicklime	2.62	mg/t	falling

Table 2: Emission factors for dolomite production

pollutant	Name of Category	EF	unit	Trend
NO _x	dolomite	1.73	kg/t	falling
SO ₂	dolomite	0.58	kg/t	falling
NMVOC	dolomite	0.041	kg/t	constant
СО	dolomite	1.940	kg/t	default
TSP	dolomite	0.034	kg/t	falling
PM ₁₀	dolomite	0.026	kg/t	falling
PM _{2.5}	dolomite	0.015	kg/t	falling
Hg	quicklime	2.63	mg/t	falling


Trends in emissions

All trends in emissions correspond to trends of emission factors in table above. No rising trends are identified.

2.A.2 - Lime Production 3/3

trends of emissions of lime industry

Emissions by pollutant / Emissionen nach Schadstoff



^{*} Base Year for PM = 1995 / Basisjahr für Feinstäube (PM) ist 1995

Emission trends in NFR 2.A.2

Recalculations

With **activity data** and **emission factors** remaining unrevised, no recalculations have been carried out compared to last year's submission.



For pollutant-specific information on recalculated emission estimates for Base Year and 2018, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

Source: German Emission Inventory (15.01.2023)

Planned improvements

At the moment, no category-specific improvements are planned.