# 2.D.3.f - Dry Cleaning

# **Short Description**

Category Code Met				ł		AD					EF				
2.D.3.f		-		Ν	NS			CS							
	NO <sub>x</sub>	NMVOC	SO <sub>2</sub>	NH3	PM <sub>2.5</sub>	<b>PM</b> <sub>10</sub>	TSP	BC	CO	Pb	Cd	Hg	Diox	PAH	НСВ
Key Categyory:	-	-/-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>T</b> = key source b	by Tre	end $\mathbf{L} = \mathbf{k}$	ey s	ourc	e by L	evel									
Methods															
	D			_	fault										
	т1				er 1 / S	Simple	e Met	hod	olog	y *					
	Т2				er 2*										
	Т3			_	er 3 / E		ed Me	etho	dolo	gy :	*				
	С			_	RINAI										
	CS			_	untry	Speci	fic								
	Μ				del										
* as described ir					n Inve	entory	Guic	lebo	ok -	20	19,	in t	he gr	oup s	pecif
AD - Data Sou			ty D	ata											
NS National Sta															
RS Regional Sta															
IS Internationa															
PS Plant Specifi															
As Associations		-													
<b>Q</b> specific Que		naires (or	sur	veys	<u>)</u>										
M Model / Mod	elled				_										
<b>C</b> Confidential															
EF - Emission I															
<b>D</b> Default (EME	P Gu	idebook)													
<b>C</b> Confidential															
CS Country Spe															
PS Plant Specifie		а													
M Model / Mode	اء ما ا														

This source category comprises NMVOC emissions from **Solvent application for professional textile cleaning**. The German inventory summarizes hydrocarbon solvents and perchloroethylene as solvent.

'NMVOC' is defined in accordance with the VOC definition found in the EC solvents directive. For purposes of the definition of solvents, the term 'solvent use' is also defined in accordance with the EC solvents directive.

# Method

#### **General procedure**

NMVOC emissions are calculated in accordance with a product-consumption-oriented approach. In this approach, solventbased products or solvents are allocated to the source category, and then the relevant NMVOC emissions are calculated from those solvent quantities via specific emission factors. Thus, the use of this method is possible with the following valid input figures for each product group:

- Quantities of VOC-containing (pre-) products and agents used in the report year,
- The VOC concentrations in these products (substances and preparations),

• The relevant application and emission conditions (or the resulting specific emission factor).

The quantity of the solvent-based (pre-)product corresponds to the domestic consumption which is the sum of domestic production plus import minus export.

NMVOC Emission = domestic consumption of a certain product \* solvent content \* specific emission factor

The calculated NMVOC emissions of different product groups for a source category are then aggregated. The product / substance quantities used are determined at the product-group level with the help of production and foreign-trade statistics. Where possible, the so-determined domestic-consumption quantities are then further verified via cross-checking with industry statistics.

# **Discussion of emission trends**

#### **General information**

Since 1990, so the data, NMVOC emissions from use of solvents and solvent-containing products in general have decreased by nearly 55%. The main emissions reductions have been achieved in the years since 1999. This successful reduction has occurred especially as a result of regulatory provisions such as the 31st Ordinance on the execution of the Federal Immissions Control Act (Ordinance on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain facilities – 31. BImSchV), the 2nd such ordinance (Ordinance on the limitation of emissions of highly volatile halogenated organic compounds – 2. BImSchV) and the TA Luft.

#### **Specific information**

Until 1999, data of the present source categories 2.D.3.e and f were treated as one source group. Source group 2.D.3.f accounts for about 0.2% of total NMVOC emissions from solvent-based products and remained stable in the last 15 years.

### Uncertainties

The overall uncertainty of emissions caused by applications of this source group is estimated at 50%.

# Recalculations

Routinely the NMVOC emissions of the last reported year must be actualized in the next reporting cycle as the final data of the foreign trade statistics are regularly only available after the publication of the respective reporting year has been completed.

For the year 2020, this adjustment is postponed to the submission 2024 because for some applications the calculation bases also have to be reviewed for further years and this review could not be completed in time for the 2023 reporting.



For **pollutant-specific information on recalculated emission estimates for Base Year and 2019**, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

# **Planned improvements**

At the moment, no category-specific improvements are planned.