

11.B - Forest Fires

Short Description

In Germany's forests prescribed burning is not applied. Therefore, all forest fires are categorized as wildfires (include emissions from forest fires occurring naturally or caused by humans). - Note that emissions reported here are not accounted for the national totals.

Method	AD	EF	Key Category
CS, T2, T1	CS	D	<i>not included in key category analysis</i>

Legend T = key source by Trend / L = key source by Level

Methods D: Default RA: Reference Approach T1: Tier 1 / Simple Methodology * T2: Tier 2* T3: Tier 3 / Detailed Methodology
* C: CORINAIR CS: Country Specific M: Model as described in the EMEP/CORINAIR Emission Inventory Guidebook - 2019, in the group specific chapters.

AD:- *Data Source for Activity Data* NS: National Statistics RS: Regional Statistics IS: International Statistics PS: Plant Specific data AS: Associations, business organisations Q: specific questionnaires, surveys

EF - Emission Factors D: Default (EMEP Guidebook) C: Confidential CS: Country Specific PS: Plant Specific data

Methodology

For calculating the emissions of wildfires a country specific Tier2 approach was used. The mass of carbon emitted $M(C)$ was calculated using the adapted equation follows the methodology of Seiler and Crutzen (1980) ¹⁾.



$$M(C) = 0.45 * A * B * \beta$$

where:

0.45 = average fraction of carbon in fuel wood;

A = forest area burnt in [m^2];

B = mean above-ground biomass of fuel material per unit area in [kg/m^2];

β = burning efficiency (fraction burnt) of the above-ground biomass.

The data on forest areas burnt for the period 1990 to 2021 have been taken from the German forest fire statistic (BLE, 2022)²⁾ managed by the Federal Agency for Agriculture and Food. The mean above-ground biomass of fuel material was determined from the pools above ground biomass, dead wood and litter. The mean above-ground biomass and dead wood biomass was derived for each year by linear extrapolation and interpolation between the

- German National Forest Inventories of 1987, 2002, 2012 (Bundeswaldinventuren 1987, 2002, 2012),
- [the inventory study 2008](#) and,
- [the carbon inventory 2017](#).

Biomass of Litter was derived for each year by linear interpolation between 1990 and 2006 and extrapolation from 2007 based on the both Forest soil inventories (BZE I Wald (1990)³⁾ and BZE II Wald (2006)⁴⁾.

Pursuant to König (2007)⁵⁾, 80% of the forest fires in Germany are surface fires and 20% crown fires. In accordance to the IPCC Good Practice Guidance for LULUCF (2003) a burning efficiency of 0.15 was used for surface fires and an efficiency of 0.45 was used for crown fires.

The emissions for the pollutants were calculated by multiplying the mass of carbon with the respective emission factors from table 3-3 (EMEP/EEA, 2019)⁶⁾.

For the calculation of particulate emissions (TSP, PM_{10} and $\text{PM}_{2.5}$) the burnt biomass was multiplied with the respective emission factors from table 3-5 (EMEP/EEA, 2019). Those particulate emission factors have been estimated by averaging the emission factors from the US Environmental Protection Agency (USEPA, 1996)⁷⁾ methodology, since no better information is available. Those emission factors are assumed to be the same for all types of forest.

The Guidebook does not indicate whether EFs have considered the condensable component (with or without).

Activity data

The data on forest areas burnt for the period 1990 to 2021 are based on the German forest fire statistic (BLE, 2021) managed by the Federal Agency for Agriculture and Food.

Table 1: Area of forest burnt from 1990 until the latest reporting year, in [ha]

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
1,606	920	4,908	1,493	1,114	592	1,381	599	397	415	581
2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
122	122	1,315	274	183	482	256	539	757	522	214
2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
269	199	120	526	283	395	2,349	2,711	368	148	

Emission factors

For the year 2021 the estimated emission factors from table 2 were applied.

Table 2: Emission factors applied for 2021

	EF₂₀₂₁
NO _x	155.19
CO	5,535.19
NMVOC	488.86
SO _x	37.25
NH ₃	41.9
TSP	879.42
PM ₁₀	569.04

	EF₂₀₂₁
PM _{2.5}	465.58
BC	41.90

In addition, a large-scale fire, which occurred in September 2018, is reported under 11.B. A detailed description can be found in the NIR 2020 in Chapter 6.8.2.5 ⁸⁾, because a large amount of CO₂ emissions were released.

The burned area of the drained moor, which is used as a military facility, covered 1,221 ha. This fire was extensively investigated and documented by the Federal Office for Infrastructure, Environmental Protection and Services of the German Armed Forces. The emissions are calculated according to IPCC GL (2006), chapter 2, form 2.27 ⁹⁾.

The product $M_b \times C_f$ is set to 336 t d.m. ha⁻¹ according to Table 2.6 and equation 2.7, 2013 IPCC Wetlands Supplement ¹⁰⁾, i.e. it is assumed that the moor was completely drained during the fire.



- M_b = mass of fuel available for combustion, tonnes ha⁻¹ (i.e. mass of dry organic soil fuel)
- C_f = combustion factor, dimensionless

For calculating CO emissions, the EF according to Table 2.7, 2013 IPCC Wetlands Supplement 207 g (kg dm)⁻¹, is taken into account, resulting in 85 kt CO. For other emissions from land fires on drained organic soils no Tier-1 emission factors exist and emissions are therefore not reported (NO).

Recalculations

Recalculations were made for the complete time series due to the methodology changes (the inclusion of the burning biomass of dead wood and litter, which has not been considered until now). No recalculation was made for the large-scale fire in 2018.

Table 3: Recalculation of air pollutant emissions from 1990 until the latest reported year, in [kt]

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
Black Carbon (BC)											
current submission	0,061	0,022	0,022	0,007	0,020	0,021	0,011	0,016	0,096	0,112	0,015
previous submission	0,047	0,017	0,017	0,005	0,016	0,017	0,009	0,013	0,078	0,091	0,013
absolute change	0,014	0,005	0,005	0,001	0,004	0,004	0,002	0,003	0,018	0,020	0,003
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Carbon monoxide											
current submission	8,043	2,949	2,879	0,911	2,641	2,771	1,506	2,114	12,684	14,762	2,019
previous submission	6,165	2,284	2,253	0,722	2,125	2,257	1,228	1,724	10,357	12,067	1,652
absolute change	1,878	0,665	0,627	0,189	0,515	0,515	0,278	0,390	2,327	2,695	0,367
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Ammonia											
current submission	0,061	0,022	0,022	0,007	0,020	0,021	0,012	0,016	0,098	0,112	0,015
previous submission	0,047	0,017	0,017	0,005	0,016	0,017	0,009	0,013	0,078	0,091	0,013
absolute change	0,014	0,005	0,005	0,001	0,004	0,004	0,002	0,003	0,020	0,020	0,003
relative change	30,0%	28,7%	27,7%	26,8%	27,2%	25,8%	25,5%	25,7%	25,5%	22,3%	22,2%
Non-Methane Volatile Organic Compounds (NMVOC)											
current submission	0,710	0,260	0,254	0,080	0,233	0,245	0,133	0,187	1,120	1,304	0,178
previous submission	0,545	0,202	0,199	0,064	0,188	0,199	0,108	0,152	0,915	1,066	0,146
absolute change	0,166	0,059	0,055	0,017	0,046	0,045	0,025	0,034	0,205	0,238	0,032
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Nitrogen oxides											
current submission	0,226	0,083	0,081	0,026	0,074	0,078	0,042	0,059	0,356	0,414	0,057
previous submission	0,173	0,064	0,063	0,020	0,060	0,063	0,034	0,048	0,290	0,338	0,046
absolute change	0,053	0,019	0,018	0,005	0,014	0,014	0,008	0,011	0,065	0,076	0,010

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Particulate Matter <10µm (PM ₁₀)											
current submission	0,827	0,303	0,296	0,094	0,271	0,285	0,155	0,217	1,304	1,518	0,208
previous submission	0,634	0,235	0,232	0,074	0,218	0,232	0,126	0,177	1,065	1,241	0,170
absolute change	0,193	0,068	0,064	0,019	0,053	0,053	0,029	0,040	0,239	0,277	0,038
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Particulate Matter <2.5µm (PM _{2.5})											
current submission	0,677	0,248	0,242	0,077	0,222	0,233	0,127	0,178	1,067	1,242	0,170
previous submission	0,519	0,192	0,189	0,061	0,179	0,190	0,103	0,145	0,871	1,015	0,139
absolute change	0,158	0,056	0,053	0,016	0,043	0,043	0,023	0,033	0,196	0,227	0,031
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Sulphur dioxide											
current submission	0,054	0,020	0,019	0,006	0,018	0,019	0,010	0,014	0,085	0,099	0,014
previous submission	0,041	0,015	0,015	0,005	0,014	0,015	0,008	0,012	0,070	0,081	0,011
absolute change	0,013	0,004	0,004	0,001	0,003	0,003	0,002	0,003	0,016	0,018	0,002
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%
Total suspended particles (TSP)											
current submission	1,278	0,469	0,457	0,145	0,420	0,440	0,239	0,336	2,015	2,345	0,321
previous submission	0,980	0,363	0,358	0,115	0,338	0,359	0,195	0,274	1,646	1,917	0,262
absolute change	0,298	0,106	0,100	0,030	0,082	0,082	0,044	0,062	0,370	0,428	0,058
relative change	30,5%	29,1%	27,8%	26,3%	24,2%	22,8%	22,7%	22,6%	22,5%	22,3%	22,2%

¹⁾ Seiler, Wolfgang, and Paul J. Crutzen. "Estimates of gross and net fluxes of carbon between the biosphere and the atmosphere from biomass burning." Climatic change 2.3 (1980): 207-247.

²⁾ BLE (Bundesanstalt für Landwirtschaft und Ernährung), (2022, 30. Juni), 2022: Waldbrandstatistik der Bundesrepublik Deutschland für das Jahr 2021, Bonn: 21 p. Retrieved July 2022,
https://www.ble.de/DE/BZL/Daten-Berichte/Wald/wald_node.html

³⁾ WOLFF, B. & RIEK, W. (1997): Deutscher Waldbodenbericht 1996 - Ergebnisse der bundesweiten Bodenzustandserhebung im Wald (BZE) 1987 - 1993. Hrsg.: Bundesministerium für Ernährung, Landwirtschaft und Forsten, Bonn, Bd. 1 u. 2., 144 S.,
<https://www.bmel-statistik.de/fileadmin/daten/FHB-0320205-1996.pdf>

⁴⁾ WELLBROCK, N., AYDIN, C.-T., BLOCK, J., BUSSIAN, B., DECKERT, M., DIEKMANN, O., EVERS, J., FETZER, K. D., GAUER, J., GEHRMANN, J., KÖLLING, C., KÖNIG, N., LIESEBACH, M., MARIN, J., MEIWES, K. J., MILBERT, G., RABEN, G., RIEK, W., SCHÄFFER, W., SCHWERHOFF, J., ULLRICH, T., UTERMANN, J., VOLZ, H.-A., WEIGEL, A. & WOLFF, B. (2006): Bodenzustandserhebung im Wald (BZE II) Arbeitsanleitung für die Außenaufnahmen. Bundesministerium für Ernährung, Landwirtschaft und Verbraucherschutz, Berlin, 413 S.
<https://www.bmel.de/DE/themen/wald/wald-in-deutschland/bodenzustandserhebung.html>

⁵⁾ König, H.-C., 2007. Waldbrandschutz - Kompendium für Forst und Feuerwehr. 1. Fachverlag Matthias Grimm, Berlin, 197 S.

⁶⁾ EMEP/EEA, 2019:
<https://www.eea.europa.eu/publications/emep-eea-guidebook-2019/part-b-sectoral-guidance-chapters/11-natural-sources/11-b-forest-fires/view>

⁷⁾ USEPA, 1996: Compilation of Air Pollutant Emission Factors Vol.1. Stationary, Point and Area Sources. Report AP-42, fifth edition

⁸⁾ NIR (2020): National Inventory Report 2020 for the German Greenhouse Gas Inventory 1990-2018. Available in April 2020

⁹⁾ IPCC (Intergovernmental Panel on Climate Change) (2006): 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4: Agriculture, Forestry and Other Land Use. Eds.: Eggleston S., Buendia L., Miwa K., Ngara T., Tanabe K. (Eds). IEA/OECD, IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.
<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>

¹⁰⁾ IPCC (Intergovernmental Panel on Climate Change) (2014b): 2013 Supplement to the IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G.(eds). Published: IPCC, Switzerland <http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html>