5.D.1 - Domestic & Commercial Wastewater Handling

Short description

Category Code	Method				AD				EF						
5.D.1	T1				NS					D					
Key Category	SO2	NO×	ΝНз	NMVOC	СО	ВС	Pb	Hg	Cd	Diox	PAH	нсв	TSP	PM ₁₀	PM ₂ 5
5.D.1	-	-	-	-/-	-	-	-	-	-	-	-	-	-	-	-

T = key source by Trend L = key source by Level

Default
Tier 1 / Simple Methodology *
Tier 2*
Tier 3 / Detailed Methodology *
CORINAIR
Country Specific
Model

^{*} as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in the group specific chapters.

	The state of the s
AD	- Data Source for Activity Data
NS	National Statistics
RS	Regional Statistics
IS	International Statistics
PS	Plant Specific data
As	Associations, business organisations
Q	specific Questionnaires (or surveys)
М	Model / Modelled
С	Confidential

EF	- Emission Factors
D	Default (EMEP Guidebook)
С	Confidential
CS	Country Specific
PS	Plant Specific data
М	Model / Modelled

In category **5.D.1**, <u>NMVOC emissions</u> from domestic and commercial wastewater handling are reported. The domestic section is covered by wastewaters of municipal origin (large centralised plants; ranging from 1000 up to >100.000 resident values). The commercial section is covered by industrial and commercial wastewaters, co-treated in municipal wwt-plants.

Method

Emissions reported under this category are calculated using the Tier 1 approach of the EMEP/EEA Guidebook 2016, where the emission factor (EF) is 15 mg/m³ wastewater (Part B, 5.D, chap. 3.2.2, Table 3-1, p. 7). This EF is multiplied with the total amount of wastewater (AD) treated in domestic and commercial wwt-plants, following the equation:

Emissions (NMVOC) = **AD** x **EF** (ibid., chap. 3.2.1)

Activity data

Total volumes of treated municipal wastewater are derived by the German statistical agency (Statistisches Bundesamt, Fachserie 19, Reihe 2.1.2 ¹⁾). The data source is published on a three-year basis with new data only for the respective year of the update. The availability of the data starts in 1991 with an exception for the following update, which was for 1995.

Missing data are inter- or extrapolated

Emisson factors

See method

Uncertainties

The AD from Statistisches Bundesamt have an uncertainty of $\pm 3\%$ (normal distribution) whereas the uncertainty for the EF, due to its range (5/50 mg/m³), is -70 / +210 % and the distribution lognormal.

Recalculations

Recalculations were not necessary

Planned improvements

Currently no improvements are planned.

1)

Statistisches Bundesamt, Fachserie 19, Reihe 2.1.2