

Explanation of Key Trends - Heavy Metals



Please note: Data for heavy metals may have issues such as missing sources. It features considerably higher uncertainties than data for other pollutants covered in this report.

Obligations

The 1998 Aarhus Protocol on Heavy Metals under the CLRTAP entered into force late in 2003. It targets three particularly harmful metals: cadmium, lead and mercury. According to one of the basic obligations, Germany has to reduce its emissions for these three metals below their levels in 1990. The Protocol aims to cut emissions from industrial sources (iron and steel industry, non-ferrous metal industry), combustion processes (power generation, road transport) and waste incineration. It defines stringent limit values for emissions from stationary sources and suggests best available techniques (BAT) for these sources, such as special filters or scrubbers for combustion sources or mercury-free processes. The Protocol requires Parties to phase out leaded petrol. It also introduces measures to lower heavy metal emissions from other products, such as mercury in batteries, and proposes the introduction of management measures for other mercury-containing products, such as electrical components (thermostats, switches), measuring devices (thermometers, manometers, barometers), fluorescent lamps, dental amalgam, pesticides and paint.


Main drivers

Emission of priority heavy metals (cadmium, lead and mercury) decreased significantly since 1990. Values show reductions by about 58 to 89% compared to the base year. Most of the achievements originate from the early 1990's though. Recent years saw hardly any improvements in heavy metal emissions.

The main sources for most heavy metals is fuel combustion: Energy Industries (NFR 1.A.1) and Industrial Processes (NFR 2) like Metal Industries (NFR 2.C) emit the majority of arsenic, cadmium, chrome, lead, mercury and Nickel. But copper and zinc emissions are mostly governed by the Transport (NFR 1.A.3) sector, resulting mostly from brake and tyre wear. Thus, trends are connected directly with the annual mileage. Selenium on the other hand originates mainly from Mineral Industry (NFR 2.A) and to a lesser degree from Transport (NFR 1.A.3). Other sources are still to be investigated but generally expected to add little to the total trend.

Trends

The figure below shows emission trends for heavy metals:



Heavy metal emission trends