

1.B.1 - Solid Fuels



Category Code	Method	AD	EF												
1.B.1.a	T2, M	AS	CS												
1.B.1.b	T2, T3	AS	CS												
Key Category	NO _x	NM VOC	SO ₂	NH ₃	PM _{2.5}	PM ₁₀	TSP	BC	CO	PB	Cd	Hg	Diox	PAH	HCB
1.B.1.a	-	-/-	-	-	-/-	-/-	-/-	-	-	-	-	-	-	-	-
1.B.1.b	-/-	-/-	-/-	-/-	-/-	-/-	L/T	-/-	-/-	-	-	-/-	-/-	-/T	-

Method(s) applied	
D	Default
T1	Tier 1 / Simple Methodology *
T2	Tier 2*
T3	Tier 3 / Detailed Methodology *
C	CORINAIR
CS	Country Specific
M	Model
* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in category chapters.	
(source for) Activity Data	
NS	National Statistics
RS	Regional Statistics
IS	International Statistics
PS	Plant Specific
As	Associations, business organisations
Q	specific Questionnaires (or surveys)
M	Model / Modelled
C	Confidential
(source for) Emission Factors	
D	Default (EMEP Guidebook)
CS	Country Specific
PS	Plant Specific
M	Model / Modelled
C	Confidential

The source category Solid fuels (1.B.1) consists of two sub-source subcategories - the source subcategory Coal mining (1.B.1.a) and the source subcategory Coal transformation (1.B.1.b). This chapter discusses fugitive emissions from coal mining, coal handling, including door leakages from coke ovens and quenching (emissions from the furnace are covered by

Emissions from lignite production other than listed below are reported by plant operator. Particle emission factors were verified in a research project (Emissionen und Maßnahmenanalyse Feinstaub 2000-2020) ⁴⁾.

Table 4: Emission factors applied for lignite-coke production

Pollutant	Unit	Value
TSP	kg/t	0.1
PM ₁₀	kg/t	0.048
PM _{2.5}	kg/t	0.013
PAH	mg/t	55
PCDD/F	µg/t	0.03

Hard coal coke production

The activity rates for hard coal coke production have been taken from the *Statistik der Kohlenwirtschafts's* website (in German only) ⁵⁾.

Table 5: Annual amounts of hard coal coke produced, in [Mt]

1990	1995	2000	2005	2010	2015	2020	2021
18.5	11.1	9.1	8.4	8.2	8.8	7.9	8.2

The emission factors for hard coal coke production have been obtained from the research project "Emission factors for the iron and steel industry, for purposes of emissions reporting" ("Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung") ⁶⁾.

Table 6: Emission factors for hard coal coke production

Pollutant	Unit	Value
CO	kg/t	0.015
NH ₃	kg/t	0.000243
NMVOC	kg/t	0.096
SO ₂	kg/t	0.004
TSP	kg/t	0.011
PM ₁₀	kg/t	0.004
PM _{2.5}	kg/t	0.004
PAH	mg/t	55
PCDD/F	µg/t	0.0015

There are many potential sources of PAH emissions from coking plants. The dominant emission sources are leakages from coke oven doors and from charging operations. As there is limited data available on PAH emissions, the uncertainties of the estimated emission factors are very high. It should also be taken into account that emissions from coke production greatly vary between different coke production plants. The emission factors for benzo[a]pyrene and mixed PAH have been revised by research projects in 2010 ⁷⁾⁸⁾. Split factors for Black Carbon (BC) are based on the EMEP Guidebook 2016 ⁹⁾.

Charcoal production

Small quantities of charcoal are produced in Germany – by one major charcoal-factory operator and in a number of demonstration charcoal kilns. The pertinent quantities are determined by the Federal Statistical Office and are subject to confidentiality requirements. The emission factors were obtained from US EPA 1995 ¹⁰⁾.

Use of charcoal (includes wood only) and barbecue coal (includes wood and lignite briquetts) is reported under [2.G. - Use of Charcoal for barbecues](#). The production of lignite briquettes is reported under 1.B.1.b.

Decommissioned hard-coal mines

NM VOC Emissions from decommissioned hard-coal mines play a role in this sub- source category. When a hard-coal mine is decommissioned, mine gas can escape from neighbouring rock, and from coal remaining in the mine, into the mine's network of shafts and passageways. Since the mine is no longer artificially ventilated, the mine gas collects and can then reach the surface via gas pathways in the overlying rock or via the mine's own shafts and passageways. Such mine gas was long seen primarily as a negative environmental factor. Recently, increasing attention has been given to the gas' positive characteristics as a fuel (due to its high methane content, it is used for energy recovery). In the past, use of mine gas was rarely cost-effective. This situation changed fundamentally in 2000 with the Renewable Energy Sources Act (EEG). Although mine gas is a fossil fuel in finite supply, its use supports climate protection, and thus the gas was included in the EEG. The Act requires network operators to accept, and provide specified compensation for, electricity generated with mine gas and fed into the grid. The NM VOC emissions from decommissioned hard-coal mines have been calculated in the research project "Potential for release and utilisation of mine gas" ("Potential zur Freisetzung und Verwertung von Grubengas")¹¹⁾. The relevant calculations were carried out for all mining-relevant deposits in Germany.

Table 7: NM VOC emission factor for decommissioned hard-coal mines, in [kg/m³]

EF
0.001599

Recalculations



For more details please refer to the super-ordinate chapter [1.B - Fugitive Emissions from fossil fuels](#)

Planned improvements

It is planned to include pant specific data from charcoal production.

References

- ¹⁾ Statistik der Kohlenwirtschaft (2019) [External Link](#) (last pageview: March 2021)
- ²⁾ Dokumentation zur Berechnung des PM10-Austrags aus dem Tagebau Hambach im Jahr 2013 und Ableitung eines Emissionsfaktors (2021) [External Link](#) (last pageview: March 2023)
- ³⁾ Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP) [External Link](#)
- ⁴⁾ Federal Environment Agency research project No. 204 42 202/2 "Emissionen und Maßnahmenanalyse Feinstaub 2000-2020", published in 2007 [External Link](#)
- ⁶⁾ Hensmann et al. 2011
- ⁷⁾ Federal Environment Agency and DFIU research project "Anpassung der deutschen Methodik zur rechnerischen Emissionsermittlung an internationale Richtlinien, Teilbericht Prioritäre Quellen", 2010 (not available online)
- ⁸⁾ Federal Environment Agency and BFI research project No. 3707 42 301 "Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung", 2011 [External Link](#)
- ⁹⁾ EMEP/EEA Air Pollutant Emission Inventory Guidebook 2016; published in 2016 [External Link](#) (last pageview: Dec 2016)
- ¹⁰⁾ Neulicht, R. (1995): Emission Factor Documentation for AP-42 Section 10.7 "Charcoal". [External Link](#)
- ¹¹⁾ Meiners, H. (2014): Potential zur Freisetzung und Verwertung von Grubengas