# 5.C.2 - Open Burning of Waste

<b>Category Code</b>	Method					AD					EF				
5.C.2	CS			Q				D, CS							
Key Category	SO <sub>2</sub>	NOx	ΝН₃	NMVOC	СО	вс	Pb	Hg	Cd	Diox	PAH	нсв	TSP	PM <sub>10</sub>	PM <sub>2</sub> 5
5.C.2	-/-	-/-	-	-/-	-/-	-/-	-/-	-	-/-	-/-	-/-	-	-/-	-/-	-/T

7 7											<u> </u>
Method(s) applied											
D	Default										
T1	Tier 1 / S	imple M	letho	dol	ogy	*					
T2	Tier 2*										
T3	Tier 3 / Detailed Methodology *										
С	CORINAI	R									
CS	Country	Specific									
M	Model										
* as described in the EMEP/E	EA Emissi	on Inve	ntory	/ Gu	idek	ook	- 201	9, in (	categ	ory c	hapters
(source for) Activity Data											
NS	National Statistics										
RS	Regional Statistics										
IS	Internati	International Statistics									
PS	Plant Spe	Plant Specific									
As	Associations, business organisations										
Q	specific Questionnaires (or surveys)										
М	Model / Modelled										
С	Confidential										
(source for) Emission Fac	tors										
D	Default (EMEP Guidebook)										
CS	Country Specific										
PS	Plant Specific										
М	Model / N	Model / Modelled									
С	Confiden	Confidential									

Within NFR sub-category 5.C.2 - Open Burning of Waste, the German emissions inventory provides emissions from registered bonfires and other wooden materials burnt outdoors. Emissions from bonfires are key source for PM2.5 and PM10, but in principle of minor priority due to discontinuous appearance.

Please see chapter regarding farming/plantation waste: https://thg.thuenen.de/iir-de/sector/agriculture/field\_burning/start - this is banned by law in Germany. So there is no gap of reporting.

Emissions from open burning of wood and green waste for traditional purposes, so-called bonfires such as Easter fires, are reported model-based. In addition to biogenic carbon dioxide, emissions of NOx, SO2, CO, NMVOC, particulate matter (PM2.5, PM10 and TSP), Polycyclic Aromatic Hydrocarbons (PAHs) and Heavy Metals are covered so far.

## Method

For developing of a estimation frame a survey regarding the number of such bonfires was carried out by an expert work <sup>1)</sup>. As the result, questionnaires from municipalities and statistical projections for Germany for the year 2016 were checked. The project has shown a declining trend since 1990. On the basis of expert judgement, a further reduction of emissions in the future is expected.

As discussed on Review 2020 regarding all relevant sources: A comparison shows that the volume of bonfires is significantly higher than the volume of campfires. In terms of number, however, the two types of fires are similar. Due to the large fluctuations of the minimum/maximum values, the median was proposed in study. In our view the estimation of bonfires emissions is conservative and completly.

# **Activity data**

Activity data for this category are based on data from a step by step calculation: After the evaluation of the questionaires an extrapolation of the volume and the number of bonfires was made for Germany. For the years since 2019, it became visible that, in addition to the model-based continuous decrease in activities, special aspects must be taken into account: Because of the restrictions on public activities during the pandemic, modeling of less traditional events was searched for. Two types of fires were already classified in the expert project: camp fires in the more private sector and, most importantly, Easter Fires in the more public sector. The calculations are now considered separately and the camp fires are modeled with a continued steady decline. The mainly Easter fires follow an approach about general percentage decreases and additionally in 2019 five percentage points decrease corresponding to various cancels due to forest fire risk. In 2020, an additional 70 percent decrease was modeled due to cancellations for pandemic response (no complete cancellation in Germany because there were exceptions and follow-up events). The following values are the result of evaluation:

Type of fire (quantity of wooden wastes)	Unit	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
All bonfires in sum	t	431,394	414,276	397,157	380,038	362,919	345,800	342,376	339,466	336,556	319,531	135,170

### **Emission factors**

As discussed on Review 2020 regarding EF used and referenced: We use different EF from different references instead the EF of Table 3-1 Tier 1 emission factors for source category 5.C.2 Small-scale waste burning, because the Tier 1 EF seem not suitable for the burning of wooden wastes. We consider both fresh wood (garden and park waste) and dry wood (without coatings etc.). We have tried to find relevant parallels, especially because of the burning of fresh wood with regard to forest fires. But regarding the EF from GB 2019 we will evaluate the use as shown in the following table:


pollutant	Figure	unit	Current reference	Planned improvement
СО		•	GB 2016 small combustion Table 3-6: Tier 1 emission factors for NFR source category 1.A.4.b, using biomass	to use EF from GB 2019 5.C.2, table 3-2: 48.8 kg/ t
NOx		_	Research results from literature: wood burning as it was documented in Ireland's IIR	to use EF from GB 2019 5.C.2, table 3-2: 1.38 kg/ t
SO2	0.2	kg/ t	Research results from literature: wood burning as it was documented in Ireland's IIR	to use EF from GB 2019 5.C.2, table 3-2: 0.03 kg/ t
NMVOC	47.0	kg/ t	not correct used, error in data handling	to use EF from GB 2019 5.C.2, table 3-2: 1.47 kg/ t
TSP	17.0	kg/ t	GB 2016 Forest fires, table 3-1, according 'wood burned'	to use EF from GB 2019 5.C.2, table 3-2: 4.31
PM10	11.0	kg/ t	GB 2016 Forest fires, table 3-1, according 'wood burned'	to use EF from GB 2019 5.C.2, table 3-2: 4.13 kg/ t
PM2.5	9.0		GB 2016 Forest fires, table 3-1, according 'wood burned'	to use EF from GB 2019 5.C.2, table 3-2: 3.76 kg/ t
ВС	0.81	kg/ t	GB 2016 Forest fires, table 3-1, according 'wood burned'	to use EF from GB 2019 5.C.2, table 3-2: 28% of PM2.5
PCDD/F	10.0	μg/ t	GB 2019 5.C.2, table 3-1	No further
PAH	3.39	g/ t	sum of single compounts	No further
BaP	1.3	g/ t	IIR Ireland 1)	No further (GB with dry matter problem)
BbF	1.5	g/ t	IIR Ireland <sup>2)</sup>	No further (GB with dry matter problem)
BkF	0.5	g/ t	IIR Ireland <sup>3)</sup>	No further (GB with dry matter problem)
IxP	0.09	g/ t	IIR Ireland	No further, Gap in GB
Pb	0.32	g/ t	GB 2019 5.C.2, table 3-2	No further
Cd	0.13	g/ t	GB 2019 5.C.2, table 3-2	No further

# **Trends in emissions**

All trends in emissions correspond to trends of AD. No rising trends are to identify.

## Trends of Emissions in Germany in NFR category Bonfires

Emissions by pollutant / Emissionen nach Schadstoff



<sup>\*</sup> Base Year for PM = 1995 / Basisjahr für Feinstäube (PM) ist 1995

Quelle: German Environment Agency, National inventory for the German reporting on atmospheric emissions since 1990, (11/2023)

#### **Emission trends of bonfires**

# Recalculations

Due to evaluation of activity data recalculations have been carried out compared to last year's submission. Mainly the values of 2019 were changed, all other years only minor.



For pollutant-specific information on recalculated emission estimates for Base Year and 2019, please see the pollutant specific recalculation tables following chapter 8.1 - Recalculations.

<sup>&</sup>lt;sup>1)</sup> EF is referenced to a former research project called 'Use of charcoal, tobacco etc.'. This was a literatur research, which is only available via UBA library in German. The EF is relating wood burning as it was documented in Ireland's IIR

<sup>&</sup>lt;sup>2)</sup> EF is referenced to a former research project called 'Use of charcoal, tobacco etc.'. This was a literatur research, which is only available via UBA library in German. The EF is relating wood burning as it was documented in Ireland's IIR

<sup>&</sup>lt;sup>3)</sup> EF is referenced to a former research project called 'Use of charcoal, tobacco etc.'. This was a literatur research, which is only available via UBA library in German. The EF is relating wood burning as it was documented in Ireland's IIR

Management: Erhebung der Größen und Zusammensetzung von Brauchtums- und Lagerfeuern durch kommunale Befragungen; URL:

 $https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2018-02-19\_texte\_11-2018\_lager-brauchtumsfeuer.pdf; UBA-Texte\_11/2018$