

## 1.B.1 - Solid Fuels



### Short description

Category Code	Method	AD	EF
1.B.1.a	T2, M	AS	CS
1.B.1.b	T2, T3	AS	CS
Method(s) applied			
D	Default		
T1	Tier 1 / Simple Methodology *		
T2	Tier 2*		
T3	Tier 3 / Detailed Methodology *		
C	CORINAIR		
CS	Country Specific		
M	Model		
* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in category chapters.			
(source for) Activity Data			
NS	National Statistics		
RS	Regional Statistics		
IS	International Statistics		
PS	Plant Specific		
As	Associations, business organisations		
Q	specific Questionnaires (or surveys)		
M	Model / Modelled		
C	Confidential		
(source for) Emission Factors			
D	Default (EMEP Guidebook)		
CS	Country Specific		
PS	Plant Specific		
M	Model / Modelled		
C	Confidential		

Key Category	NO <sub>x</sub>	NM VOC	SO <sub>2</sub>	NH <sub>3</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC	CO	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	PCDD/F	B(a)P	B(b)F	B(k)F	I(x)P	PAHs	HCB
1.B.1.a	NA	-/-	NA	NA	-/-	-/T	-/-	NA	NA	NA	NA	NA	-/-	-/-	-/-	-/-	-/-	-/-	NA	-	-	-	-	NA	NA
1.B.1.b	-/-	-/-	-/-	-/-	-/-	-/-	L/T	-/-	-/-	NA	NA	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	L/T	-/T	-/T	-/T	-/T	NA

<b>L/-</b>	key source by <b>Level</b> only
<b>-/T</b>	key source by <b>Trend</b> only
<b>L/T</b>	key source by both <b>Level</b> and <b>Trend</b>
<b>-/-</b>	no key source for this pollutant
<b>IE</b>	emission of specific pollutant <b>Included Elsewhere</b> (i.e. in another category)
<b>NE</b>	emission of specific pollutant <b>Not Estimated</b> (yet)
<b>NA</b>	specific pollutant not emitted from this source or activity = <b>Not Applicable</b>
<b>*</b>	no analysis done

Source category **1.B.1 - Solid fuels** consists of two sub-source subcategories – the source subcategory Coal mining (1.B.1.a) and the source subcategory Coal transformation (1.B.1.b). This chapter discusses fugitive emissions from coal mining, coal handling, including door leakages from coke ovens and quenching (emissions from the furnace are covered by category 1.A.1.c), and emissions from the beneficiation of solid fuels. In the mining sector, a distinction is made between open-pit mines, in which raw materials are extracted from pits open to the surface, and closed-pit mines, in which seams are mined underground. In Germany, hard coal used to be mined in closed-pits only (until 2018), while lignite is mined in four coal fields since 2003 with the open-pit method only.

Until 2018 Germany produced significant amounts of hard coal in underground mines. Since 2019, hard coal has been imported exclusively. NMVOC emissions are considered insignificant as the coal outgased along the import route. Particle emissions from loading and unloading as well as storage are reported under [2.L\(a\) - Handling of Bulk Products](#)“

## Methodology

### Lignite production



Activity data for lignite production have been taken from the *Statistik der Kohlenwirtschaft*'s website (in German only) <sup>1)</sup>. Extracted coal is moved directly to processing and to power stations.

The emission factors used for calculating emissions from lignite production (TSP, PM<sub>10</sub> and PM<sub>2.5</sub>) already include possible emissions from transport and storage. The PM<sub>10</sub> factor is taken from a 2021 research project <sup>2)</sup>. TSP and PM<sub>2.5</sub> have been reduced the same way on the basis of the previous values <sup>3)</sup>.

Table 1: Annual amounts of extracted raw lignite, in [Mt]

1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023
357	193	168	178	169	178	172	171	166	131	107	126	131	102

Table 2: Emission factors applied for lignite extraction, in [kg/t]

Pollutant	Value
TSP	0.03
PM <sub>10</sub>	0.015
PM <sub>2.5</sub>	0.002

### Lignite coke production

Table 3: Annual amounts of lignite coke produced, in [Mt]

1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023
3.36	0.19	0.18	0.17	0.18	0.17	0.16	0.15	0.16	0.16	0.15	0.16	0.14	0.16

Emissions from lignite production other than listed below are reported by plant operator. Particle emission factors were verified in a research project (Emissionen und Maßnahmenanalyse Feinstaub 2000-2020) by Jörß et al. (2007)<sup>4)</sup>.

Table 4: Emission factors applied for lignite-coke production

Pollutant	Unit	Value
TSP	kg/t	0.1
PM <sub>10</sub>	kg/t	0.048
PM <sub>2.5</sub>	kg/t	0.013
PAH	mg/t	55
PCDD/F	µg/t	0.03

### Hard coal coke production

The activity rates for hard coal coke production have been taken from the *Statistik der Kohlenwirtschafts's* website (in German only)<sup>5)</sup>.

Table 5: Annual amounts of hard coal coke produced, in [Mt]

1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023
18.5	11.1	9.12	8.40	8.15	8.80	9.39	9.34	9.36	8.91	7.94	8.25	8.27	8.03

The emission factors for hard coal coke production have been obtained from the research project "Emission factors for the iron and steel industry, for purposes of emissions reporting" ("Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung") by Hensmann et al. (2011)<sup>6)</sup>.

Table 6: Emission factors for hard coal coke production

Pollutant	Unit	Value
CO	kg/t	0.015
NH <sub>3</sub>	kg/t	0.000243
NMVOC	kg/t	0.096
SO <sub>2</sub>	kg/t	0.004
TSP	kg/t	0.011
PM <sub>10</sub>	kg/t	0.004
PM <sub>2.5</sub>	kg/t	0.004
PAH	mg/t	55
PCDD/F	µg/t	0.0015

There are many potential sources of PAH emissions from coking plants. The dominant emission sources are leakages from coke oven doors and from charging operations. As there is limited data available on PAH emissions, the uncertainties of the estimated emission factors are very high. It should also be taken into account that emissions from coke production greatly

vary between different coke production plants. The emission factors for benzo[a]pyrene and mixed PAH have been revised by research projects in 2010 <sup>7)8)</sup>. Split factors for Black Carbon (BC) are based on the EMEP Guidebook 2016 <sup>9)</sup>.

### Decommissioned hard-coal mines

NMVOC Emissions from decommissioned hard-coal mines play a role in this sub- source category. When a hard-coal mine is decommissioned, mine gas can escape from neighbouring rock, and from coal remaining in the mine, into the mine's network of shafts and passageways. Since the mine is no longer artificially ventilated, the mine gas collects and can then reach the surface via gas pathways in the overlying rock or via the mine's own shafts and passageways. Such mine gas was long seen primarily as a negative environmental factor. Recently, increasing attention has been given to the gas' positive characteristics as a fuel (due to its high methane content, it is used for energy recovery). In the past, use of mine gas was rarely cost-effective. This situation changed fundamentally in 2000 with the Renewable Energy Sources Act (EEG). Although mine gas is a fossil fuel in finite supply, its use supports climate protection, and thus the gas was included in the EEG. The Act requires network operators to accept, and provide specified compensation for, electricity generated with mine gas and fed into the grid.

The NMVOC emissions from decommissioned hard-coal mines have been calculated in the research project "Potential for release and utilisation of mine gas" ("Potential zur Freisetzung und Verwertung von Grubengas") <sup>10)</sup>. The relevant calculations were carried out for all mining-relevant deposits in Germany.

Table 7: NMVOC emission factor for decommissioned hard-coal mines, in [kg/m<sup>13</sup>]

EF
0.001599

## Recalculations



For more details please refer to the super-ordinate chapter [1.B - Fugitive Emissions from fossil fuels](#)

## Planned improvements



At the moment, no further category-specific improvements are planned.

<sup>1)</sup> Statistik der Kohlenwirtschaft (2019) [External Link](#) (last pageview: March 2025)

<sup>2)</sup> Dokumentation zur Berechnung des PM10-Austrags aus dem Tagebau Hambach im Jahr 2013 und Ableitung eines Emissionsfaktors (2021) [External Link](#) (last pageview: March 2025)

<sup>3)</sup> Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP) [External Link](#)

<sup>4)</sup> Jörß et al. (2007): Jörß, W.; Handke, V. (Institut für Zukunftsstudien und Technologiebewertung (IZT), Berlin) mit Beiträgen von Lambrecht, U., Dünnebeil, Fr. (Institut für Energie und Umweltforschung Heidelberg (ifeu)): Emissionen und Maßnahmenanalyse Feinstaub 2000–2020. UBA-TEXTE Nr. 38/200; URL: <https://www.umweltbundesamt.de/publikationen/emissionen-massnahmenanalyse-feinstaub-2000-2020>; on order of the Umweltbundesamt; Dessau-Roßlau, 2007.

<sup>6)</sup> Hensmann et al. (2011): Hensmann, M., Haardt, S., Ebert, D.: Emission factors for the iron and steel industry, for purposes of emissions reporting" ("Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung"; <https://www.umweltbundesamt.de/sites/default/files/medien/publikation/long/4362.pdf>; Dessau-Roßlau, October 2012

<sup>7)</sup> Federal Environment Agency and DFIU research project "Anpassung der deutschen Methodik zur rechnerischen Emissionsermittlung an internationale Richtlinien, Teilbericht Prioritäre Quellen", 2010 (not available online)

<sup>8)</sup> Federal Environment Agency and BFI research project No. 3707 42 301 "Emissionsfaktoren zur Eisen- und Stahlindustrie für die Emissionsberichterstattung", 2011 [External Link](#)

---

<sup>9)</sup> EMEP/EEA Air Pollutant Emission Inventory Guidebook 2016; published in 2016 [External Link](#) (last pageview: March 2025)

<sup>10)</sup> Meiners, H. (2014): Potential zur Freisetzung und Verwertung von Grubengas