5.B.1 - Biological Treatment of Waste: Composting

Short description

Within NFR category **5.B.1**, <u>ammonia (NH₃) emissions from composting</u> of organic wastes are reported.

| NFR Code | Method | AD | EF | | | | | |
|--------------------------|----------------------------|--------------------------------------|----------------------|--|--|--|--|--|
| 5.B.1 | CS | NS | CS | | | | | |
| Method(s) applied | | | | | | | | |
| D | Default | Default | | | | | | |
| T1 | Tier 1 / Simple Metho | Tier 1 / Simple Methodology * | | | | | | |
| T2 | Tier 2* | Tier 2* | | | | | | |
| Т3 | Tier 3 / Detailed Met | Tier 3 / Detailed Methodology * | | | | | | |
| С | CORINAIR | CORINAIR | | | | | | |
| CS | Country Specific | Country Specific | | | | | | |
| М | Model | Model | | | | | | |
| * as described in the EM | EP/EEA Emission Inventory | v Guidebook - 2019, ii | n category chapters. | | | | | |
| (source for) Activity D | ata | | | | | | | |
| NS | National Statistics | National Statistics | | | | | | |
| RS | Regional Statistics | Regional Statistics | | | | | | |
| IS | International Statistic | International Statistics | | | | | | |
| PS | Plant Specific | Plant Specific | | | | | | |
| As | Associations, busines | Associations, business organisations | | | | | | |
| Q | specific Questionnair | specific Questionnaires (or surveys) | | | | | | |
| М | Model / Modelled | Model / Modelled | | | | | | |
| С | Confidential | Confidential | | | | | | |
| (source for) Emission | Factors | | | | | | | |
| D | Default (EMEP Guide | Default (EMEP Guidebook) | | | | | | |
| CS | Country Specific | Country Specific | | | | | | |
| PS | Plant Specific | Plant Specific | | | | | | |
| М | Model / Modelled | Model / Modelled | | | | | | |
| С | Confidential | | | | | | | |

| NO _x | NMVOC | SO ₂ | NH ₃ | PM _{2.5} | PM ₁₀ | TSP | BC | CO | Heavy Metals | POPs |
|-----------------|--|-----------------|-----------------|-------------------|------------------|-----|----|----|--------------|------|
| NA | NA | NA | -/- | NA | NA | NA | NA | NA | NA | NA |
| L/- | L/- key source by Level only | | | | | | | | | |
| -/T | T key source by Trend only | | | | | | | | | |
| L/T | Key source by both Level and Trend | | | | | | | | | |
| -/- | no key source for this pollutant | | | | | | | | | |
| IE | emission of specific pollutant Included Elsewhere (i.e. in another category) | | | | | | | | | |
| NE | emission of specific pollutant N ot E stimated (yet) | | | | | | | | | |
| NA | specific pollutant not emitted from this source or activity = Not Applicable | | | | | | | | | |
| * | no analysis done | | | | | | | | | |

Separately collected organic waste (biowaste) from e.g. households, public garden and park service, food industry, restaurants, canteens and from agriculture can be treated in two different ways: aerobic treatment (composting) and anaerobic treatment (biogas production).

The aim of the treatment reported under this category is the production of compost, leading to the recycling of nutrients and organic matter.

The produced compost is used as fertilizer or soil improver in agriculture or horticulture and also in private gardening. In Germany about 50% of the organic waste is treated in composting plants and ammonia (NH_3) is an important emission to air.

Method

Emissions from composting are not a key source and of minor priority.

Activity Data

Official statistical data (Statistisches Bundesamt, GENESIS, Table Nr. 32111-0003 - Erhebung der Abfallentsorgung; ¹) are used for the estimation. The data are published on a yearly basis with an exception for the current year of reporting. The activity data for the current year of reporting are obtained, initially, by extrapolating the trend of the last 3 years into the future. In the following year, when the actual activity data for the given year becomes available, they replace the data that were extrapolated. This procedure has only a very small impact on the total emissions in the relevant current report year.

Emission factors

The emission factor used for calculating NH_3 emissions is based on emission data from a research project ²). The NH_3 -EF is 222 g/t and used for the whole time series. The use of abatement technologies (such as biofilters) are taken into account.

Uncertainties

The AD from Statistisches Bundesamt have an uncertainty of $\pm 2\%$ whereas the uncertainty for the EF is -59/+130% (ibid.).

Recalculations

When preparing the current inventory data, statistical data are only available for the previous reporting year. The current reporting year must therefore be extrapolated on the basis of the previous year. The result of this approach is revised by the correct data in the following year. For this reason, annual recalculations are required for the previous year.

Table 1: Revised biowaste activity data, in [kt]

| | 2022 |
|---------------------|-------|
| current submission | 8.174 |
| previous submission | 9.092 |

Table 2: Accordingly revised NH₃ emissions, in [t]

| | 2022 |
|---------------------|-------|
| current submission | 1.815 |
| previous submission | 2.018 |



For **pollutant-specific information on recalculated emission estimates for Base Year and for the current year, please see the recalculation tables following chapter 8.1 - Recalculations.

Planned improvements

Currently no improvements are planned.

¹⁾ Statistisches Bundesamt, GENESIS, Table Nr. 32111-0003 - Erhebung der Abfallentsorgung; Wiesbaden; https://www-genesis.destatis.de/genesis//online?operation=table&code=32111-0003&bypass=true&levelindex=0&levelid=1 630573034654#abreadcrumb

²⁾ Carsten Cuhls, Birte Mähl, Joachim Clemens; gewitra Ingenieurgesellschaft für Wissenstransfer mbH: Ermittlung der Emissionssituation bei der Verwertung von Bioabfällen;

https://www.umweltbundesamt.de/publikationen/ermittlung-der-emissionssituation-bei-der; im Auftrag des Umweltbundesamtes, April 2015