

Explanation of Key Trends - Sulfur Dioxide

Obligations

Under the terms of the UN ECE Geneva Convention on Long-range Transboundary Air Pollution Control (CLTRAP, 1979), the Federal Republic of Germany was obliged by the UN ECE Helsinki Protocol to reduce its annual sulphur emissions by at least 30% by 1993, as compared to 1980 levels. In 1993, the SO₂ emissions were 2.9 Mt, compared to approximately 7.5 Mt in 1980. This represents a reduction of 61%. The second UN ECE protocol on the reduction of sulphur emissions obliged Germany to reduce SO₂ emissions to 1,300 kt by 2000, and to 990 kt by 2005. The targets set for 2000 & 2005 have already been achieved by 1998.

More recently, Germany has made a commitment under the multicomponent protocol to further minimise SO₂ emissions. Since 2010, it is no longer permissible to exceed a National Emission Ceiling of 550 kt SO₂ for Germany as whole. The revised Gothenburg Protocol and the revised NEC Directive both define emission reduction targets relative to a 2005 base year, mandating 21 % (2020) and 58 % (2030) reductions respectively.

While Germany's compliance with these obligations is not discussed here, further information on this subject can be found in [Chapter 9 - Projections](#) and [Chapter 11 - Adjustments and Emission Ceiling Exceedance](#).

Main drivers

SO₂ **total emissions** show a falling trend from 1990 onwards, with **emission reductions of over 95% since 1990**.

By far the largest proportion of SO₂ is produced by the oxidation of the sulphur contained in the fuels used in combustion processes, **Fuel Combustion (NFR 1.A)** with a 96% share of total SO₂ emissions in 1990 and a 97% reduction between 1990 and 2023.

In 1990, the biggest source of emissions therein is **Public Electricity and Heat Production (NFR 1.A.1.a)** with about 47% of the emissions from **Fuel Combustion (NFR 1.A)**. Other sources substantially influencing the SO₂ emission trend are **Manufacturing Industries and Construction (NFR 1.A.2)** and **Other Sectors (NFR 1.A.4**, including commercial/institutional and residential sources), each adding about one fifth of 1990 emissions from Fuel Combustion (NFR 1.A).


All of these sub-categories show a reduction >95 per cent between 1990 and 2023, due to stricter regulations of West Germany that applied to the New German Länder after the German Reunification and changed the fuel mix from sulphur-rich solid fuels to liquid and gaseous fuels.

Table: SO₂ emissions 1990-2023, in kilotonnes [kt]

															Trend: latest compared to	
1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	1990	previous year
5 461	1 743	643	468	397	341	316	307	294	258	239	249	247	210	199	-96.4%	-5.3%

Sulfur Dioxide / Schwefeldioxid


Emissions per Sector / Sektorale Emissionen



Quelle: German Emission Inventory (04.02.2026)

Sulfur Dioxide / Schwefeldioxid

Emissions per Sector / Sektorale Emissionen



Quelle: German Emission Inventory (04.02.2026)