





occurred. In 2024, dairy cattle numbers are 56.5 % of 1990 numbers, while the total population of other cattle is at 52.3 % of 1990. Swine numbers decreased until 1995 and then increased slightly. Since 2014 a new decrease occurred which became significant between 2020 and 2022 (total swine numbers were reduced by around 18 % within two years). 2024 swine numbers are 66.6 % of 1990 numbers. The 2024 numbers of horses, sheep and goats are, respectively, at 99.6 %, 55.9 % and 183.5 % of 1990.

Figures for broilers and turkeys are showing a massive increase since 1990. Since the year 2013, there have been only minor changes of total poultry numbers. In total, 2024 poultry population figures are at 147.2 % of 1990.

Emissions of deer, rabbits, ostrich and fur-bearing animals are reported since submission 2024. The underlying animal numbers of these categories were estimated in different ways because there are no surveys which collect those animal numbers. However, the impact of those animal categories on the total emissions is small.

A detailed description of the animal numbers used can be found in Vos et al. (2026), chapter 2.3.

Table 1: Population of animals, in [1,000 individuals]

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>dairy cattle</b>	6,354.6	5,229.4	4,569.8	4,236.4	4,183.1	4,284.6	4,217.7	4,199.0	4,100.9	4,011.7	3,921.4	3,832.7	3,809.7	3,712.8	3,589.4
<b>other cattle</b>	13,133.4	10,660.5	9,968.9	8,800.4	8,628.7	8,350.8	8,248.9	8,082.2	7,848.2	7,627.9	7,380.5	7,206.9	7,187.2	7,123.4	6,871.9
<b>buffalo</b>	NO	NO	IE												
<b>mules and asses</b>	IE														
<b>horses</b>	1,373.5	1,743.9	1,373.7	1,398.1	1,269.9	1,233.1	1,215.4	1,223.4	1,231.5	1,239.5	1,247.6	1,277.8	1,308.0	1,338.2	1,368.3
<b>sheep</b>	3,266.1	2,990.7	2,743.3	2,643.1	2,245.0	1,866.9	1,851.0	1,863.2	1,846.0	1,813.6	1,780.3	1,794.8	1,805.7	1,847.6	1,824.5
<b>goats</b>	90.0	100.0	140.0	170.0	149.9	135.9	138.8	142.8	146.9	150.9	154.9	157.5	160.0	162.6	165.2
<b>swine</b>	26,502.5	20,387.3	21,767.7	22,742.8	22,244.4	22,978.5	22,761.2	22,920.8	22,019.2	21,625.8	21,622.0	19,728.6	17,692.3	17,525.3	17,642.7
<b>laying hens</b>	53,450.5	47,575.8	48,640.0	43,641.6	41,700.5	50,619.3	51,935.5	52,571.1	53,206.6	53,842.1	54,477.6	54,921.5	55,365.4	55,809.3	56,084.2
<b>broilers</b>	35,393.0	46,625.9	61,940.7	76,045.0	98,389.7	94,909.4	93,791.3	93,458.7	93,126.1	92,793.5	92,461.0	91,004.5	89,548.1	88,091.6	88,091.6
<b>turkeys</b>	5,029.2	6,742.0	8,893.1	10,611.1	11,344.0	12,658.5	12,359.9	12,164.7	11,969.5	11,774.3	11,579.1	10,718.9	9,858.6	8,998.3	8,998.3
<b>pullets</b>	17,210.8	16,149.2	17,284.1	16,050.9	14,827.0	13,828.3	12,921.8	12,736.3	12,550.7	12,365.1	12,179.6	12,253.0	12,326.5	12,399.9	12,399.9
<b>ducks</b>	2,013.7	1,933.7	2,055.7	2,352.2	3,164.3	2,410.8	2,236.4	2,209.1	2,181.9	2,154.6	2,127.4	1,949.3	1,771.2	1,593.1	1,593.1
<b>geese</b>	781.5	617.0	404.8	329.5	278.1	400.8	329.0	327.7	326.3	324.9	323.5	354.2	385.0	415.7	415.7
<b>deer</b>	155.8	204.0	252.3	261.5	270.0	277.4	278.9	280.4	281.9	283.3	284.8	286.3	287.8	289.3	290.8
<b>rabbits</b>	1,851.4	1,565.6	1,268.9	997.0	864.2	720.7	691.2	642.7	608.3	593.9	548.4	470.0	430.6	422.8	422.5
<b>ostrich</b>	NO	1.2	2.5	3.7	4.9	7.7	7.4	7.4	7.9	7.4	7.9	6.1	5.1	4.6	4.0
<b>fur animals</b>	179.9	179.9	179.9	153.5	121.7	34.4	24.7	15.0	5.3	5.3	NO	NO	NO	NO	NO

## Additional data

Emission calculations in accordance with a Tier 2 or Tier 3 method require data on animal performance (animal weight, weight gain, milk yield, milk protein content, milk fat content, numbers of births, numbers of eggs and weights of eggs) and on the relevant feeding details (phase feeding, feed components, protein and energy content, digestibility and feed efficiency). To subdivide officially recorded total numbers of turkeys into roosters and hens, the respective population percentages need to be known. Details on data requirements for the modelling of emissions from livestock husbandry in the German inventory can be found in Vos et al. (2026), Chapter 2.

Most of the data regarding feed and performance is not available from official statistics and was obtained from literature, from publications by agricultural associations, from guidelines for agricultural consulting in Germany and from expert judgments.

For 1991, 1995 and 1999, frequency distributions of feeding strategies, husbandry systems (shares of pasturing/stabling; shares of various housing methods), storage types as well as techniques of farm manure spreading were obtained with the help of the RAUMIS agricultural sector model (Regionalisiertes Agrar- und UmweltInformationssystem für Deutschland/ Regionalised agricultural and environmental information system for Germany). RAUMIS has been developed and is operated by the Institute of Rural Studies of the Thünen Institute (Federal Research Institute for Rural Areas, Forestry and Fisheries). For an introduction to RAUMIS see Weingarten (1995)<sup>2)</sup>; a detailed description is provided in Henrichsmeyer et al. (1996)<sup>3)</sup>.

RAUMIS did not model complete time series but only selected years. RAUMIS data for the years 1991, 1995, and 1999 are used in the inventory for the respective years. For 1990, the data for 1991 is adopted, for the intervening years (1992-1994 and 1996-1998) data gaps were closed by linear interpolation on district level.

For the year 2009, respective data are used that were derived from the 2010 official agricultural census and the simultaneous survey of agricultural production methods (Landwirtschaftliche Zählung 2010, Statistisches Bundesamt/ Federal Statistical Office) as well as the 2011 survey on manure application practices (Erhebung über Wirtschaftsdüngerausbringung, Statistisches Bundesamt/ Federal Statistical Office). The gaps between the latest RAUMIS model data (1999) and the first official data (2009) were closed by linear interpolation on district level. For the year 2015, data on techniques of farm manure spreading from the 2016 official agricultural census (Agrarstrukturenerhebung 2016, Statistisches Bundesamt / Federal Statistical Office) are used. For the year 2019 data from the 2020 official agricultural census (Landwirtschaftszählung 2020, LW20) are used for housing systems, storage systems and manure spreading systems.

For 2010 to 2018 the housing and storage systems data was linearly interpolated between the censuses of 2010 and 2020. The data on manure spreading techniques was linearly interpolated between the census data from 2009 and 2015, and for 2016 to 2018 between the censuses conducted in 2016 and 2020. In addition, it was taken into account that, as of 2012, slurry spread on bare soil has to be incorporated within four hours. The data from the 2020 official agricultural census (DESTATIS, 2020)<sup>4</sup> is being used for subsequent years until more current data is available.

For a description of the RAUMIS data, the data from official surveys and additional data from other sources see Vos et al. (2026), Chapter 2.5. Time series of frequency distributions of housing systems, storage systems and application techniques as well as the corresponding emission factors are provided in NID 2026<sup>5</sup>, Chapter 17.3.1.

Since submission 2026 transports of manure, energy crops and digestates between NUTS 3 regions are simulated in the calculation model. This does not have major influence on the emissions of the whole country, but changes the distribution of emissions between single NUTS 3 regions. For details on the methodology, see Vos et al. (2026), Chapter 1.2.

## NH<sub>3</sub> and NO<sub>x</sub>

### Method

#### N in manure management

##### N excretion

In order to determine NH<sub>3</sub> and NO<sub>x</sub> emissions from manure management of a specific animal category, the individual N excretion rate must be known as well as, for NH<sub>3</sub>, the TAN content of the N excretions. Default excretion rates are provided by IPCC Guidelines and default TAN contents can be found in the EMEP Guidebook, 2023<sup>6</sup>. However, the German agricultural emission inventory uses N mass balances to calculate the N excretions and the TAN contents of almost all reported animal categories. N mass balance calculations (see below) consider N intake with feed, N retention due to growth, N contained in milk and eggs, and N in offspring. Table 2 presents national means of N excretions and TAN contents. For methodological details and mass balance input data see Vos et al. (2026), Chapter 4.2 as well as Chapter 4.1.2. For dairy cows N excretion is estimated based on milk urea content since submission 2026, for details see Vos et al. (2026) Chapter 4.1.2. This allows an estimation that is based on the actual feeding of the dairy cows, thus it is possible to depict the effect of feeding changes in N excretions. Milk urea contents are obtained from the agricultural information services company vit (Vereinigtes Informationssysteme Tierhaltung w.V.) and directly by four state control associations, who operate a separate database. Each record includes farm-aggregated information regarding herd size, milk yield, milk urea, fat and protein contents, milking day, and racial composition of the herd. For the time between 2005 and 2022 over 10 million milk urea contents at herd-level were obtained. For the purpose of the inventory these were aggregated on district level, weighted by herd size. For the years from 1990-2004 the values of 2005 were taken, for 2023 and 2024 the values of 2022.

Table 2: National means of N excretions and TAN contents


	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>mean N excretions in kg per animal place</b>															
<b>dairy cattle</b>	106.3	110.8	114.3	117.5	114.3	115.2	116.8	118.7	120.3	121.4	119.7	117.0	116.1	118.6	120.2
<b>other cattle</b>	37.9	39.9	41.3	41.2	42.1	42.5	42.5	42.7	42.9	43.4	43.7	43.9	43.9	44.2	44.4
<b>horses</b>	48.2	48.1	49.0	48.8	48.8	54.0	55.0	56.1	57.1	58.2	59.2	59.2	59.2	59.2	59.2
<b>sheep</b>	7.7	7.7	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8
<b>goats</b>	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0
<b>swine</b>	12.8	13.1	13.0	12.8	12.6	12.4	12.4	12.3	12.2	12.1	12.1	12.3	12.2	12.2	12.3

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>laying hens</b>	0.81	0.78	0.76	0.79	0.86	0.88	0.89	0.89	0.89	0.89	0.90	0.90	0.90	0.90	0.91
<b>broilers</b>	0.48	0.37	0.37	0.36	0.35	0.40	0.40	0.40	0.41	0.40	0.39	0.39	0.40	0.41	0.42
<b>turkeys</b>	2.0	2.0	2.0	2.2	2.2	2.3	2.3	2.3	2.2	2.2	2.1	2.1	2.1	2.1	2.1
<b>pullets</b>	0.38	0.35	0.32	0.33	0.33	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34
<b>ducks</b>	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61	0.61
<b>geese</b>	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70
<b>deer</b>	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00
<b>rabbits</b>	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80
<b>ostrich</b>	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60	15.60
<b>fur animals</b>	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59	4.59
<b>mean TAN contents in %</b>															
<b>dairy cattle</b>	51.3	48.9	47.3	45.9	44.6	43.7	43.3	43.4	42.9	42.3	41.5	41.1	41.4	40.5	39.9
<b>other cattle</b>	65.5	65.7	65.7	65.7	66.0	66.3	66.4	66.4	66.4	66.4	66.4	66.3	66.3	66.3	66.3
<b>horses</b>	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0
<b>sheep</b>	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0
<b>goats</b>	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0
<b>swine</b>	72.0	71.7	71.1	71.8	72.3	71.5	71.3	71.2	71.0	70.9	70.7	70.8	70.6	70.5	70.4
<b>laying hens</b>	70.2	69.6	69.0	69.3	70.0	70.2	70.1	70.1	70.2	70.2	70.1	70.1	70.2	70.3	70.2
<b>broilers</b>	60.8	58.9	56.4	53.5	50.0	46.9	46.5	46.1	45.7	45.2	44.8	44.8	44.8	44.8	44.8
<b>turkeys</b>	64.7	64.7	63.0	63.9	63.0	63.5	63.5	63.5	63.0	63.0	62.1	62.1	62.1	62.1	62.1
<b>pullets</b>	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4	69.4
<b>ducks</b>	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9	49.9
<b>geese</b>	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0
<b>deer</b>	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0
<b>rabbits</b>	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0
<b>ostrich</b>	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0	70.0
<b>fur animals</b>	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0	60.0

#### N mass flow and emission assessment

The calculation of the emissions of NH<sub>3</sub>, N<sub>2</sub>O, NO<sub>x</sub> and N<sub>2</sub> from German animal husbandry is based on the so-called N mass flow approach (e. g. Dämmgen and Hutchings, 2008<sup>7)</sup>). This approach differentiates between N excreted with faeces (organic nitrogen Norg, i. e. undigested feed N) and urine (total ammoniacal nitrogen TAN, i. e. fraction of feed N metabolized). The N flow within the manure management system is treated as depicted in the figure below. This method reconciles the requirements of both the Atmospheric Emission Inventory Guidebook for NH<sub>3</sub> emissions (EMEP, 2023), and the IPCC guidelines for greenhouse gas emissions (IPCC (2006)<sup>8)</sup>. Reidy et al. (2008),<sup>9)</sup> showed for several European countries (Germany, the Netherlands, Switzerland, United Kingdom) that their N-flow based inventory models yielded, in spite of national peculiarities, comparable results as long as standardised data sets for the input variables were used.

Not explicitly shown in the N mass flow scheme is air scrubbing in housing and anaerobic digestion of manure. These issues are separately described further below. Note that emissions from grazing and application are reported in sector 3.D.



General scheme of N flows in animal husbandry

$m$ : mass from which emissions may occur. Narrow broken arrows: TAN (total ammoniacal nitrogen); narrow continuous arrows: organic N. The horizontal arrows denote the process of immobilisation in systems with bedding occurring in the house, and the process of mineralisation during storage, which occurs in any case. Broad arrows denote N-emissions assigned to manure management ( $E_{yard}$  NH<sub>3</sub> emissions from yards;  $E_{house}$  NH<sub>3</sub> emissions from house;  $E_{storage}$  NH<sub>3</sub>, N<sub>2</sub>O, NO<sub>x</sub> and N<sub>2</sub> emissions from storage;  $E_{applic}$  NH<sub>3</sub> emissions during and after spreading;  $E_{graz}$  NH<sub>3</sub>, N<sub>2</sub>O, NO<sub>x</sub> and N<sub>2</sub> emissions during and after grazing;  $E_{soil}$  N<sub>2</sub>O, NO<sub>x</sub> and N<sub>2</sub> emissions from soil resulting from manure input).

The model allows tracing of the pathways of the two N fractions after excretion. The various locations where excretion may take place are considered. The partial mass flows through the livestock systems are represented. During storage N<sub>org</sub> can be transformed into TAN and vice versa. Both, the way and the magnitude of such transformations may be influenced by manure treatment processes like, e. g., anaerobic digestion where a considerable fraction of N<sub>org</sub> is mineralized to TAN. For details see Vos et al. (2026), Chapter 4.2. Wherever NH<sub>3</sub> is emitted, its formation is related to the amount of the TAN present. N<sub>2</sub>O emissions are related to the total amount of N available (N<sub>org</sub> + TAN). NO<sub>x</sub> emissions (i. e. NO emissions) are calculated proportionally to the N<sub>2</sub>O emissions, see section 'Emission factors'. Note that the N<sub>2</sub>O, NO<sub>x</sub> and N<sub>2</sub> emissions from the various storage systems include the respective emissions from the related housing systems.

### Air scrubber systems in swine and poultry housings

For pig and poultry production the inventory model considers the effect of air scrubbing. Data on frequencies of air scrubbing facilities and the removal efficiency are provided by KTBL (Kuratorium für Technik und Bauwesen in der Landwirtschaft / Association for Technology and Structures in Agriculture) supplemented by data from the 2020 agricultural census. The average removal efficiency of NH<sub>3</sub> is 80 % for swine and 70 % for poultry, while for TSP and PM<sub>10</sub> the rates are set to 90 % and for PM<sub>2.5</sub> to 70 % for both animal categories. For swine two types of air scrubbers are distinguished: systems of “first class” that remove both NH<sub>3</sub> and particles, and “second class” systems that remove only particles reliably and have a NH<sub>3</sub> removal efficiency of 20%.

According to KTBL, 7.6 % of all pig places were equipped with “first class” systems in 2024, another 12.6 % were equipped with “second class” systems. For poultry 0.9 % of all laying hen places and 2.5 % of all broiler places were equipped with air scrubbers that remove both NH<sub>3</sub> and particles. The amounts of NH<sub>3</sub>-N removed by air scrubbing are completely added to the pools of total N and TAN for landspreading. For details see Vos et al. (2026), Chapter 4.2.2.

### Anaerobic digestion of manure



	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>fur animals</b>	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123	1.123
<b>Nitrogen oxides</b>															
<b>dairy cattle</b>	0.163	0.154	0.163	0.168	0.158	0.146	0.146	0.147	0.148	0.148	0.143	0.137	0.136	0.140	0.142
<b>other cattle</b>	0.064	0.068	0.071	0.076	0.079	0.077	0.077	0.077	0.078	0.078	0.078	0.077	0.077	0.078	0.078
<b>horses</b>	0.169	0.169	0.172	0.171	0.171	0.184	0.187	0.190	0.193	0.196	0.198	0.198	0.198	0.198	0.198
<b>sheep</b>	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012	0.012
<b>goats</b>	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
<b>swine</b>	0.013	0.015	0.015	0.016	0.016	0.014	0.014	0.013	0.013	0.013	0.012	0.012	0.012	0.012	0.012
<b>laying hens</b>	0.00027	0.00026	0.00026	0.00030	0.00035	0.00037	0.00036	0.00036	0.00023	0.00023	0.00024	0.00024	0.00023	0.00024	0.00024
<b>broilers</b>	0.00016	0.00012	0.00012	0.00013	0.00014	0.00016	0.00016	0.00016	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011	0.00011
<b>turkeys</b>	0.00068	0.00068	0.00070	0.00085	0.00095	0.00091	0.00094	0.00094	0.00069	0.00066	0.00061	0.00061	0.00060	0.00058	0.00060
<b>pullets</b>	0.00013	0.00012	0.00011	0.00013	0.00015	0.00014	0.00014	0.00014	0.00009	0.00009	0.00009	0.00009	0.00009	0.00010	0.00010
<b>ducks</b>	0.00024	0.00024	0.00024	0.00025	0.00026	0.00027	0.00028	0.00027	0.00021	0.00021	0.00022	0.00022	0.00021	0.00021	0.00021
<b>geese</b>	0.00025	0.00025	0.00025	0.00029	0.00032	0.00032	0.00030	0.00030	0.00023	0.00023	0.00024	0.00024	0.00023	0.00022	0.00022
<b>deer</b>	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
<b>rabbits</b>	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269	0.00269
<b>ostrich</b>	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126	0.00126
<b>fur animals</b>	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508	0.01508

**Trend discussion for Key Sources**

Dairy cattle, other cattle and swine are key sources of NH<sub>3</sub> emissions from manure management. The time series of the total NH<sub>3</sub> emissions from all three categories are predominantly driven by the development of the animal numbers, see Table 1. However, the effect of decreasing animal numbers is partly compensated by the continuously increasing animal performance. This leads to increasing N excretions per animal, see Table 2, which, in principle, is reflected by increasing implied emission factors, see Table 3. For swine the IEF is decreasing over time due to lower raw protein contents in feed and the use of air scrubbing systems that, to a high degree, remove NH<sub>3</sub> from the housings.

For NO<sub>x</sub> there are no key categories.

**Recalculations**

All timeseries of the emission inventory have completely been recalculated. Tables 4 and 5 compare the recalculated time series for NH<sub>3</sub> and NO<sub>x</sub> from 3B with the respective data of last year’s submission.

For NH<sub>3</sub> there are many reasons for very different emissions compared to last year’s submission. For dairy cows the new methodology to calculate N and TAN excretions (see [main page of the agricultural sector recalculation No. 3](#)) results in general in lower emissions. The upward correction of historic milk yields (**recalculation No. 4**) results in higher emissions. In combination these two recalculations result in lower emissions, especially in more recent years. The subdivision of the sows category in gilts and old sows (**recalculation No. 5**) is the main reason for lower emissions from swine. The adjusted N excretion for horses after 2010 (**recalculation No. 6**) is the main reason for higher emissions from other animals. The addition of substrate transports to biogas plants (**recalculation No. 1**) has a smaller impact on emissions than the other recalculations. This recalculation is the main reason for the changes for emissions from other cattle and poultry but it also affects dairy cattle and swine emissions. Many of the other recalculations have much smaller effects. Overall, the changes result in lower emissions compared with last year’s submission.

The total emissions of NO<sub>x</sub> for all years up to 2019 are higher and thereafter a little bit lower than those of submission 2025. The main reasons for this are the recalculations done for dairy cattle (**recalculations No. 3 and No. 4**). Up to the year 2019 the effect of the adjusted milk yields (resulting in higher emissions) is higher than the effect of the new methodology to calculate N emissions which leads to lower N excretions especially in more recent years. All other reasons listed above regarding NH<sub>3</sub> have similar effects on NO<sub>x</sub> emissions. Further details on recalculations are described in Vos et al. (2026), Chapter 1.3.

Table 4: Comparison of NH<sub>3</sub> emissions [kt] with previous submission

<b>NFR Total emissions</b>
----------------------------

	Submission	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>Total</b>	<b>current</b>	297.22	250.13	245.22	244.46	235.19	231.40	227.54	226.84	220.30	215.27	209.68	199.90	190.21	188.78	187.67
	<b>previous</b>	300.67	252.12	248.41	248.80	242.06	237.98	233.87	231.70	224.95	220.40	216.51	208.17	197.88	196.54	
	<b>absolute change</b>	-3.45	-1.99	-3.20	-4.33	-6.87	-6.58	-6.33	-4.86	-4.65	-5.14	-6.83	-8.27	-7.67	-7.76	
	<b>relative change [%]</b>	-1.15	-0.79	-1.29	-1.74	-2.84	-2.76	-2.71	-2.10	-2.07	-2.33	-3.15	-3.97	-3.88	-3.95	
<b>Dairy cattle</b>	<b>current</b>	58.81	50.52	46.08	45.56	44.83	46.63	46.48	47.61	46.96	46.11	43.41	40.83	40.65	39.50	38.05
	<b>previous</b>	60.08	50.88	47.58	47.96	49.01	51.16	50.96	50.81	50.34	50.41	49.61	48.53	48.11	47.56	
<b>Other cattle</b>	<b>current</b>	75.09	61.37	58.84	54.44	57.03	52.31	51.04	49.59	47.76	46.24	44.92	43.89	43.75	43.65	42.16
	<b>previous</b>	75.09	61.37	58.84	54.43	57.07	52.45	51.18	49.73	47.86	46.37	45.09	44.10	43.95	43.75	
<b>Swine</b>	<b>current</b>	118.02	88.90	92.30	94.41	87.70	84.59	82.66	82.07	78.05	75.42	74.52	68.60	59.81	59.50	60.67
	<b>previous</b>	120.20	90.53	93.99	96.33	90.18	87.72	85.80	85.33	81.27	78.46	77.66	71.70	62.66	62.12	
<b>poultry</b>	<b>current</b>	23.31	22.67	26.15	28.06	25.82	27.64	27.13	26.92	26.51	26.09	25.03	24.31	23.25	22.86	23.06
	<b>previous</b>	23.31	22.67	26.16	28.07	26.00	27.82	27.37	27.17	26.72	26.32	25.23	24.50	23.43	22.93	
<b>Other animals</b>	<b>current</b>	22.00	26.67	21.85	22.00	19.81	20.24	20.23	20.64	21.02	21.42	21.80	22.27	22.75	23.27	23.74
	<b>previous</b>	22.00	26.67	21.85	22.00	19.81	18.82	18.56	18.66	18.75	18.84	18.92	19.32	19.73	20.18	

Table 5: Comparison of NO<sub>x</sub> emissions [kt] with previous submission

		<b>NFR Total emissions</b>														
	Submission	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>Total</b>	<b>current</b>	2.537	2.211	2.087	2.062	1.997	1.891	1.867	1.854	1.800	1.762	1.710	1.633	1.597	1.607	2.537
	<b>previous</b>	2.403	2.119	2.023	2.014	1.980	1.878	1.848	1.822	1.783	1.750	1.716	1.658	1.619	1.621	
	<b>absolute change</b>	0.13	0.09	0.06	0.05	0.02	0.01	0.02	0.03	0.02	0.01	-0.01	-0.02	-0.02	-0.01	
	<b>relative change [%]</b>	5.60	4.35	3.20	2.38	0.84	0.67	1.01	1.77	0.97	0.65	-0.40	-1.48	-1.39	-0.86	
<b>Dairy cattle</b>	<b>current</b>	1.036	0.807	0.743	0.712	0.660	0.623	0.618	0.618	0.608	0.593	0.562	0.523	0.519	0.520	0.508
	<b>previous</b>	0.892	0.708	0.672	0.658	0.634	0.607	0.600	0.590	0.585	0.580	0.570	0.549	0.544	0.545	
<b>Other cattle</b>	<b>current</b>	0.843	0.726	0.711	0.668	0.683	0.647	0.637	0.625	0.609	0.595	0.576	0.557	0.553	0.556	0.537
	<b>previous</b>	0.843	0.726	0.711	0.668	0.684	0.652	0.642	0.630	0.613	0.600	0.582	0.565	0.561	0.558	
<b>Swine</b>	<b>current</b>	0.351	0.313	0.326	0.366	0.358	0.316	0.308	0.303	0.286	0.271	0.266	0.241	0.208	0.209	0.213
	<b>previous</b>	0.360	0.319	0.332	0.372	0.367	0.332	0.322	0.318	0.300	0.286	0.281	0.258	0.224	0.222	
<b>poultry</b>	<b>current</b>	0.026	0.025	0.029	0.035	0.043	0.048	0.048	0.048	0.033	0.032	0.032	0.032	0.030	0.030	0.030
	<b>previous</b>	0.026	0.025	0.029	0.034	0.042	0.048	0.047	0.047	0.046	0.044	0.042	0.040	0.040	0.039	
<b>Other animals</b>	<b>current</b>	0.282	0.340	0.279	0.281	0.253	0.256	0.256	0.261	0.265	0.270	0.275	0.280	0.286	0.293	1.249
	<b>previous</b>	0.282	0.340	0.279	0.281	0.253	0.240	0.236	0.238	0.239	0.240	0.241	0.246	0.251	0.257	

**Planned improvements**

No improvements are planned at present.

**NM VOC**

In 2024, NMVOC emissions from manure management amount to 291.7 kt which is 97.1 % of total NMVOC emissions from the agricultural sector. 84.1 % thereof originate from cattle, 15.9 % from other animals.

**Method**

The Tier 2 methodology provided by EMEP (2023)-3B-26 was used to assess the NMVOC emissions from manure management for dairy cattle and other cattle. For all other animals the Tier 1 methodology (EMEP (2023)-3B-17) was used. The use of the Tier 2 methodology yields NMVOC emissions which formally could be reported in the sectors 3.D.a.2.a (application of manure to soils) and 3.D.a.3 (grazing emissions). However, to be congruent with the NMVOC emissions for other animal categories, Germany reports these emissions in the NMVOC emissions reported from manure management (3.B). For the NFR codes 3.D.a.2.a and 3.D.a.3 the key note IE is used for NMVOC emissions.



	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>fur animals</b>	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941	1.941

### Trend discussion for Key Sources

Dairy cattle and other cattle are key sources of NMVOC emissions from manure management. The total NMVOC emissions from both animal categories strongly correlate with the animal numbers given in Table 1 (dairy cattle:  $R^2 = 0.90$ ; other cattle:  $R^2 = 0.99$ ).

### Recalculations

All timeseries of the emission inventory have completely been recalculated. Table 7 compares the recalculated time series of the NMVOC emissions from 3.B with the respective data of last year's submission. The recalculated total emissions are higher. For dairy cattle and other cattle emissions are higher due to changes of  $NH_3$  emissions which have impact on the Tier 2 methodology which is applied for cattle NMVOC emissions. For other animals there are no changes compared with the previous submission. Further details on recalculations are described in Vos et al. (2026), Chapter 1.3.

Table 7: Comparison of NMVOC emissions [kt] with previous submission

		<b>NFR Total emissions</b>														
	<b>Submission</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2010</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>	<b>2022</b>	<b>2023</b>	<b>2024</b>
<b>Total</b>	<b>current</b>	418.62	362.01	346.77	325.36	324.20	325.33	322.01	318.62	314.16	311.13	306.95	300.63	296.10	296.66	291.74
	<b>previous</b>	415.22	357.45	342.28	320.98	320.35	322.59	318.93	315.55	311.20	307.95	303.73	297.43	293.26	292.11	
	<b>absolute change</b>	3.39	4.56	4.49	4.38	3.85	2.74	3.07	3.07	2.96	3.18	3.22	3.20	2.84	4.55	
	<b>relative change [%]</b>	0.82	1.27	1.31	1.36	1.20	0.85	0.96	0.97	0.95	1.03	1.06	1.07	0.97	1.56	
<b>Dairy cattle</b>	<b>current</b>	209.16	185.28	175.89	168.29	168.54	175.23	174.31	173.08	172.19	171.79	170.21	166.94	164.48	166.02	163.60
	<b>previous</b>	205.77	180.73	171.39	163.89	164.40	171.96	170.78	169.61	168.80	168.25	166.61	163.33	161.27	161.23	
<b>Other cattle</b>	<b>current</b>	162.85	131.10	123.68	107.78	105.33	98.83	97.03	94.80	91.85	89.49	87.02	85.52	85.15	84.63	81.81
	<b>previous</b>	162.85	131.10	123.68	107.80	105.62	99.36	97.49	95.21	92.27	89.86	87.40	85.93	85.52	84.87	
<b>Other animals</b>	<b>current</b>	46.61	45.63	47.21	49.29	50.33	51.27	50.67	50.73	50.12	49.84	49.72	48.17	46.47	46.01	46.33
	<b>previous</b>	46.61	45.63	47.21	49.29	50.33	51.27	50.67	50.73	50.12	49.83	49.72	48.17	46.47	46.01	

### Planned improvements

No improvements are planned at present.

## Particle emissions

In 2024, TSP emissions from manure management amount to 60.4 % of total emissions from the agricultural sector. Of these emissions 23.9 % originate from cattle, 32.9 % from pigs, and 41.6 % from poultry.

34.7 % of total  $PM_{10}$  emissions from the agricultural sector are caused by manure management, where 33.3 % originate from cattle, 15.3 % from pigs, and 47.0 % from poultry.

66.4 % of total  $PM_{2.5}$  emissions from the agricultural sector are caused by manure management, where 76.5 % originate from cattle, 2.4 % from pigs, and 16.7 % from poultry.

### Method

EMEP (2013-3B-26) provided a Tier 2 methodology. In the 2023 Guidebook (EMEP, 2023), this methodology has been replaced by a Tier 1 methodology. However, EF for cattle derived with the EMEP 2013 Tier 2 methodology remained unchanged. Therefore, the EMEP 2013<sup>11)</sup> methodology was kept for cattle. For swine the EMEP 2013 methodology was formally kept but the EMEP 2023 Tier 1 EF was used both for slurry and solid based manure management systems. In case the EMEP 2023 EFs are simply rounded EMEP 2013 EFs, the unrounded EMEP 2013 EFs were kept. For rabbits the EFs from



	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
<b>pullets</b>	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200
<b>ducks</b>	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400	0.1400
<b>geese</b>	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400	0.2400
<b>deer</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
<b>rabbits</b>	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107	0.0107
<b>ostrich</b>	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112
<b>fur animals</b>	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080
<b>PM<sub>2.5</sub></b>															
<b>dairy cattle</b>	0.3616	0.4181	0.4338	0.4393	0.4465	0.4582	0.4610	0.4636	0.4663	0.4690	0.4690	0.4690	0.4690	0.4690	0.4693
<b>other cattle</b>	0.1574	0.1548	0.1520	0.1487	0.1456	0.1444	0.1443	0.1442	0.1442	0.1440	0.1439	0.1440	0.1442	0.1439	0.1437
<b>horses</b>	0.1027	0.1026	0.1039	0.1038	0.1036	0.1037	0.1037	0.1037	0.1037	0.1037	0.1038	0.1037	0.1037	0.1037	0.1037
<b>sheep</b>	0.0059	0.0059	0.0060	0.0060	0.0060	0.0059	0.0059	0.0059	0.0059	0.0059	0.0059	0.0059	0.0059	0.0059	0.0059
<b>goats</b>	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112	0.0112
<b>swine</b>	0.0056	0.0057	0.0056	0.0055	0.0053	0.0050	0.0050	0.0049	0.0049	0.0049	0.0048	0.0048	0.0047	0.0047	0.0047
<b>laying hens</b>	0.0030	0.0030	0.0030	0.0030	0.0030	0.0029	0.0029	0.0029	0.0029	0.0029	0.0029	0.0029	0.0029	0.0029	0.0029
<b>broilers</b>	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020
<b>turkeys</b>	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200	0.0200
<b>pullets</b>	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020
<b>ducks</b>	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180	0.0180
<b>geese</b>	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320	0.0320
<b>deer</b>	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
<b>rabbits</b>	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021	0.0021
<b>ostrich</b>	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034	0.0034
<b>fur animals</b>	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040

### Trend discussion for Key Sources

Swine and laying hens are key sources of TSP emissions from manure management. The total TSP emissions from swine mainly follow the animal numbers given in Table 1 for the earlier years of the time series. However, due to increases in places equipped with air scrubbing and different emission factors of the different housing systems of the five swine subcategories (sows (divided in gilts and old sows), weaners, fattening pigs, boars) and the varying population shares in those housing systems the R<sup>2</sup> of the linear regression is lower than 1 (0.79). For laying hens (R<sup>2</sup> = 0.98) and broilers (R<sup>2</sup> = 0.99), due to the low prevalence of air scrubbing systems TSP emissions almost perfectly correlate with the animal numbers provided in Table 1.

### Recalculations

The following table 9 shows the effects of recalculations on emissions of particulate matter. Minimal differences compared with the previous submission are due to the correction of the number of animal places equipped with air scrubbers (**recalculation No. 11**), see [main page of the agricultural sector](#). Further details on recalculations are described in Vos et al. (2026), Chapter 1.3.

Table 9: Comparison of particle emissions (TSP, PM<sub>10</sub> & PM<sub>2.5</sub>) [kt] with previous submission

<b>TSP, PM<sub>10</sub>, PM<sub>2.5</sub> emissions from manure management, in kt</b>																
	<b>Submission</b>	<b>1990</b>	<b>1995</b>	<b>2000</b>	<b>2005</b>	<b>2010</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>	<b>2022</b>	<b>2023</b>	<b>2024</b>
<b>TSP</b>	<b>current</b>	50.36	43.31	44.13	43.37	42.70	43.91	43.56	43.53	42.51	41.85	41.67	39.64	37.71	37.32	37.23
	<b>previous</b>	50.36	43.31	44.13	43.37	42.70	43.91	43.57	43.54	42.51	41.80	41.67	39.63	37.71	37.32	
	<b>absolute change</b>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	
	<b>relative change [%]</b>	0.00	0.00	0.00	0.00	-0.01	-0.01	-0.01	0.00	0.00	0.13	0.00	0.00	0.00	0.00	

PM <sub>10</sub>	current	14.50	13.12	13.25	13.09	13.32	13.60	13.42	13.37	13.12	12.93	12.79	12.32	11.91	11.69	11.58
	previous	14.50	13.12	13.25	13.09	13.32	13.60	13.42	13.37	13.12	12.93	12.79	12.32	11.91	11.69	
	absolute change	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
	relative change [%]	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	
PM <sub>2.5</sub>	current	5.11	4.61	4.32	4.04	4.03	4.10	4.05	4.03	3.95	3.88	3.79	3.70	3.66	3.58	3.49
	previous	5.11	4.61	4.32	4.04	4.03	4.10	4.05	4.03	3.95	3.88	3.79	3.70	3.66	3.58	
	absolute change	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	relative change [%]	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	



For **pollutant-specific information on recalculated emission estimates for Base Year and 2023**, please see the pollutant specific recalculation tables following [chapter 9.1 - Recalculations](#).

## Planned improvements



At the moment, no category-specific improvements are planned.

## Uncertainty

Details are described in [chapter 1.7](#).

<sup>1)</sup>

Vos, C., Rösemann, C., Haenel, H.-D., Dämmgen, U., Döring, U., Wulf, S., Eurich-Menden, B., Döhler, H., Steuer, B., Osterburg, B., & Fuß, R. (2026) Calculations of gaseous and particulate emissions from German agriculture 1990 – 2024 : Report on methods and data (RMD) Submission 2026. [www.eminv-agriculture.de](http://www.eminv-agriculture.de)

<sup>2)</sup>

Weingarten, 1995: Das „Regionalisierte Agrar- und Umweltinformationssystem für die Bundesrepublik Deutschland“ (RAUMIS). Berichte über die Landwirtschaft Band 73, 272-302.

<sup>3)</sup>

Henrichsmeyer, W.; Cypris, Ch.; Löhe, W.; Meuth, M.; Isermeyer F; Heinrich, I.; Schefski, A.; Neander, E.; Fasterding, F.; Neumann, M.; Nieberg, H., 1996: Entwicklung des gesamtdeutschen Agrarsektormodells RAUMIS96. Endbericht zum Kooperationsprojekt. Forschungsbericht für das BMELF (94 HS 021), Bonn, Braunschweig.

<sup>4)</sup>

Statistisches Bundesamt (2020): LW20, Landwirtschaftszählung 2020. <https://www-genesis.destatis.de/datenbank/online/statistic/41141/details>

<sup>5)</sup>

NID (2026): National Inventory Report 2026 for the German Greenhouse Gas Inventory 1990-2024. Available in April 2026.

<sup>6)</sup>

EMEP/EEA air pollutant emission inventory guidebook 2023, EEA Report No 06/2023, <https://www.eea.europa.eu/en/analysis/publications/emep-eea-guidebook-2023>.

<sup>7)</sup>

Dämmgen U., Hutchings N.J. (2008): Emissions of gaseous nitrogen species from manure management - a new approach. Environmental Pollution 154, 488-497.

<sup>8)</sup>

IPCC – Intergovernmental Panel on Climate Change (2006): 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4 Agriculture, Forestry and Other Land Use.

<sup>9)</sup>

Reidy B. et al. (2008): Reidy B., Dämmgen U., Döhler H., Eurich-Menden B., Hutchings N.J., Luesink H.H., Menzi H., Misselbrook T.H., Monteny G.-J., Webb J. (2008): Comparison of models used for the calculation of national NH<sub>3</sub> emission inventories from agriculture: liquid manure systems. Atmospheric Environment 42, 3452-3467.

<sup>10)</sup>

IPCC – Intergovernmental Panel on Climate Change (2019): 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4 Agriculture, Forestry and Other Land Use.

<sup>11)</sup>

EMEP (2013): EMEP/EEA air pollutant emission inventory guidebook – 2013

<sup>12)</sup>  
Huis In't Velt, J.W.H., Dousma, F., Nijboer, G.M. (2011): Gaseous Emissions and fine dust from rabbit housing systems. Livestock research Wageningen, Report 459.