

5.C.2 - Open Burning of Waste

Category Code	Method	AD	EF
5.C.2	CS	Q	D, CS
Method(s) applied			
D	Default		
T1	Tier 1 / Simple Methodology *		
T2	Tier 2*		
T3	Tier 3 / Detailed Methodology *		
C	CORINAIR		
CS	Country Specific		
M	Model		
* as described in the EMEP/EEA Emission Inventory Guidebook - 2019, in category chapters.			
(source for) Activity Data			
NS	National Statistics		
RS	Regional Statistics		
IS	International Statistics		
PS	Plant Specific		
As	Associations, business organisations		
Q	specific Questionnaires (or surveys)		
M	Model / Modelled		
C	Confidential		
(source for) Emission Factors			
D	Default (EMEP Guidebook)		
CS	Country Specific		
PS	Plant Specific		
M	Model / Modelled		
C	Confidential		

NO _x	NM VOC	SO ₂	NH ₃	PM _{2.5}	PM ₁₀	TSP	BC	CO	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	PCDD/F	B(a)P	B(b)F	B(k)F	I(x)P	PAH1-4	HCB	PCBs																																																																																																																																																																																																																									
-/-	-/-	-/-	NE	-/-	-/-	-/-	-/-	-/-	-/-	-/-	NE	NE	NE	NE	NE	NE	NE	-/-	-/-	-/-	-/-	-/-	-/-	NE	NE																																																																																																																																																																																																																									
<table border="1"> <tr> <td>L/-</td> <td colspan="26">key source by Level only</td> </tr> <tr> <td>-/T</td> <td colspan="26">key source by Trend only</td> </tr> <tr> <td>L/T</td> <td colspan="26">key source by both Level and Trend</td> </tr> <tr> <td>-/-</td> <td colspan="26">no key source for this pollutant</td> </tr> <tr> <td>IE</td> <td colspan="26">emission of specific pollutant Included Elsewhere (i.e. in another category)</td> </tr> <tr> <td>NE</td> <td colspan="26">emission of specific pollutant Not Estimated (yet)</td> </tr> <tr> <td>NA</td> <td colspan="26">specific pollutant not emitted from this source or activity = Not Applicable</td> </tr> <tr> <td>*</td> <td colspan="26">no analysis done</td> </tr> </table>																											L/-	key source by L evel only																										-/T	key source by T rend only																										L/T	key source by both L evel and T rend																										-/-	no key source for this pollutant																										IE	emission of specific pollutant I ncluded E lsewhere (i.e. in another category)																										NE	emission of specific pollutant N ot E stimated (yet)																										NA	specific pollutant not emitted from this source or activity = N ot A pplicable																										*	no analysis done																									
L/-	key source by L evel only																																																																																																																																																																																																																																																	
-/T	key source by T rend only																																																																																																																																																																																																																																																	
L/T	key source by both L evel and T rend																																																																																																																																																																																																																																																	
-/-	no key source for this pollutant																																																																																																																																																																																																																																																	
IE	emission of specific pollutant I ncluded E lsewhere (i.e. in another category)																																																																																																																																																																																																																																																	
NE	emission of specific pollutant N ot E stimated (yet)																																																																																																																																																																																																																																																	
NA	specific pollutant not emitted from this source or activity = N ot A pplicable																																																																																																																																																																																																																																																	
*	no analysis done																																																																																																																																																																																																																																																	

Within NFR sub-category 5.C.2 - Open Burning of Waste, the German emissions inventory provides only emissions from allowed bonfires and from other wooden materials burnt outdoors. Emissions from bonfires are key source for PM_{2.5} and PM₁₀, but in principle of minor priority due to discontinuous appearance.

Please see chapter regarding farming/plantation waste: [3.F - Field burning of agricultural residues](#) - this is banned by law in Germany. So there is no gap of reporting.

Emissions from open burning of wood and green waste for traditional purposes, so-called bonfires such as Easter fires, are reported model-based. In addition to biogenic carbon dioxide, emissions of NO_x, SO₂, CO, NMVOC, particulate matter (PM_{2.5}, PM₁₀ and TSP), Polycyclic Aromatic Hydrocarbons (PAHs) and Heavy Metals are covered so far.

Method

For developing of a estimation frame a survey regarding the number of such bonfires was carried out by an expert work ¹⁾. As the result, questionnaires from municipalities and statistical projections for Germany for the year 2016 were checked. The

project has shown a declining trend since 1990. On the basis of expert judgement, a further reduction of emissions in the future is expected.

As discussed on Review 2020 regarding all relevant sources: A comparison shows that the volume of bonfires is significantly higher than the volume of campfires. In terms of number, however, the two types of fires are similar. Due to the large fluctuations of the minimum/maximum values, the median was proposed in study. In our view the estimation of bonfires emissions is conservative and completely.

Activity data

Activity data for this category are based on data from a step by step calculation: After the evaluation of the questionnaires an extrapolation of the volume and the number of bonfires was made for Germany. For the years since 2019, it became visible that, in addition to the model-based continuous decrease in activities, special aspects must be taken into account: Because of the restrictions on public activities during the pandemic, modeling of less traditional events was searched for.

Two types of fires were already classified in the expert project: camp fires in the more private sector and, most importantly, Easter Fires in the more public sector. The calculations are now considered separately and the camp fires are modeled with a continued steady decline.

Here, Easter fires follow an approach about general percentage decreases and additionally in 2019 five percentage points decrease corresponding to various cancels due to forest fire risk. In 2020 and 2021, an additional 70 percent decrease was modeled due to cancellations for pandemic response (but no complete cancellation in Germany because there were exceptions and follow-up events). For 2022 no restrictions were modelled, only the known slight decreasing trend. The following values are the result of evaluation:

Table 1: Total annual mass of bonfires, in metric tonnes [t]

1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
431,394	414,275	397,157	380,038	362,919	345,800	342,376	339,466	336,556	319,531	135,170	134,297	324,915	322,005	319,094

Emission factors

As discussed on Review 2020 regarding EF used and referenced: We use different EF from different references suitable for the burning of wooden wastes. We consider both fresh wood (garden and park waste) and dry wood (without coatings etc.). We have tried to find relevant parallels, for example because of the burning of fresh wood with regard to forest fires. But the majority of EF are derived from the EMEP/EEA Guidebook 2023 for 5.C.2, evaluated and corrected in use, as shown in the following table:


	value	unit	Current reference
CO	48.8	kg/t	EMEP/EEA Guidebook 2023, Chapter 5.C.2, table 3-2: Tier 2 emission factors for source category 5.C.2 Small-scale waste burning, forest residues
NO_x	1.38	kg/t	
SO₂	0.03	kg/t	
NMVO	1.47	kg/t	
TSP	4.31	kg/t	
PM₁₀	4.13	kg/t	
PM_{2.5}	3.76	kg/t	
BC	1,05	kg/t	EMEP/EEA Guidebook 2023, Chapter 5.C.2, table 3-2: f-BC = 28% of PM2.5
PCDD/F	10.0	µg/t	EMEP/EEA Guidebook 2023, Chapter 5.C.2, table 3-1 - Tier 1 emission factors for source category 5.C.2 Small-scale waste burning
PAH	3.39	g/t	sum of single compounds
B[a]P	1.3	g/t	IIR Ireland: EFs are referenced to a former research project called 'Use of charcoal, tobacco etc.', a literature research, which is only available via the UBA library and in German. The EFs are relating to wood burning as it was documented in Ireland's IIR. ²⁾
B[b]F	1.5	g/t	
B[k]F	0.5	g/t	
I[...]P	0.09	g/t	
Pb	0.32	g/t	EMEP/EEA Guidebook 2023, Chapter 5.C.2, table 3-2: Tier 2 emission factors for source category 5.C.2 Small-scale waste burning, forest residues
Cd	0.13	g/t	

Trends in emissions

All trends in emissions correspond to trends of AD. No emissions reduction measures are identifiable in the long term.

Trends of Emissions in Germany in NFR-category Bonfires

Emissions by pollutant / Emissionen nach Schadstoff



* Base Year for PM = 1995 / Basisjahr für Feinstäube (PM) ist 1995

Quelle: German Environment Agency, National inventory for the German reporting on atmospheric emissions since 1990, (03/2026)

Emission trends of bonfires

Recalculations



With **activity data and emission factors remaining unrevised**, no recalculations were carried out compared to the previous submission.

Planned improvements



At the moment, no category-specific improvements are planned.

¹⁾ Wagner & Steinmetzer, 2018: Jörg Wagner, Sonja Steinmetzer, INTECUS GmbH Abfallwirtschaft und umweltintegratives Management: Erhebung der Größen und Zusammensetzung von Brauchtums- und Lagerfeuern durch kommunale Befragungen; URL: https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2018-02-19_texte_11-2018_lager-brauchtumsfeuer.pdf; UBA-Texte 11/2018